
Exercise sheet 8: Donsker's invariance principle (v2)

Exercise 1 — Recurrence and Donsker.

You know that almost surely the random walk on \mathbb{Z}^2 visits 0 infinitely often. Is it the case for the bidimensional Brownian motion? What does Donsker's invariance principle tell us here?

Exercise 2 — Skorokhod's embedding.

Let X be a centered random variable with variance 1.

- (1) Argue for the existence of a sequence of random times T_k such that $(T_k - T_{k-1})_k$ is an i.i.d. sequence of mean 1 and $(S_k)_k = (B_{T_k})_k$ is a random walk whose increments are distributed like X . Define $(\tilde{S}_t^n)_t = (\frac{S_{[nt]}}{\sqrt{n}})_t$ its (properly interpolated) rescaled version.
- (2) Let $\phi_n(t) = n^{-1}T_{[nt]}$. Show that almost surely this random function converges uniformly on every compact to the identity $\mathbb{R}_+ \rightarrow \mathbb{R}_+$.
- (3) Show that $\tilde{S}^n = (t \mapsto n^{-1/2}B_{nt}) \circ \phi_n$ at points that are multiples of $1/n$. Deduce that $\|\tilde{S}^n - (t \mapsto n^{-1/2}B_{nt})\|_{[0,T]}$ goes to 0 in probability for every T .
- (4) Deduce Donsker's theorem.

Exercise 3 — Donsker's theorem for bridges.

In this exercise, let $b(n, p, k)$ denote the probability that a binomial of parameters (n, p) equals k , and f denote the standard Gaussian density. We will make use of the following *local limit theorem*, which is a refinement of the central limit theorem.

Theorem. (De Moivre–Laplace) As $n \rightarrow \infty$,

$$\sup_{k \in \mathbb{Z}} \left| \sqrt{p(1-p)n} b(n, p, k) - f\left(\frac{k - np}{\sqrt{p(1-p)n}}\right) \right| = o(n^{-1/2}).$$

Recall (a few sessions back) that the Brownian bridge β (from 0 to 0) has the following property: for every integrable function H and $\epsilon > 0$,

$$\mathbb{E} [H(\beta_{[0,1-\epsilon]})] = \mathbb{E} \left[H(B_{[0,1-\epsilon]}) \frac{\epsilon^{-1/2} f(B_{1-\epsilon})}{f(0)} \right].$$

Define the simple random walk S and its interpolated and rescaled version \tilde{S}^n . Our goal is to show that the distribution of \tilde{S}^{2n} **given that it is a bridge** (i.e. $\tilde{S}_{2n} = 0$), converges to that of β as $n \rightarrow \infty$.

- (1) Let H be a bounded continuous or positive measurable function. Fix $n \geq 1$ and $k_n \leq n - 1$. Show that

$$\mathbb{E}[H(\tilde{S}_{[0, k_n/n]}^{2n}) \mid \tilde{S}_{2n} = 0] = \mathbb{E} \left[H(\tilde{S}_{[0, k_n/n]}^{2n}) \frac{b(\frac{1}{2}, 2n - 2k_n, n - k_n + \frac{\sqrt{2n}}{2} \tilde{S}_{k/n})}{b(\frac{1}{2}, 2n, n)} \right]$$

- (2) Suppose that $k_n/n \rightarrow 1 - \epsilon$. Denote by A_n the fraction appearing in the right-hand side. Show that A_n is bounded by $b(\frac{1}{2}, 2n - 2k_n, n - k_n)/b(\frac{1}{2}, 2n, n)$, which (deterministic) bound converges to a constant. Deduce that the tightness criterion (the bound on $\mathbb{E}[|\tilde{S}_t^{2n} - \tilde{S}_s^{2n}|^4]$) that applies to \tilde{S}^{2n} still applies to the conditioned version.
- (3) Show that there exists a deterministic $o(n^{-1/2})$ such that almost surely we have $\left| A_n - \frac{\epsilon^{-1/2} f(\tilde{S}_{k/n}^{2n})}{f(0)} \right| = o(n^{-1/2})$. Deduce that all finite-dimensional marginals of \tilde{S}^{2n} given $\tilde{S}^{2n} = 0$ converge to that of the Brownian bridge.
- (4) Conclude.