

ORIENTABLE HOMOTOPY MODULES

FRÉDÉRIC DÉGLISE

CONTENTS

Introduction	1
Notations and conventions	3
1. The conjecture of Morel	3
1.1. The homotopy t -structure	3
1.2. Homotopy modules	4
1.3. Homotopy modules with transfers	6
2. Preparations	6
2.1. The theory of cycle modules	6
2.2. Modules and ring spectra	8
2.3. On the Gysin triangle and morphism	10
2.4. Coniveau spectral sequence	14
3. The proof	16
3.1. The (weak) \mathbf{H} -module structure	16
3.2. The associated cycle premodule	16
3.3. The associated cycle module	19
3.4. The remaining isomorphism	21
4. Some further comments	22
4.1. Monoidal structure	22
4.2. Weakly orientable spectra	24
4.3. Coniveau spectral sequence	24
4.4. Cohomology spectral sequence and cycle classes	25
References	27

ABSTRACT. We prove a conjecture of Morel identifying Voevodsky's homotopy invariant sheaves with transfers with spectra in the stable homotopy category which are concentrated in degree zero for the homotopy t -structure and have a trivial action of the Hopf map. This is done by relating these two kind of objects to Rost's cycle modules. Applications to algebraic cobordism and construction of cycle classes are given.

INTRODUCTION

In [Mor04] and [Mor06], F. Morel started a thorough analysis of the stable homotopy category of schemes over a field culminating in the computation of the zero-th stable homotopy groups of the zero sphere $\pi_0(S^0)_*$ in a joint work with M. Hopkins. The result involves a mixture of the Witt ring of the field and its Milnor K-theory. This paper is built on the idea of Morel that the Witt part contains the obstruction to orientation in stable homotopy. Indeed, let us recall that

Date: May 2010.

1991 Mathematics Subject Classification. 14F42, 19E15, 19D45.

Key words and phrases. Stable homotopy theory of schemes, motivic cohomology and transfers, algebraic cobordism.

Partially supported by the ANR (grant No. ANR-07-BLAN-042).

$\pi_0(S^0)_*$ is generated as an algebra by the units of the field k and the Hopf map¹ η . As in topology, the Hopf map η is the first obstruction for a ring spectrum to be oriented.²

On the other hand the units of k generate a subring of $\pi_0(S^0)_*$ which turns out to be exactly the Milnor K-theory of k , or, in contemporaneous terms, the part of motivic cohomology where the degree is equal to the twist. It is understood now, though still conjectural, that motivic cohomology is the universal oriented cohomology with additive formal group law.

To build something out of these general remarks, one has to go more deeply into motivic homotopy theory. The category $DM^{eff}(k)$ of V. Voevodsky's motivic complexes over a perfect field k is built out of the so called homotopy invariant sheaves with transfers who have the distinctive property that their cohomology is homotopy invariant.

Starting from the point that they form the heart of a natural t-structure on $DM^{eff}(k)$, Morel introduced the homotopy t-structure on the stable homotopy category $SH(k)$, analog of the previous one, and identified its heart as some graded sheaves without transfers but still with homotopy invariant cohomology. They are called *homotopy modules* by Morel (see definition 1.2.2) and we denote their category by $\Pi_*(k)$. These sheaves have to be thought as *stable homotopy groups*: in fact, taking the zero-th homology of S^0 with respect to the homotopy t-structure gives a homotopy module $\pi_0(S^0)_*$ whose fiber at the point k is precisely the graded abelian group $\pi_0(S^0)_*$. Note that, as a consequence, any homotopy module has a natural action of the Hopf map η seen as an element of $\pi_0(S^0)_*(k)$.

To relate these two kind of sheaves, it is more accurate to introduce the non effective (*i.e. stable*) version of $DM^{eff}(k)$, simply denoted by $DM(k)$. The canonical t-structure on $DM^{eff}(k)$ extends to a t-structure on $DM(k)$ whose heart $\Pi_*^{tr}(k)$ is now made by some graded homotopy invariant sheaves with transfers which we call *homotopy modules with transfers* (see definition 1.3.2).

Then the natural map $DM(k) \rightarrow SH(k)$ induces a natural functor

$$\gamma_* : \Pi_*^{tr}(k) \rightarrow \Pi_*(k)$$

which basically does nothing more than forgetting the transfers. In this article, we prove the following conjecture of Morel:

- Theorem.** (1) *The functor γ_* is fully faithful and its essential image consists of the homotopy modules on which η acts as 0.*
 (2) *The functor γ_* is monoidal and homotopy modules with transfers can be described as homotopy modules with an action of the unramified Milnor K-theory sheaf.*

Actually the first part was conjectured by Morel ([Mor04, conj. 3, p. 71]) and the second one is a remark we made, deduced from the first one.

Let us now come back to the opening point of this introduction: it turns out from this theorem that, for homotopy modules seen as objects of $SH(k)$, η is precisely the obstruction to be orientable (see corollary 4.1.4 for the precise statement). Moreover, the Milnor part of $\pi_0(S^0)_*$ – *i.e.* the unramified Milnor K-theory sheaf – is the universal homotopy module on which η acts as 0.

This theorem relies on our previous work on homotopy modules with transfers where we show that they are completely determined by the system of their fibers over finitely generated extensions of k , which can be described precisely as a cycle module in the sense of M. Rost. Actually, according to [Dég09, th. 3.4], this defines in fact an equivalence of categories: this gives a way to construct transfers on sheaves out of a more elementary algebraic structure on their fibers. The proof of our theorem thus consists in showing that the fibers of a homotopy module with trivial action of η gives a cycle module; this was actually the original form of the conjecture of Morel.

This is done by appealing to our work on the Gysin triangle [Dég08b] in the framework of modules over ring spectra: indeed, the main idea of the proof is that a homotopy module with trivial action of η has a (weak) structure of a module over the motivic cohomology ring spectrum

¹See paragraph 1.2.6.

²See remark 1.2.8.

H. After having recalled the central definitions involved in the formulation of the previous theorem in section 1, we have dedicated section 2 to recall the main technical tools which will be involved in the proof: cycle modules, modules over ring spectra, Gysin triangles and morphisms, the coniveau spectral sequence and the computation of its differentials. This enables us, in section 3, to give a neat proof of the first part of our theorem though it uses all the previous technical tools. In the last section, we prove the additional fact made by the point (2) in the above and we collect a few additional ones which illustrate our result. Notably, we use the coniveau spectral sequence to obtain the following results:

- (Cor. 4.3.4) Let **MGL** be Voevodsky’s algebraic cobordism spectrum. Then for any smooth connected scheme of dimension d , the canonical map

$$\mathbf{MGL}^{2d,d}(X) \rightarrow CH_0(X),$$

with target the 0-cycles modulo rational equivalence, is an isomorphism.

- (Prop. 4.4.5) Let \mathbb{E} be a monoid in $SH(k)$ satisfying the following assumptions:
 - (a) η acts trivially on $\pi_0(\mathbb{E})_*$.
 - (b) For any negative integers n, m , any smooth connected scheme X , and any cohomological class $\rho \in \mathbb{E}^{n,m}(X)$, there exists a non empty open $U \subset X$ such that $\rho|_U = 0$.
 Then \mathbb{E} admits an orientation whose associated formal group law is additive. Moreover, for any integer $n \geq 0$ and any smooth scheme X , there exists a canonical cycle class map

$$CH^n(X) \rightarrow \mathbb{E}^{2n,n}(X)$$

which satisfies all the usual properties.

NOTATIONS AND CONVENTIONS

In this article, k is a perfect field. Any scheme is assumed to be a *separated k -scheme* unless stated otherwise. Such a scheme will be called a smooth scheme if it is *smooth of finite type*. We denote by Sm_k (resp. $Sm_{k,\bullet}$) the category of smooth schemes (resp. pointed smooth schemes). Given a smooth scheme X , we put $X_+ = X \sqcup \text{Spec}(k)$ considered as a pointed scheme by the obvious k -point.

Following Morel, our convention for t -structures in triangulated categories is *homological* – see in particular Definition 1.1.3.

Our monoidal categories, as well as functors between them, are always assumed to be symmetric monoidal. Similarly, monoids are always assumed to be commutative.

In diagrams involving shifts or twists of morphisms, we do not indicate them in the label of the arrows – this does not lead to any confusion as they are indicated in the source and target of the arrows.

Graded (resp. bigraded) means \mathbb{Z} -graded (resp. \mathbb{Z}^2 -graded).

1. THE CONJECTURE OF MOREL

1.1. The homotopy t -structure.

1.1.1. We will denote by $SH(k)$ the *stable homotopy category over k* of Morel and Voevodsky (see [Mor04] and [Jar00]). Objects of $SH(k)$ will be simply called *spectra*.³ It is a triangulated category with a canonical functor:

$$Sm_{k,\bullet} \rightarrow SH(k), X \mapsto \Sigma^\infty X.$$

It has a closed monoidal structure; the tensor product denoted by \wedge is characterized by the property: $\Sigma^\infty(X \times_k Y)_+ = \Sigma^\infty X_+ \wedge \Sigma^\infty Y_+$. We denote by $S^0 = \Sigma^\infty \text{Spec}(k)_+$ the unit of this tensor product. By construction of $SH(k)$, the object $\Sigma^\infty \mathbb{G}_m$ is invertible for the tensor product. For any integer $n \in \mathbb{Z}$, we will denote by $S^{n,n}$ its n -th tensor power. Moreover, for any couple $(i, n) \in \mathbb{Z}^2$, we put $S^{i,n} = S^{n,n}[i - n]$.

³ They are called \mathbb{P}^1 -spectrum in [Mor04] because they have to be distinguished from S^1 -spectra. As we will not use S^1 -spectra in this work, there will be no risk of confusion.

1.1.2. Consider a spectrum \mathbb{E} . For any smooth k -scheme X and any couple $(i, n) \in \mathbb{Z}^2$, we put $\mathbb{E}^{i,n}(X) = \mathrm{Hom}_{SH(k)}(\Sigma^\infty X_+, S^{i,n} \wedge \mathbb{E})$. This defines a bigraded cohomology theory on Sm_k . We let $\pi_i(\mathbb{E})_n$ be the Nisnevich sheaf of abelian groups on Sm_k associated with the presheaf:

$$X \mapsto \mathbb{E}^{n-i,n}(X).$$

For a fixed integer $i \in \mathbb{Z}$, $\pi_i(\mathbb{E})_*$ will be considered as an abelian \mathbb{Z} -graded sheaf. Recall the definition 5.2.1 of [Mor04]:

Definition 1.1.3. A spectrum \mathbb{E} will be said to be *non negative* (resp. *non positive*) if for any $i < 0$ (resp. $i > 0$) $\pi_i(\mathbb{E})_* = 0$. We let $SH(k)_{\geq 0}$ (resp. $SH(k)_{\leq 0}$) be the full subcategory of $SH(k)$ made of non negative (resp. nonpositive) objects of $SH(k)$.

As proved in [Mor04, 5.2.3], this defines a t-structure on the triangulated category $SH(k)$ with *homological conventions*. This means the following properties:

- (1) The inclusion functor $o_+ : SH(k)_{\geq 0} \rightarrow SH(k)$ (resp. $o_- : SH(k)_{\leq 0} \rightarrow SH(k)$) admits a right adjoint t_+ (resp. left adjoint t_-). For any spectrum \mathbb{E} and any integer $i \in \mathbb{Z}$, we put:

$$\begin{aligned} E_{\geq 0} &= o_+ t_+(\mathbb{E}), & E_{\leq 0} &= o_- t_-(\mathbb{E}), \\ E_{\geq i} &= (E[-i])_{\geq 0}[i], & E_{\leq i} &= (E[-i])_{\leq 0}[i], \\ E_{> i} &= E_{\geq i+1}, & E_{< i} &= (E)_{\leq i-1}. \end{aligned}$$

- (2) For any spectra \mathbb{E}, \mathbb{F} ,

$$(1.1.3.1) \quad \mathrm{Hom}_{SH(k)}(\mathbb{E}_{> 0}, \mathbb{F}_{\leq 0}) = 0.$$

- (3) For any object \mathbb{E} in $SH(k)$, there is a unique distinguished triangle in $SH(k)$:

$$(1.1.3.2) \quad \mathbb{E}_{\geq 0} \rightarrow \mathbb{E} \rightarrow \mathbb{E}_{< 0} \rightarrow \mathbb{E}_{\geq 0}[1]$$

where the first two maps are given by the adjunctions of point (1).

Remark 1.1.4. The key point for the previous results is the stable \mathbb{A}^1 -connectivity theorem of Morel (see [Mor04, th. 4.2.10]). Recall this theorem also implies that for any smooth k -scheme X and any pair $(i, n) \in \mathbb{N} \times \mathbb{Z}$, the spectrum $S^{i+n,n} \wedge \Sigma^\infty X_+$ is non negative. Then, according to the previous definition, the spectra of this shape constitute an essentially small family of generators for the localizing subcategory $SH(k)_{\geq 0}$ of $SH(k)$.

1.2. Homotopy modules.

1.2.1. Given an abelian Nisnevich sheaf F on Sm_k , we denote by $F_{-1}(X)$ the kernel of the morphism $F(X \times_k \mathbb{G}_m) \rightarrow F(X)$ induced by the unit section of \mathbb{G}_m . Following the terminology of Morel, we will say that F is *strictly homotopy invariant* if the Nisnevich cohomology presheaf $H_{\mathrm{Nis}}^*(-, F)$ is homotopy invariant.

Definition 1.2.2 (Morel). A *homotopy module* is a pair (F_*, ϵ_*) where F_* is a \mathbb{Z} -graded abelian Nisnevich sheaf on Sm_k which is strictly homotopy invariant and $\epsilon_n : F_n \rightarrow (F_{n+1})_{-1}$ is an isomorphism. A morphism of homotopy modules is a homogeneous natural transformation of \mathbb{Z} -graded sheaves compatible with the given isomorphisms.

We denote by $\Pi_*(k)$ the category of homotopy modules.

1.2.3. For any spectrum \mathbb{E} , $\pi_0(\mathbb{E})_*$ has a canonical structure of a homotopy module. Moreover, the functor $\mathbb{E} \mapsto \pi_0(\mathbb{E})_*$ induces an equivalence of categories between the heart of $SH(k)$ for the homotopy t-structure and the category $\Pi_*(k)$ (see [Mor04, 5.2.6]). As in *loc. cit.* we will denote its quasi-inverse by:

$$(1.2.3.1) \quad H : \Pi_*(k) \rightarrow \pi_0(SH(k)).$$

Note this result implies that $\Pi_*(k)$ is a Grothendieck abelian category with generators of shape

$$(1.2.3.2) \quad \pi_0(X)_* \{n\} = \pi_0(S^{n,n} \wedge \Sigma^\infty X_+)_*$$

for a smooth k -scheme and an integer $n \in \mathbb{Z}$. It admits a monoidal structure defined by

$$(1.2.3.3) \quad F_* \otimes G_* := \pi_0(H(F_*) \wedge H(G_*))_* \simeq (H(F_*) \wedge H(G_*))_{\geq 0}.$$

The isomorphism follows from the fact that the tensor product on $SH(k)$ preserves non negative spectra (according to remark 1.1.4).

Note that $\pi_0(S^0)_*$ is the unit object for this monoidal structure. Given a homotopy module $F_* = (F_*, \epsilon_*)$ and an integer $n \in \mathbb{Z}$, we will denote by $F_*\{n\}$ the homotopy module whose i -th graded term is $(F_{i+n}, \epsilon_{i+n})$. Then:

$$(1.2.3.4) \quad F_*\{n\} = \pi_0(\mathbb{G}_m^{\wedge, n})_* \otimes F_*$$

so that this notation agrees with that of (1.2.3.2).

Remark 1.2.4. Let F_* be a homotopy module. Then according to the construction of the spectrum $\mathbb{F} := H(F_*)$, for any couple $(i, n) \in \mathbb{Z}^2$, we get an isomorphism:

$$(1.2.4.1) \quad \mathbb{F}^{i, n}(X) \simeq H_{\text{Nis}}^{i-n}(X, F_n),$$

natural in X .

Example 1.2.5. Let \mathbf{H} be the spectrum representing motivic cohomology.⁴ Then the homotopy module $\pi_0(\mathbf{H})_*$ is the sheaf of unramified Milnor K-theory \underline{K}_*^M on Sm_k .⁵

1.2.6. Define the *Hopf map* in $SH(k)$ as the morphism $\eta : \mathbb{G}_m \rightarrow S^0$ obtained by applying the tensor product with $S^{-2, -1}$ to the map induced by

$$\mathbb{A}_k^2 - \{0\} \rightarrow \mathbb{P}_k^1, (x, y) \mapsto [x, y].$$

According to [Mor04, 6.2.4],⁶ there exists a canonical exact sequence in $\Pi_*(k)$ of shape:

$$(1.2.6.1) \quad \pi_0(\mathbb{G}_m)_* \xrightarrow{\eta_*} \pi_0(S^0)_* \rightarrow \underline{K}_*^M \rightarrow 0.$$

Definition 1.2.7. A homotopy module F_* is said to be *orientable* if the map induced by η :

$$1 \otimes \eta_* : F_*\{1\} \rightarrow F_*$$

is zero. We will denote by $\Pi_*^{\eta=0}(k)$ the full subcategory of $\Pi_*(k)$ made by the orientable homotopy modules.

Note that η is in fact an element of $\pi_0(S^0)_{-1}(k)$. Given a homotopy module F_* , a smooth scheme X and an integer $n \geq 0$, $H_{\text{Nis}}^n(X, F_*)$ has a canonical structure of a graded $\pi_0(S^0)_*(k)$ -module. Then the following conditions are equivalent:

- (i) F_* is orientable.
- (ii) For any smooth scheme X and any integer $n \geq 0$, the action of η on $H_{\text{Nis}}^n(X, F_*)$ is zero.

Remark 1.2.8. (1) Consider a linear embedding $i : \mathbb{P}^1 \rightarrow \mathbb{P}^2$. Then from [Mor04, 6.2.1], the sequence

$$\Sigma^\infty(\mathbb{A}^2 - \{0\}) \xrightarrow{S^{2,1} \wedge \eta} \Sigma^\infty \mathbb{P}^1 \xrightarrow{i_*} \Sigma^\infty \mathbb{P}^2$$

is homotopy exact in $SH(k)$. In particular, one sees that a homotopy module F_* is orientable if $i^* : H_{\text{Nis}}^*(\mathbb{P}^2, F_*) \rightarrow H_{\text{Nis}}^*(\mathbb{P}^1, F_*)$ is split – actually the reciprocal statement holds (see Corollary 4.1.4).

- (2) Recall that modulo 2-torsion, if -1 is a sum of squares in k , any homotopy module is orientable (see [Mor04, 6.3.5]).

⁴ See [Voe98] or example 2.2.6.

⁵ This follows from two arguments: the identification of the unstable motivic cohomology of bidegree (n, n) of a field E with the n -th Milnor ring $K_n^M(E)$; the cancellation theorem of Voevodsky to identify unstable motivic cohomology with the stable one.

⁶ See [Mor06] for a proof (stated as Corollary 21 in the introduction, which can be deduced from the results of section 2.3).

1.3. Homotopy modules with transfers.

1.3.1. Recall Voevodsky has introduced the category Sm_k^{cor} whose objects are the smooth schemes and morphisms are the *finite correspondences*. Taking the graph of a morphism of smooth schemes induces a functor $\gamma : Sm_k \rightarrow Sm_k^{cor}$. A Nisnevich sheaf with transfers is an abelian presheaf F on Sm_k^{cor} such that $F \circ \gamma$ is a Nisnevich sheaf. Note that the construction F_{-1} of paragraph 1.2.1 applied to a sheaf with transfers F gives in fact a sheaf with transfers, still denoted by F_{-1} . In [Dég09, 1.15], we have introduced the following definition:

Definition 1.3.2. A *homotopy module with transfers* is a pair (F_*, ϵ_*) where F_* is a \mathbb{Z} -graded abelian Nisnevich sheaf with transfers which is strictly homotopy invariant and $\epsilon_n : F_n \rightarrow (F_{n+1})_{-1}$ is an isomorphism of sheaves with transfers. A morphism of homotopy modules with transfers is a homogeneous natural transformation of \mathbb{Z} -graded sheaves with transfers compatible with the given isomorphisms.

We denote by $\Pi_*^{tr}(k)$ the category of homotopy modules with transfers.

1.3.3. Let (F_*, ϵ_*) be a homotopy module with transfers. Obviously, the functor $\gamma_*(F_*) = F_* \circ \gamma$ together with the natural isomorphism $\epsilon_* \cdot \gamma$ is a homotopy module. Recall from [Dég09, 1.3] that one can attach to F_* a cohomological functor:

$$\varphi : DM_{gm}(k)^{op} \rightarrow \mathcal{A}b$$

from Voevodsky's category of geometric motives to the category of abelian groups such that:

$$\varphi(M(X)(n)[n+i]) = H_{Nis}^i(X, F_{-n}).$$

According to remark 1.2.8, the projective bundle theorem in $DM_{gm}(k)$ implies that $\gamma_*(F_*)$ is orientable. Thus we have obtained a canonical functor:

$$(1.3.3.1) \quad \gamma'_* : \Pi_*^{tr}(k) \rightarrow \Pi_*^{\eta=0}(k).$$

One can see that γ'_* is faithful. The main point of this note is the following theorem:

Theorem 1.3.4. *The functor γ'_* introduced above is an equivalence of categories.*

1.3.5. This theorem is an equivalent form of the conjecture [Mor04, conj. 3, p. 160] of Morel. Indeed, recall the main theorem of [Dég09] establishes an equivalence of categories between the category of homotopy modules with transfers and the category $\mathcal{M}Cycl(k)$ of cycle modules defined by M. Rost in [Ros96]. Indeed, from [Dég09, 3.3], we get quasi-inverse equivalences of categories:

$$(1.3.5.1) \quad \rho : \Pi_*^{tr}(k) \xleftrightarrow{\sim} \mathcal{M}Cycl(k) : A^0.$$

In fact, we will prove the previous theorem by proving the following equivalent form:

Theorem 1.3.6. *There exists a canonical functor*

$$\tilde{\rho} : \Pi_*^{\eta=0}(k) \rightarrow \mathcal{M}Cycl(k)$$

and a natural isomorphism of endofunctors of $\Pi_^{\eta=0}(k)$:*

$$\epsilon : 1 \rightarrow (\gamma'_* \circ A^0 \circ \tilde{\rho}).$$

2. PREPARATIONS

2.1. The theory of cycle modules. We shortly recall the theory of cycle modules by M. Rost (see [Ros96]) in a way that will facilitate the proof of our main result.

2.1.1. Cycle premodules.

2.1.1. A *function field* will be an extension field E of k of finite transcendental degree. A *valued function field* (E, v) will be a function field E together with a discrete valuation $v : E^\times \rightarrow \mathbb{Z}$ whose ring of integers \mathcal{O}_v is essentially of finite type over k . In this latter situation, we will denote by $\kappa(v)$ the residue field of \mathcal{O}_v .

Given a function field E , we will denote by $K_*^M(E)$ the Milnor K-theory of E . Expanding a remark of Rost ([Ros96, (1.10)]), we introduce the additive category $\tilde{\mathcal{E}}_k$ as follows: its objects are

the couples (E, n) where E is a function field and $n \in \mathbb{Z}$ an integer; the morphisms are defined by the following generators and relations:

Generators :

- (D1) $\varphi_* : (E, n) \rightarrow (L, n)$ for a field extension $\varphi : E \rightarrow L$, $n \in \mathbb{Z}$.
- (D2) $\varphi^* : (L, n) \rightarrow (E, n)$ for a finite field extension $\varphi : E \rightarrow L$, $n \in \mathbb{Z}$.
- (D3) $\gamma_x : (E, n) \rightarrow (E, n+r)$, for any $x \in K_r^M(E)$, $n \in \mathbb{Z}$.
- (D4) $\partial_v : (E, n) \rightarrow (\kappa(v), n-1)$, for any valued function field (E, v) , $n \in \mathbb{Z}$.

Relations :

- (R0) For all $x \in K_*^M(E)$, $y \in K^M(E)$, $\gamma_x \circ \gamma_y = \gamma_{x \cdot y}$.
- (R1a) $(\psi \circ \varphi)_* = \psi_* \circ \varphi_*$.
- (R1b) $(\psi \circ \varphi)^* = \varphi^* \circ \psi^*$.
- (R1c) Consider finite field extensions $\varphi : K \rightarrow E$, $\psi : K \rightarrow L$ and put $R = E \otimes_K L$. For any point $z \in \text{Spec}(R)$, let $\bar{\varphi}_z : L \rightarrow R/z$ and $\bar{\psi}_z : E \rightarrow R/z$ be the induced morphisms. Under these notations, the relation (R1c) is:

$$\psi_* \varphi^* = \sum_{z \in \text{Spec}(R)} \text{lg}(R_z) \cdot (\bar{\varphi}_z)^* (\bar{\psi}_z)_*$$
where lg stands for the length of an Artinian ring.
- (R2a) For a field extension $\varphi : E \rightarrow L$, $x \in K_*^M(E)$, $\varphi_* \circ \gamma_x = \gamma_{\varphi_*(x)} \circ \varphi_*$.
- (R2b) For a finite field extension $\varphi : E \rightarrow L$, $x \in K_*^M(E)$, $\varphi^* \circ \gamma_{\varphi_*(x)} = \gamma_x \circ \varphi^*$.
- (R2c) For a finite field extension $\varphi : E \rightarrow L$, $y \in K_*^M(L)$, $\varphi^* \circ \gamma_y \circ \varphi_* = \gamma_{\varphi^*(y)}$.
- (R3a) Let $\varphi : E \rightarrow L$ be a morphism, where (L, v) and (E, w) are valued function fields such that $v|_{E^\times} = e \cdot w$ for a positive integer e . Let $\bar{\varphi} : \kappa(w) \rightarrow \kappa(v)$ be the induced morphism. Under these notations, relation (R3a) is: $\partial_v \circ \varphi_* = e \cdot \bar{\varphi}_* \circ \partial_w$.
- (R3b) Let $\varphi : E \rightarrow L$ be a finite field extension where (E, v) is a valued function field. For any valuation w on L , we let $\bar{\varphi}_w : \kappa(v) \rightarrow \kappa(w)$ be the induced morphism. Then relation (R3b) is: $\partial_v \circ \varphi^* = \sum_{w/v} \bar{\varphi}_w^* \circ \partial_w$.
- (R3c) For any $\varphi : E \rightarrow L$, any valuation v on L , trivial on E^\times : $\partial_v \circ \varphi_* = 0$.
- (R3d) For a valued function field (E, v) , a prime π of v : $\partial_v \circ \gamma_{\{-\pi\}} \circ \varphi_* = \bar{\varphi}_*$.
- (R3e) For a valued field (E, v) and a unit $u \in E^\times$: $\partial_v \circ \gamma_{\{u\}} = -\gamma_{\{u\}} \circ \partial_v$.

Recall the following definition – [Ros96, (1.1)]:

Definition 2.1.2. A cycle premodule is an additive covariant functor $M : \tilde{\mathcal{E}}_k \rightarrow \mathcal{A}b$. We denote by $\mathcal{A}b^{\tilde{\mathcal{E}}_k}$ the category of such functors with natural transformations as morphisms.

2.1.2. *Cycle modules.*

2.1.3. Consider a cycle premodule M , a scheme X essentially of finite type over k , and an integer $p \in \mathbb{Z}$. According to *loc. cit.*, §5, we set:

$$(2.1.3.1) \quad C^p(X; M) = \bigoplus_{x \in X^p} M(\kappa_x)$$

where κ_x denotes the residue field of x in X . This is a graded abelian group and we put: $C^p(X; M)_n = \bigoplus_{x \in X^p} M_{n-p}(\kappa_x)$.⁷

Consider a couple $(x, y) \in X^{(p)} \times X^{(p+1)}$. Assume that y is a specialization of x . Let Z be the reduced closure of x in X and $\tilde{Z} \xrightarrow{f} Z$ be its normalization. Each point $t \in f^{-1}(y)$ corresponds to a discrete valuation v_t on κ_x with residue field κ_t . We denote by $\varphi_t : \kappa_y \rightarrow \kappa_t$ the morphism induced by f . Then, we define according to [Ros96, (2.1.0)] the following homogeneous morphism of graded abelian groups:

$$(2.1.3.2) \quad \partial_y^x = \sum_{t \in f^{-1}(y)} \varphi_t^* \circ \partial_{v_t} : C^p(X; M)_n \rightarrow C^{p+1}(X; M)_n$$

If y is not a specialization of x , we put: $\partial_y^x = 0$.

⁷ It is denoted by $C^p(X; M, n)$ in *loc. cit.*

Definition 2.1.4. Consider the hypothesis and notations above. We introduce the following property of the cycle premodule M :

(FD_X) For any $x \in X$ and any $\rho \in M(\kappa_x)$, $\partial_y^x(\rho) = 0$ for all but finitely many $y \in X$.

Assume **(FD_X)** is satisfied. Then for any integer p , we define according to [Ros96, (3.2)] the following morphism:

$$(2.1.4.1) \quad d_X^p = \sum_{(x,y) \in X^{(p)} \times X^{(p+1)}} \partial_y^x : C^p(X; M)_n \rightarrow C^{p+1}(X; M)_n,$$

and introduce the following further property of M :

(C_X) For any integer $p \geq 0$, $d_X^{p+1} \circ d_X^p = 0$.

2.1.5. Thus, under **(FD_X)** and **(C_X)**, the map d_X^* is a differential on $C^*(X; M)$ and we get the complex of cycles in X with coefficients in M defined by Rost in *loc. cit.* In fact, the complex $C^*(X; M)$ is graded according to (2.1.4.1).

In the conditions of the above definition, the following properties are clear from the definition:

- Let Y be a closed subscheme (resp. open subscheme, localized scheme) of X . Then **(FD_X)** implies **(FD_Y)** and **(C_X)** implies **(C_Y)**.
- Let $X = \cup_{i \in I} U_i$ be an open cover of X . Then **(FD_X)** is equivalent to **(FD_{U_i)}** for any $i \in I$.

As any affine algebraic k -scheme X can be embedded into \mathbb{A}_k^n for n sufficiently large, we deduce easily from these facts the following lemma:

Lemma 2.1.6. *Let M be a cycle premodule. Then the following conditions are equivalent:*

- (i) M satisfies **(FD_X)** for any X .
- (ii) M satisfies **(FD_{Aⁿ})** for any integer $n \geq 0$.

Assume these equivalent conditions are satisfied. Then, the following conditions are equivalent:

- (iii) M satisfies **(C_X)** for any X .
- (iv) M satisfies **(C_{Aⁿ})** for any integer $n \geq 0$.

The following definition is equivalent to that of [Ros96, (2.1)] (see in particular [Ros96, (3.3)]):

Definition 2.1.7. A cycle premodule M which satisfies the equivalent conditions (i)-(iv) of the previous lemma is called a cycle module.

2.1.8. Given a cycle module M and a scheme X essentially of finite type over k , we denote by $A^p(X; M)$ the p -th cohomology group of the complex $C^*(X; M)$. This group is graded according to the graduation of $C^*(X; M)$.

Recall that according to one of the main constructions of Rost, $A^*(X; M)$ is contravariant with respect to morphisms of smooth schemes. Moreover, according to [Dég08c], the presheaf

$$X \mapsto A^0(X; M)$$

has a canonical structure of a homotopy module with transfers (def. 1.3.2) which actually defines the functor A^0 of (1.3.5.1).

2.2. Modules and ring spectra.

2.2.1. Recall that a *ring spectrum* is a (commutative) monoid \mathbf{R} of the monoidal category $SH(k)$.

A module over the ring spectrum \mathbf{R} is a \mathbf{R} -module in $SH(k)$ in the classical sense: a spectrum \mathbb{E} equipped with a multiplication map $\gamma_{\mathbb{E}} : \mathbf{R} \wedge \mathbb{E} \rightarrow \mathbb{E}$ satisfying the usual identities – see [ML98].⁸

Given two \mathbf{R} -modules \mathbb{E} and \mathbb{F} , a morphism of \mathbf{R} -modules is a morphism $f : \mathbb{E} \rightarrow \mathbb{F}$ in $SH(k)$ such that the following diagram is commutative:

$$\begin{array}{ccc} \mathbf{R} \wedge \mathbb{E} & \xrightarrow{1_{\mathbf{R}} \wedge f} & \mathbf{R} \wedge \mathbb{F} \\ \gamma_{\mathbb{E}} \downarrow & & \downarrow \gamma_{\mathbb{F}} \\ \mathbb{E} & \xrightarrow{f} & \mathbb{F}. \end{array}$$

⁸As \mathbf{R} is commutative, we will not distinguished the left and right \mathbf{R} -modules.

This defines an additive category denoted by $\mathbf{R}\text{-mod}^w$.

Given any spectrum \mathbb{E} , $\mathbf{R} \wedge \mathbb{E}$ has an obvious structure of a \mathbf{R} -module. This defines a functor which is obviously left adjoint to the inclusion functor. Thus we get an adjunction of categories:

$$(2.2.1.1) \quad L_{\mathbf{R}}^w : SH(k) \rightleftarrows \mathbf{R}\text{-mod}^w : \mathcal{O}_{\mathbf{R}}^w.$$

Remark 2.2.2. The category $\mathbf{R}\text{-mod}^w$ has a poor structure. For example, it is not possible in general to define a tensor product or a triangulated structure on $\mathbf{R}\text{-mod}^w$. This motivates the definitions which follow.

2.2.3. Recall also from [Jar00] that $SH(k)$ is the homotopy category of a monoidal model category: the category of symmetric spectra denoted by $Sp^{\Sigma}(k)$. A *strict ring spectrum* \mathbf{R} is a commutative monoid object in the category $Sp^{\Sigma}(k)$. Given such an object we can form the category $\mathbf{R}\text{-mod}^{Sp}$ of \mathbf{R} -modules with respect to the monoidal category $Sp^{\Sigma}(k)$ and consider the natural adjunction

$$(2.2.3.1) \quad Sp^{\Sigma}(k) \rightleftarrows \mathbf{R}\text{-mod}^{Sp}$$

where the right adjoint is the forgetful functor. An object of $\mathbf{R}\text{-mod}^{Sp}$ will be called a *strict module over \mathbf{R}* .

The monoidal model category $Sp^{\Sigma}(k)$ satisfies the monoid axiom of Schwede and Shipley: this implies that the category of strict \mathbf{R} -modules admits a (symmetric) monoidal model structure such that the right adjoint of (2.2.3.1) preserves and detects fibrations and weak equivalences (cf [SS00, 4.1]).

Definition 2.2.4. We denote by $\mathbf{R}\text{-mod}$ the homotopy category associated with the model category $\mathbf{R}\text{-mod}^{Sp}$ described above.

2.2.5. Note that $\mathbf{R}\text{-mod}$ is a monoidal triangulated category. The Quillen adjunction (2.2.3.1) induces a canonical adjunction:

$$(2.2.5.1) \quad L_{\mathbf{R}} : SH(k) \rightleftarrows \mathbf{R}\text{-mod} : \mathcal{O}_{\mathbf{R}}.$$

By construction, the functor $L_{\mathbf{R}}$ is triangulated and monoidal. Given a smooth scheme X and a couple $(i, n) \in \mathbb{Z}^2$, we will put:

$$(2.2.5.2) \quad \mathbf{R}(X)(n)[i] := L_{\mathbf{R}}(S^{i,n} \wedge \Sigma^{\infty} X_+).$$

Note that the \mathbf{R} -modules of the above shape are compact and form a family of generators for the triangulated category $\mathbf{R}\text{-mod}$.

The functor $\mathcal{O}_{\mathbf{R}}$ is triangulated and conservative. Because it is the right adjoint of a monoidal functor, it is weakly monoidal. In other words, for any strict \mathbf{R} -modules M and N , we get a canonical map in $SH(k)$:

$$\mathcal{O}_{\mathbf{R}}(M \otimes_{\mathbf{R}} N) \rightarrow \mathcal{O}_{\mathbf{R}}(M) \wedge \mathcal{O}_{\mathbf{R}}(N).$$

This implies that $\mathcal{O}_{\mathbf{R}}$ maps into the subcategory $\mathbf{R}\text{-mod}^w$ giving a functor

$$(2.2.5.3) \quad \mathcal{O}_{\mathbf{R}}^l : \mathbf{R}\text{-mod} \rightarrow \mathbf{R}\text{-mod}^w.$$

For any spectrum \mathbb{E} , $\mathcal{O}_{\mathbf{R}} L_{\mathbf{R}}(\mathbb{E}) = \mathbf{R} \wedge \mathbb{E}$. Thus the following diagram commutes:

$$(2.2.5.4) \quad \begin{array}{ccc} & SH(k) & \\ L_{\mathbf{R}} \swarrow & & \searrow L_{\mathbf{R}}^w \\ \mathbf{R}\text{-mod} & \xrightarrow{\mathcal{O}_{\mathbf{R}}^l} & \mathbf{R}\text{-mod}^w \end{array}$$

Example 2.2.6. The spectrum \mathbf{H} representing motivic cohomology has a canonical structure of a strict ring spectrum. This follows from the adjunction of triangulated categories

$$(2.2.6.1) \quad \gamma^* : SH(k) \rightleftarrows DM(k) : \gamma_*$$

where $DM(k)$ is the category of stable motivic complexes over k . In fact, by the very definition,

$$\mathbf{H} = \gamma_*(\mathbb{1})$$

where $\mathbb{1}$ is the unit object for the monoidal structure on $DM(k)$. To prove that \mathbf{H} is a strict ring spectrum, the argument is that γ^* is induced by a monoidal left Quillen functor between the underlying monoidal model categories (see [CD09] for details).

Note also that any object of $DM(k)$ admits a canonical structure of a strict \mathbf{H} -module. In fact, the previous adjunction induces a canonical adjunction

$$(2.2.6.2) \quad \tilde{\gamma}^* : \mathbf{H}\text{-mod} \rightleftarrows DM(k) : \tilde{\gamma}_*$$

(see again [CD09]).

2.3. On the Gysin triangle and morphism.

2.3.1. The proof of theorem 3.2.2 requires some of the results of [Dég08b]. We recall them to the reader to make the proof more intelligible.

Recall that for any smooth scheme X , we have⁹ an isomorphism of abelian groups:

$$(2.3.1.1) \quad c_1 : \text{Pic}(X) \rightarrow \mathbf{H}^{2,1}(X).$$

Considering the left adjoint of (2.2.5.1), we get a monoidal functor:

$$(2.3.1.2) \quad SH(k) \rightarrow \mathbf{H}\text{-mod}$$

and this functor shows the category $\mathbf{H}\text{-mod}$ satisfies the axioms of paragraph 2.1 in *loc. cit.* – see example 2.12 of *loc. cit.*

Remark 2.3.2. Note that, because (2.3.1.1) is a morphism of groups, the formal group law attached to the category $\mathbf{H}\text{-mod}$ in [Dég08b, 3.7] is just the additive formal group law $F(x, y) = x + y$. In particular, the power series in the indeterminate x

$$[2]_F = F(x, x), [3]_F = F(x, F(x, x)), \dots$$

are simply given by the formula: $[r]_F = r \cdot x$ for any integer r .

2.3.3. *The Gysin triangle:* For any smooth schemes X, Z and any closed immersion $i : Z \rightarrow X$ of pure codimension c , with complementary open immersion j , there exists a canonical distinguished triangle (see [Dég08b, def. 4.6]):

$$(2.3.3.1) \quad \underline{\mathbf{H}}(Z - X) \xrightarrow{j_*} \underline{\mathbf{H}}(X) \xrightarrow{i^*} \underline{\mathbf{H}}(Z)(c)[2c] \xrightarrow{\partial_{X,Z}} \underline{\mathbf{H}}(X - Z)[1].$$

The map i^* (resp. $\partial_{X,Z}$) is called the Gysin (resp. residue) morphism associated with i . Consider moreover a commutative square of smooth schemes

$$(2.3.3.2) \quad \begin{array}{ccc} T & \xrightarrow{k} & Y \\ q \downarrow & & \downarrow p \\ Z & \xrightarrow{i} & X \end{array}$$

such that i and k are closed immersions of pure codimension c and T is the reduced scheme associated with $Z \times_X Y$. Let $h : (Y - T) \rightarrow (X - Z)$ be the morphism induced by p . Then the following formulas hold:

(G1a) Assume $T = Z \times_X Y$. Then, according to [Dég08b, prop. 4.10], the following diagram is commutative:

$$\begin{array}{ccccc} \underline{\mathbf{H}}(Y) & \xrightarrow{k^*} & \underline{\mathbf{H}}(T)(c)[2c] & \xrightarrow{\partial_{Y,T}} & \underline{\mathbf{H}}(Y - T)[1] \\ p_* \downarrow & & q_* \downarrow & & \downarrow h_* \\ \underline{\mathbf{H}}(X) & \xrightarrow{i^*} & \underline{\mathbf{H}}(Z)(c)[2c] & \xrightarrow{\partial_{X,Z}} & \underline{\mathbf{H}}(X - Z)[1]. \end{array}$$

(G1b) Assume $Z \times_X Y$ is irreducible and let e be its geometric multiplicity. Then according to [Dég08b, 4.26] combined with remark 2.3.2, the following diagram is commutative:

$$\begin{array}{ccccc} \underline{\mathbf{H}}(Y) & \xrightarrow{k^*} & \underline{\mathbf{H}}(T)(c)[2c] & \xrightarrow{\partial_{Y,T}} & \underline{\mathbf{H}}(Y - T)[1] \\ p_* \downarrow & & e \cdot q_* \downarrow & & \downarrow h_* \\ \underline{\mathbf{H}}(X) & \xrightarrow{i^*} & \underline{\mathbf{H}}(Z)(c)[2c] & \xrightarrow{\partial_{X,Z}} & \underline{\mathbf{H}}(X - Z)[1]. \end{array}$$

⁹Note that, as in example 1.2.5, one uses the cancellation theorem of Voevodsky to get this isomorphism.

Remark 2.3.4. The Gysin morphism i_* in the triangle (2.3.3.1) induces a pushout in motivic cohomology:

$$i_* : \mathbf{H}^{n,m}(Z) \rightarrow \mathbf{H}^{n+2c,m+c}(X).$$

Considering the unit element of the graded algebra $\mathbf{H}^{*,*}(Z)$, we define the fundamental class associated with i as the element $\eta_X(Z) := i_*(1) \in \mathbf{H}^{2c,c}(X)$.

Assume $c = 1$. Then Z defines an effective Cartier divisor of X which corresponds uniquely to a line bundle L over X together with a section $s : X \rightarrow L$. Moreover, s is transversal to the zero section s_0 and the following square is cartesian:

$$\begin{array}{ccc} Z & \xrightarrow{i} & X \\ i \downarrow & & \downarrow s \\ X & \xrightarrow{s_0} & L \end{array}$$

Then according to [Dég08b, 4.21], $\eta_X(Z) = c_1(L)$ with the notation of (2.3.1.1).¹⁰

2.3.5. Consider a smooth scheme X such that $X = X_1 \sqcup X_2$ and let $x_1 : X_1 \rightarrow X$ be the canonical open and closed immersion. By additivity, $\underline{\mathbf{H}}(X) = \underline{\mathbf{H}}(X_1) \oplus \underline{\mathbf{H}}(X_2)$. Moreover, x_{1*} is the obvious split inclusion which corresponds to this decomposition. Correspondingly, it follows from (G1a) that x_1^* is the obvious split epimorphism. One can complete [Dég08b] by the following lemma which describes the additivity properties of the Gysin triangle:

Lemma 2.3.6. *Consider smooth schemes X, Z and a closed immersion $\nu : Z \rightarrow X$ of pure codimension n .*

Consider the canonical decompositions $Z = \sqcup_{i \in I} Z_i$ and $X = \sqcup_{j \in J} X_j$ into connected components and put $\hat{Z}_j = Z \times_X X_j$. We also consider the obvious inclusions:

$$z_i : Z_i \rightarrow Z, x_j : X_j \rightarrow X, u_j : (X_j - \hat{Z}_j) \rightarrow (X - Z)$$

and the morphisms ν_{ji}, ∂_{ij} uniquely defined by the following commutative diagram:

$$\begin{array}{ccccc} \underline{\mathbf{H}}(X) & \xrightarrow{\nu^*} & \underline{\mathbf{H}}(Z)(n)[2n] & \xrightarrow{\partial_{X,Z}} & \underline{\mathbf{H}}(X - Z)[1] \\ \uparrow \sim & & \uparrow \sim & & \uparrow \sim \\ \oplus_{j \in J} \underline{\mathbf{H}}(X_j) & \xrightarrow{(\nu_{ji})_{j \in J, i \in I}} & \oplus_{i \in I} \underline{\mathbf{H}}(Z_i)(n)[2n] & \xrightarrow{(\partial_{ij})_{i \in I, j \in J}} & \oplus_{j \in J} \underline{\mathbf{H}}(X_j - \hat{Z}_j)[1]. \end{array}$$

Then, for any couple $(i, j) \in I \times J$:

- (1) if $Z_i \subset X_j$, we let $\nu_i^j : Z_i \rightarrow X_j$ be the obvious inclusion and we put: $Z'_i = \hat{Z}_j - Z_i$. Then $\nu_{ji} = (\nu_i^j)^*$ and $\partial_{ij} = \partial_{X_j - Z'_i, Z_i}$.
- (2) Otherwise, $\nu_{ji} = 0$ and $\partial_{ij} = 0$.

Proof. According to the preamble 2.3.5, we get: $\nu_{ji} = z_i^* \nu^* x_{j*}, \partial_{i,j} = u_j^* \partial_{X,Z} z_{i*}$.

In the respective case (1) and (2), we consider the following cartesian squares:

$$\begin{array}{ccc} \text{(1) If } Z_i \subset X_j, & & \text{(2) otherwise,} \\ \begin{array}{ccccccc} Z_i & \xrightarrow{\nu_i^j} & X_j & \xleftarrow{\hat{z}_j} & \hat{Z}_j & \longleftarrow & Z_i \\ \parallel & & \downarrow x_j & & \downarrow \nu_i^j & & \parallel \\ Z_i & \xrightarrow{\nu_i} & X & \xleftarrow{\nu} & Z & \xleftarrow{z_i} & Z_i \end{array} & & \begin{array}{ccccccc} \emptyset & \longrightarrow & X_j & \xleftarrow{\hat{z}_j} & \hat{Z}_j & \longleftarrow & \emptyset \\ \downarrow & & \downarrow x_j & & \downarrow \nu_i^j & & \downarrow \\ Z_i & \xrightarrow{\nu_i} & X & \xleftarrow{\nu} & Z & \xleftarrow{z_i} & Z_i \end{array} \end{array}$$

¹⁰ One can prove more generally that, through the isomorphism $\mathbf{H}^{2c,c}(X) \simeq CH^c(X)$ where the left hand side denotes the Chow group of codimension c cycles, i_* agrees with the usual pushout on Chow groups (see [Dég09, Prop. 3.11]). Thus $\eta_X(Z)$ simply corresponds to the cycle of Z in X .

Then the result follows from the following computations:

$$z_i^* \nu^* x_{j*} \stackrel{(a)}{=} \nu_i^* x_{j*} \stackrel{(b)}{=} \begin{cases} (\nu_i^j)^* & \text{if } Z_i \subset X_j, \\ 0 & \text{otherwise.} \end{cases}$$

$$u_j^* \partial_{X, Z} z_{i*} \stackrel{(c)}{=} \partial_{X_j, \hat{Z}_j} \hat{z}_j^* z_{i*} \stackrel{(d)}{=} \begin{cases} \partial_{X_j, \hat{Z}_j} (z_i^j)^* \stackrel{(e)}{=} \partial_{X_j - Z'_i, Z_i} & \text{if } Z_i \subset X_j, \\ 0 & \text{otherwise.} \end{cases}$$

We give the following justifications for each equality: (a) use (G2a) ($\nu_i = \nu \circ z_i$), (b) use (G1a) applied to Δ_1 , (c) use (G2b) applied to Δ_2 , (d) use (G1a) applied to Δ_3 , (e) use (G1a). \square

2.3.7. The Gysin morphism: For any smooth schemes X and Y and any projective equidimensional morphism $f : Y \rightarrow X$ of dimension n , there exists (see [Dég08b, def. 5.12]) a canonical morphism in $\mathbf{H}\text{-mod}$ of the form

$$(2.3.7.1) \quad f^* : \mathbf{H}(X)(n)[2n] \rightarrow \mathbf{H}(Y)$$

which coincides with the morphism of (2.3.3.1) in the case where f is a closed immersion. It satisfies the following properties:

- (G2a) Whenever it makes sense, we get (see [Dég08b, prop. 5.14]): $f^* g^* = (gf)^*$.
(G2b) Consider a cartesian square of smooth schemes of shape (2.3.3.2) such that i and k (resp. p and q) are closed immersion (resp. projective equidimensional morphisms). Assume i, k, f, g have constant relative codimension n, m, s, t respectively. Put $d = n + t = m + s$. Let $h : (Y - T) \rightarrow (X - Z)$ be the morphism induced by f . Then the following square is commutative (see [Dég08b, prop. 5.15]¹¹):

$$\begin{array}{ccc} \mathbf{H}(Z)(n)[2n] & \xrightarrow{\partial_{X, Z}} & \mathbf{H}(X - Z)[1] \\ g^* \downarrow & & \downarrow h^* \\ \mathbf{H}(T)(d)[2d] & \xrightarrow{\partial_{Y, T}} & \mathbf{H}(Y - T)(s)[2s + 1]. \end{array}$$

(G2c) For any cartesian square of smooth schemes

$$(2.3.7.2) \quad \begin{array}{ccc} T & \xrightarrow{q} & Y \\ g \downarrow & \Delta & \downarrow f \\ Z & \xrightarrow{p} & X \end{array}$$

such that p and q are projective equidimensional of the same dimension, we get (see [Dég08b, 5.17(i)]):

$$p^* f_* = g_* q^*.$$

(G2d) Consider a commutative square of smooth schemes of shape (2.3.7.2) such that:

- p, q, f and g are finite equidimensional morphisms.
- T is equal to the reduced scheme associated with $T' = Z \times_X Y$.

Let $(T'_i)_{i \in I}$ be the connected components of T' . For any index $i \in I$, we let $q_i : T'_i \rightarrow X'$ (resp. $g_i : T'_i \rightarrow Y$) be the morphism induced by q (resp. g) and we denote by r_i the geometric multiplicity of T'_i . Then according to [Dég08b, prop. 5.22] combined with remark 2.3.2, we get the following formula:

$$p^* f_* = \sum_{i \in I} r_i \cdot g_{i*} q_i^*.$$

Remark 2.3.8. At one point¹², we will need property (G2b) when we only require that the square (2.3.3.2) is topologically cartesian (*i.e.* $T = (Y \times_X Z)_{red}$). In fact, the proof given in [Dég08b] requires only this assumption: one proves Proposition 5.15 by reducing to Theorem 4.32 (the

¹¹There is a misprint in *loc. cit.*: one should read $n + t = m + s$, $d = n + t$.

¹²The proof of Proposition 3.3.4.

case where p and q are closed immersions), and the proof uses only the fact that the square is topologically cartesian (in the construction of the isomorphism denoted by $\mathbf{p}_{(X;Y,Y')}$).

2.3.9. Products: Let X be a smooth scheme and $\delta : X \rightarrow X \times X$ the diagonal embedding. For any morphisms $a : \underline{\mathbf{H}}(X) \rightarrow \mathbb{E}$ and $b : \underline{\mathbf{H}}(X) \rightarrow \mathbb{F}$, we denote by $a \boxtimes_X b$ the composite morphism:

$$(2.3.9.1) \quad \underline{\mathbf{H}}(X) \xrightarrow{\delta_*} \underline{\mathbf{H}}(X) \otimes_{\mathbf{H}} \underline{\mathbf{H}}(X) \xrightarrow{a \otimes_{\mathbf{H}} b} \mathbb{E} \otimes_{\mathbf{H}} \mathbb{F}.$$

(G3a) For any morphism $f : Y \rightarrow X$ of smooth schemes and any couple (a, b) of morphisms in $R\text{-mod}$ with source $\underline{\mathbf{H}}(Y)$, we get (obviously):

$$(a \boxtimes_X b) \circ f_* = (a \circ f_*) \boxtimes_Y (b \circ f_*)$$

(G3b) For any projective morphism $f : Y \rightarrow X$ of smooth schemes and any morphism a (resp. b) in $\mathbf{H}\text{-mod}$ with source $\underline{\mathbf{H}}(X)$ (resp. $\underline{\mathbf{H}}(Y)$), we get (see [Dég08b, 5.18]):

$$\begin{aligned} [a \boxtimes_X (b \circ f_*)] \circ f^* &= (a \circ f^*) \boxtimes_Y b. \\ [(b \circ f_*) \boxtimes_X a] \circ f^* &= b \boxtimes_Y (a \circ f^*). \end{aligned}$$

Remark 2.3.10. Consider a smooth scheme X and a vector bundle E/X of rank n . Let $P = \mathbb{P}(E)$ be the associated projective bundle with projection $p : P \rightarrow X$ and λ the canonical line bundle such that $\lambda \subset p^{-1}(E)$. Then, for any integer $r \geq 0$, the r -th power of the class $c = c_1(\lambda^\vee)$ (see (2.3.1.1)) in motivic cohomology corresponds to a morphism in $\mathbf{H}\text{-mod}$ which we denote by: $c^r : \underline{\mathbf{H}}(P) \rightarrow \underline{\mathbf{H}}(r)[2r]$. The projective bundle theorem ([Dég08b, 3.2]) says that the map

$$\sum_{0 \leq r \leq n} p_* \boxtimes_P c^r : \underline{\mathbf{H}}(P) \rightarrow \bigoplus_{0 \leq r \leq n} \underline{\mathbf{H}}(X)(r)[2r].$$

is an isomorphism. Thus we get a canonical map $l_n(P) : \underline{\mathbf{H}}(X)(n)[2n] \rightarrow \underline{\mathbf{H}}(P)$.

Given a finite epimorphism $f : Y \rightarrow X$ which admits a factorization as

$$Y \xrightarrow{i} P \xrightarrow{p} X$$

where i is a closed immersion and P/X is the projective bundle considered above, we recall that the Gysin morphism f^* of (2.3.7.1) is defined as the composite map:

$$\underline{\mathbf{H}}(X)(n)[2n] \xrightarrow{l_n(P)} \underline{\mathbf{H}}(P) \xrightarrow{i^*} \underline{\mathbf{H}}(Y)(n)[2n]$$

after taking the tensor product with $\mathbf{H}(-n)[-2n]$.

Assume Y (thus X) is connected and P/X has relative dimension 1. Consider the map $f_* : \mathbf{H}^{n,m}(Y) \rightarrow \mathbf{H}^{n,m}(X)$ induced by f^* . According to the above description, we get a factorization of f_* as:

$$\mathbf{H}^{n,m}(Y) \xrightarrow{i_*} \mathbf{H}^{n+2,m+1}(P) \xrightarrow{p_*} \mathbf{H}^{n,m}(X).$$

where i_* corresponds to the morphism of remark 2.3.4. To describe the second map, recall that any class $\alpha \in \mathbf{H}^{n+2,m+1}(P)$ can be written uniquely as $\alpha = p^*(\alpha_0) + p^*(\alpha_1).c_1(\lambda^\vee)$: then $p_*(\alpha) = \alpha_1$. Thus, we now deduce from this description the following *trace formula*:

$$(2.3.10.1) \quad f_*(1) = d$$

where d is the generic degree of f . In fact, according to remark 2.3.4 and its notations, $i_*(1) = c_1(L)$. Thus the formula follows from the equality $c_1(L) = d.c_1(\lambda^\vee)$ in $\text{Pic}(P)$ modulo $\text{Pic}(X)$.

2.3.11. Consider a regular invertible function $x : X \rightarrow \mathbb{G}_m$ on a smooth scheme X . According to the canonical decomposition $\underline{\mathbf{H}}(\mathbb{G}_m) = \underline{\mathbf{H}} \oplus \underline{\mathbf{H}}(1)[1]$, it induces a morphism: $x' : \underline{\mathbf{H}}(X) \rightarrow \underline{\mathbf{H}}(1)[1]$.

Using the product (2.3.9.1), we then deduce the following morphism:

$$(2.3.11.1) \quad \gamma_x = 1_{X^*} \boxtimes_X x' : \underline{\mathbf{H}}(X) \rightarrow \underline{\mathbf{H}}(X)(1)[1].$$

If $\nu_x : X \rightarrow X \times \mathbb{G}_m$ denotes the graph of x , then γ_x is also equal to the following composite map:

$$(2.3.11.2) \quad \underline{\mathbf{H}}(X) \xrightarrow{\nu_{x^*}} \underline{\mathbf{H}}(X \times \mathbb{G}_m) = \underline{\mathbf{H}}(X) \otimes_{\mathbf{H}} \underline{\mathbf{H}}(\mathbb{G}_m) \rightarrow \underline{\mathbf{H}}(X)(1)[1].$$

We will need the following properties of this particular kind of products:

Proposition 2.3.12. *Let X be a smooth scheme and $i : Z \rightarrow X$ be the immersion of a smooth divisor. Put $U = X - Z$ and let $j : U \rightarrow X$ be the canonical open immersion.*

- (1) *Let $x : X \rightarrow \mathbb{G}_m$ be a regular invertible function, \bar{u} (resp. u) its restriction to Z (resp. U). Then the following diagram is commutative:*

$$\begin{array}{ccc} \underline{\mathbf{H}}(Z)(1)[1] & \xrightarrow{\partial_{X,Z}} & \underline{\mathbf{H}}(U) \\ \gamma_{\bar{u}} \otimes_{\mathbf{H}} Id \downarrow & & \downarrow \gamma_u \\ \underline{\mathbf{H}}(Z)(2)[2] & \xrightarrow{-\partial_{X,Z} \otimes_{\mathbf{H}} Id} & \underline{\mathbf{H}}(U)(1)[1] \end{array}$$

- (2) *Consider a regular function $\pi : X \rightarrow \mathbb{A}^1$ such that $Z = \pi^{-1}(0)$. Write again $\pi : U \rightarrow \mathbb{G}_m$ for the obvious restriction of π . Then the following diagram is commutative:*

$$\begin{array}{ccccc} \underline{\mathbf{H}}(Z)(1)[1] & \xrightarrow{\partial_{X,Z}} & \underline{\mathbf{H}}(U) & \xrightarrow{\gamma_\pi} & \underline{\mathbf{H}}(U)(1)[1] \\ & \searrow i_* & & \swarrow j_* & \\ & & \underline{\mathbf{H}}(Z)(1)[1] & & \end{array}$$

Proof. (1) Let $\nu_x, \nu_u, \nu_{\bar{u}}$ be the respective graphs of x, u and \bar{u} . Applying property (G1a), we get a commutative square:

$$\begin{array}{ccc} \underline{\mathbf{H}}(Z)(1)[1] & \xrightarrow{\partial_{X,Z}} & \underline{\mathbf{H}}(U) \\ \nu_{\bar{u}*} \otimes_{\mathbf{H}} Id \downarrow & & \downarrow \nu_{u*} \\ \underline{\mathbf{H}}(Z \times \mathbb{G}_m)(1)[1] & \xrightarrow{\partial_{X \times \mathbb{G}_m, Z \times \mathbb{G}_m}} & \underline{\mathbf{H}}(U \times \mathbb{G}_m) \end{array}$$

According to [Dég08b, 4.12], we get a commutative diagram:

$$\begin{array}{ccc} \underline{\mathbf{H}}(Z \times \mathbb{G}_m)(1)[1] & \xrightarrow{\partial_{X \times \mathbb{G}_m, Z \times \mathbb{G}_m}} & \underline{\mathbf{H}}(U \times \mathbb{G}_m) \\ \parallel & & \parallel \\ \underline{\mathbf{H}}(Z) \otimes_{\mathbf{H}} \underline{\mathbf{H}}(\mathbb{G}_m)(1)[1] & \xrightarrow{\epsilon} \underline{\mathbf{H}}(Z)(1)[1] \otimes_{\mathbf{H}} \underline{\mathbf{H}}(\mathbb{G}_m) & \xrightarrow{\partial_{X,Z} \otimes Id} \underline{\mathbf{H}}(U) \otimes_{\mathbf{H}} \underline{\mathbf{H}}(\mathbb{G}_m) \end{array}$$

where ϵ is the symmetry isomorphism for the tensor product of \mathbf{H} -mod. The result then follows from the fact $\epsilon = -1$.¹³

- (2) For this point, we refer the reader to the proof of [Dég08c, 2.6.5]. \square

2.4. Coniveau spectral sequence.

2.4.1. Recall from [Dég08a, sec. 3.1.1] that a *triangulated exact couple* in a triangulated category \mathcal{T} is the data of bigraded objects D and E of \mathcal{T} and homogeneous morphisms between them

$$(2.4.1.1) \quad \begin{array}{ccc} D & \xrightarrow{(1,-1)} & D \\ & \searrow \alpha & \swarrow \\ & E & \\ & \swarrow \gamma & \searrow \beta \\ & & E \end{array} \quad \begin{array}{l} \\ \\ (-1,0) \\ (0,0) \end{array}$$

with the bidegrees of each morphism indicated in the diagram and such that for any pair of integers (p, q) :

- (1) $D_{p,q+1} = D_{p,q}[-1]$,
(2) the following sequence is a distinguished triangle:

$$D_{p-1,q+1} \xrightarrow{\alpha_{p-1,q+1}} D_{p,q} \xrightarrow{\beta_{p,q}} E_{p,q} \xrightarrow{\gamma_{p,q}} D_{p-1,q} = D_{p-1,q+1}[1].$$

¹³ Indeed it is well known that for any classes $x, y \in \mathbf{H}^{n,m}(X) \times \mathbf{H}^{s,t}(X)$, $xy = (-1)^{n+s}yx$.

We can associate with such a triangulated exact couple a *differential* according to the formula: $d = \beta \circ \gamma$. According to point (2) above, $d^2 = 0$ and we have defined a complex in \mathcal{T} :

$$\dots \rightarrow E_{p+1,q} \xrightarrow{d_{p,q}} E_{p,q} \rightarrow \dots$$

Recall also from [Dég08a, def. 3.3] that, given a smooth scheme X , a *flag* of X is a decreasing sequence $(Z^p)_{p \in \mathbb{Z}}$ of closed subschemes of X such that for all integer $p \in \mathbb{Z}$, Z^p is of codimension greater than p in X if $p \geq 0$ and $Z^p = X$ otherwise. We denote by $\mathcal{D}(X)$ the set of flags of X , ordered by termwise inclusion.

Consider such a flag $(Z^p)_{p \in \mathbb{Z}}$. Put $U_p = X - Z^p$. By hypothesis, we get an open immersion $j_p : U_{p-1} \rightarrow U_p$. In the category of pointed sheaves of sets on Sm_k , we get an exact sequence:

$$0 \rightarrow (U_{p-1})_+ \xrightarrow{j_p} (U_p)_+ \rightarrow U_p/U_{p-1} \rightarrow 0$$

because j_p is a monomorphism (in the category of schemes). Then, for any pair of integers (p, q) , we deduce from that exact sequence a distinguished triangle in $SH(k)$:

$$(2.4.1.2) \quad \begin{aligned} \Sigma^\infty(U_{p-1})_+[-p-q] &\xrightarrow{j_{p*}} \Sigma^\infty(U_p)_+[-p-q] \rightarrow \Sigma^\infty(U_p/U_{p-1})[-p-q] \\ &\rightarrow \Sigma^\infty(U_{p-1})_+[-p-q+1], \end{aligned}$$

which in turn gives a triangulated exact couple according to the above definition.

2.4.2. In the following, we will not consider the preceding exact couple for only one flag. Rather, we remark that the triangle (2.4.1.2) is natural with respect to the inclusion of flags: thus we really get a projective system of triangles and then a projective system of triangulated exact couples.

Recall that a pro-object of a category \mathcal{C} is a (*covariant*) functor F from a left filtering category \mathcal{I} to the category \mathcal{C} . Usually, we will denote such a pro-object F by the intuitive notation " $\varprojlim_{i \in \mathcal{I}} F_i$ "

and call it the *formal projective limit* of the projective system $(F_i)_{i \in \mathcal{I}}$.

For any integer $p \in \mathbb{Z}$, we introduce the following pro-objects of $SH(k)$:

$$(2.4.2.1) \quad F_p(X) = \varprojlim_{Z^* \in \mathcal{D}(X)} \Sigma^\infty(X - Z^p)_+$$

$$(2.4.2.2) \quad Gr_p(X) = \varprojlim_{Z^* \in \mathcal{D}(X)} \Sigma^\infty(X - Z^p/X - Z^{p-1})$$

Taking the formal projective limit of the triangles (2.4.1.2) where Z^* runs over $\mathcal{D}(X)$, we get a pro-distinguished triangle¹⁴:

$$(2.4.2.3) \quad \begin{aligned} F_{p-1}(X)[-p-q] &\xrightarrow{\alpha_{p-1,q+1}} F_p(X)[-p-q] \xrightarrow{\beta_{p,q}} Gr_p(X)[-p-q] \\ &\xrightarrow{\gamma_{p,q}} F_{p-1}(X)[-p-q+1], \end{aligned}$$

Definition 2.4.3. Considering the above notations, we define the *homotopy coniveau exact couple* as data for any couple of integers (p, q) of the pro-spectra:

$$D_{p,q} = F_p(X)[-p-q], \quad E_{p,q} = Gr_p(X)[-p-q]$$

and that of the homogeneous morphisms of pro-objects α, β, γ appearing in the pro-distinguished triangle (2.4.2.3).

For short, a projective system of triangulated exact couples will be called a *pro-triangulated exact couple*.

Example 2.4.4. Consider a spectrum \mathbb{E} . We extend the functor represented by \mathbb{E} to pro-spectra as follows:

$$\bar{\varphi}_{\mathbb{E}} : (\mathbb{F}_i)_{i \in \mathcal{I}} \mapsto \varprojlim_{i \in \mathcal{I}^{\circ p}} \text{Hom}_{SH(k)}(\mathbb{F}_i, \mathbb{E}).$$

¹⁴ *i.e.* the formal projective limit of a projective system of distinguished triangles.

Applying the functor $\bar{\varphi}_{\mathbb{E}}$ to the homotopy coniveau exact couple gives an exact couple of abelian groups in the classical sense (with the conventions of [McC01, th. 2.8]) whose associated spectral sequence is:

$$(2.4.4.1) \quad E_1^{p,q}(X, \mathbb{E}) = \varinjlim_{Z^* \in \mathcal{D}(X)} \mathbb{E}^{p+q}(X - Z^p/X - Z^{p-1}) \Rightarrow \mathbb{E}^{p+q}(X).$$

This is the usual coniveau spectral sequence associated with \mathbb{E} (see [BO74], [CTHK97]). Note that it is concentrated in the band $0 \leq p \leq \dim(X)$; thus it is convergent.

Remark 2.4.5. The canonical functor

$$SH(k) \rightarrow DM(k),$$

once extended to pro-objects, sends the data defined above to the motivic coniveau exact couple considered in [Dég08a, def. 3.5].

3. THE PROOF

3.1. The (weak) \mathbf{H} -module structure.

Proposition 3.1.1. *Let F_* be an orientable homotopy module.*

Then the spectrum $H(F_)$ (see (1.2.3.2)) admits a canonical structure of \mathbf{H} -module in $SH(k)$.*

Proof. This follows from the exact sequence (1.2.6.1), example 1.2.5 and the fact that the tensor product on $SH(k)$ preserves positive objects for the homotopy t-structure. \square

3.1.2. In particular, the presheaf represented by the (weak) \mathbf{H} -module $H(F_*)$ precomposed with the functor (2.2.5.3) induces a canonical functor:

$$(3.1.2.1) \quad \varphi_F : (\mathbf{H}\text{-mod})^{op} \rightarrow \mathcal{A}b.$$

According to the commutative diagram (2.2.5.4), we get a commutative diagram:

$$(3.1.2.2) \quad \begin{array}{ccc} SH(k)^{op} & \xrightarrow{\varphi_F^0} & \mathcal{A}b \\ (L_{\mathbf{H}})^{op} \downarrow & & \nearrow \varphi_F \\ (\mathbf{H}\text{-mod})^{op} & & \end{array}$$

where φ_F^0 is the presheaf represented by the spectrum $H(F_*)$. According to the isomorphism (1.2.4.1), this implies that for any smooth scheme X and any integer $n \in \mathbb{Z}$, $\varphi_F(\mathbf{H}(X)(n)[n]) = F_{-n}(X)$.

3.2. The associated cycle premodule.

3.2.1. Let \mathcal{O} be a formally smooth essentially of finite type k -algebra. A *smooth model* of \mathcal{O} will be an affine smooth scheme $X = \text{Spec}(A)$ (of finite type) such that A is a sub- k -algebra of \mathcal{O} . Let $\mathcal{M}^{sm}(\mathcal{O}/k)$ be the set of such smooth models, ordered by the relation: $\text{Spec}(B) \leq \text{Spec}(A)$ if $A \subset B$. As k is perfect, $\mathcal{M}^{sm}(\mathcal{O}/k)$ is a non empty left filtering ordered set. We will denote by (\mathcal{O}) the pro-scheme $(X)_{X \in \mathcal{M}^{sm}(\mathcal{O}/k)}$.

Theorem 3.2.2. *Consider the above notations and the category $\tilde{\mathcal{E}}_k$ introduced in 2.1.1. There exists a canonical additive functor*

$$\underline{\mathbf{H}}^{(0)} : \tilde{\mathcal{E}}_k^{op} \rightarrow \text{pro}-(\mathbf{H}\text{-mod})$$

defined on an object (E, n) of $\tilde{\mathcal{E}}_k$ by the formula:

$$\underline{\mathbf{H}}^{(0)}(E, n) = \varprojlim_{X \in \mathcal{M}^{sm}(E/k)} \mathbf{H}(X)(-n)[-n].$$

Note this theorem follows directly from our previous work on generic motives [Dég08c] when k admits resolution of singularities because the adjunction (2.2.6.2) is then an isomorphism. However, we give a proof (see paragraphs 3.2.4 and 3.2.9) which avoids this assumption. It uses the same arguments as [Dég08c] after a generalisation of its geometric constructions to the category $\mathbf{H}\text{-mod}$ (see section 2.3). But let us first state the corollary which motivates the previous result:

Corollary 3.2.3. *Let F_* be an orientable homotopy module.*

Then one associates with F_ a canonical cycle premodule $\hat{F}_* : \tilde{\mathcal{E}}_k \rightarrow \mathcal{A}b$ defined on an object (E, n) of $\tilde{\mathcal{E}}_k$ by the formula:*

$$\hat{F}_*(E, n) = \varinjlim_{X \in \mathcal{M}^{sm}(E/k)} F_{-n}(X).$$

This defines a functor: $\tilde{\rho}' : \Pi_^{\eta=0}(k) \rightarrow \mathcal{A}b^{\tilde{\mathcal{E}}_k}$.*

Proof. In fact, the functor φ_F associated with F_* in (3.1.2.1) admits an obvious extension $\bar{\varphi}_F$ to pro-objects of $\mathbf{H}\text{-mod}$. One simply puts: $\hat{F}_* = \bar{\varphi}_F \circ \mathbf{H}^{(0)}$. This is obviously functorial in F_* . \square

3.2.4. Functorialities:

(D1) is induced by the natural functoriality.

(D2): A finite extension fields $\varphi : E \rightarrow L$ induces a morphism of pro-schemes $(\varphi) : (L) \rightarrow (E)$ such that

$$(\varphi) = \text{''}\varprojlim\text{''}_{i \in I} (f_i : Y_i \rightarrow X_i)$$

where the f_i are finite surjective morphisms, whose associated generic residual extension is L/E , and the transition morphisms in the previous formal projective limit are made by transversal squares (see [Dég08c, 5.2] for details). One defines the map $\varphi_* : \mathbf{H}(E) \rightarrow \mathbf{H}(L)$ corresponding to (D2) as the formal projective limit:

$$\text{''}\varprojlim\text{''}_{i \in I} (f_i^* : \mathbf{H}(X_i) \rightarrow \mathbf{H}(Y_i))$$

using the Gysin morphism (2.3.7.1) and property (G2c).

(D3): According to example 1.2.5, for any function field E/k and any integer $n \geq 0$,

$$(3.2.4.1) \quad \left(\varinjlim_{X \in \mathcal{M}^{sm}(E/k)} \mathbf{H}^n(X) \right) \simeq K_n^M(E).$$

Thus any symbol $\sigma \in K_n^M(E)$ corresponds to a morphism of pro-objects

$$\mathbf{H}^{(0)}(E) \rightarrow \mathbf{H}(n)[n]$$

still denoted by σ . For any smooth model X of E/k , we let $\sigma_X : \mathbf{H}^{(0)}(X) \rightarrow \mathbf{H}(n)[n]$ the component of σ corresponding to X . We define γ_σ as the formal projective limit:

$$\text{''}\varprojlim\text{''}_{X \in \mathcal{M}^{sm}(E/k)} (\sigma_X \boxtimes_X 1_{X^*})$$

with the definition given by formula (2.3.9.1).

(D4): Let (E, v) be a valued function field with ring of integers \mathcal{O}_v and residue field κ_v . There exists a smooth model X of \mathcal{O}_v and a point $x \in X$ of codimension 1 corresponding to the valuation v . Let Z be the reduced closure of x in X . Given an open neighborhood U of x in X such that $Z \cap U$ is smooth, we can write the corresponding Gysin triangle (2.3.3.1) as follows:

$$(3.2.4.2) \quad \mathbf{H}(Z \cap U)(1)[1] \xrightarrow{\partial_{U, Z \cap U}} \mathbf{H}(U - Z \cap U) \xrightarrow{j_*} \mathbf{H}(U) \rightarrow \mathbf{H}(Z \cap U)(1)[2]$$

According to property (G1a), the morphism $\partial_{U, Z \cap U}$ is functorial with respect to the open subscheme U . Taking the formal projective limit of this morphism with respect to the neighborhoods U as above, we obtain the desired map:

$$\partial_v : \mathbf{H}(\kappa_v)(1)[1] \rightarrow \mathbf{H}(E).$$

Remark 3.2.5. Note for future references that the triangle (3.2.4.2) being distinguished, we get with the above notations: $j_* \circ \partial_{U, Z \cap U} = 0$.

3.2.6. Before proving the relations, we note that one can apply the preceding construction to obtain maps in motivic cohomology. More precisely, given any function fields E/k , we put (as in corollary 3.2.3):

$$\hat{\pi}_0(\mathbf{H})_n(E) := \varinjlim_{X \in \mathcal{M}^{sm}(E/k)} \mathbf{H}^n(X).$$

As $\mathbf{H}^n(X) = \mathrm{Hom}_{\mathbf{H}\text{-mod}}(\mathbf{H}(X)(-n)[-2n], \mathbf{H})$, we thus obtain that for any extension (resp. finite extension) of function fields $\varphi : K \rightarrow E$, the map (D1) (resp. (D2)) induces a canonical morphism:

$$\begin{aligned} \varphi_* : \hat{\pi}_0(\mathbf{H})_n(K) &\rightarrow \hat{\pi}_0(\mathbf{H})_n(E), \\ \text{resp. } \varphi^* : \hat{\pi}_0(\mathbf{H})_n(E) &\rightarrow \hat{\pi}_0(\mathbf{H})_n(K). \end{aligned}$$

We will need the following lemma concerning these maps:

Lemma 3.2.7. *Consider for any function field E/k the isomorphism (3.2.4.1) of graded abelian groups:*

$$K_*^M(E) \xrightarrow{\epsilon_E} \hat{\pi}_0(\mathbf{H})_*(E).$$

Then, the following properties hold:

- (1) ϵ_E is an isomorphism of graded algebras.
- (2) For any morphism $\varphi : E \rightarrow L$ of function fields, the following square is commutative:

$$\begin{array}{ccc} K_*^M(E) & \xrightarrow{\epsilon_E} & \hat{\pi}_0(\mathbf{H})_*(E) \\ \varphi_* \downarrow & & \downarrow \varphi_* \\ K_*^M(L) & \xrightarrow{\epsilon_L} & \hat{\pi}_0(\mathbf{H})_*(L) \end{array}$$

where the left (resp. right) vertical map corresponds to the standard functoriality of Milnor K -theory (resp. data (D1)).

- (3) For any finite morphism $\varphi : E \rightarrow L$ of function fields, the following square is commutative:

$$\begin{array}{ccc} K_*^M(E) & \xrightarrow{\epsilon_E} & \hat{\pi}_0(\mathbf{H})_*(E) \\ \varphi^* \uparrow & & \uparrow \varphi^* \\ K_*^M(L) & \xrightarrow{\epsilon_L} & \hat{\pi}_0(\mathbf{H})_*(L) \end{array}$$

where the left (resp. right) vertical map corresponds to the standard functoriality of Milnor K -theory (resp. data (D2)).

Proof. Assertions (1) and (2) follow precisely from [SV00, th. 3.4]. The verification of (3) is not easy due to the abstract nature of the Gysin morphism used to define (D2). However, we can follow the argument of [SV00], lemmas 3.4.1 and 3.4.4: this means we are reduced to prove the following formulas:

Lemma 3.2.8. *Let $\varphi : K \rightarrow E$ and $\psi : K \rightarrow L$ be finite extensions of fields. Let $[E : K]$ (resp. $[E : K]_i$) be the degree (resp. inseparable degree) of E/K . Then for any elements $x \in \hat{\pi}_0(\mathbf{H})_n(E)$ and $y \in \hat{\pi}_0(\mathbf{H})_n(K)$, the following formulas hold:*

- (4) $\varphi^*(x \cdot \varphi_*(y)) = \varphi^*(x) \cdot y$, $\varphi^*(\varphi_*(y) \cdot x) = y \cdot \varphi^*(x)$.
- (5) $\varphi^* \varphi_*(x) = [E : K] \cdot x$.
- (6) Put $R = E \otimes_K L$. Then, $\psi_* \varphi^*(x) = \sum_{z \in \mathrm{Spec}(R)} \mathrm{lg}(R_z) \cdot \bar{\varphi}_z^* \bar{\psi}_{z*}(x)$, with the notations of (R1c).
- (7) Assume L/K is normal. Then, $\psi_* \varphi^*(x) = [E : K]_i \cdot \sum_{j \in \mathrm{Hom}_K(E, L)} j_*(x)$.

In fact, point (7) implies point (3) for the graded part of degree 1 because according to point (2), $\psi_* : \pi_0(\mathbf{H})_1(K) \rightarrow \pi_0(\mathbf{H})_1(L)$ is injective. Then the proof of [SV00, lem. 3.4.4] allows to deduce point (3): this proof indeed uses only the preceding fact together with points (4) and (5).

Let us now prove the preceding lemma. Points (4) (resp. (6)) follow from the definition of (D2) and property (G3b) (resp. (G2d)). Point (7) is then an easy consequence of (6) using elementary Galois theory. The difficult part is point (5).

According to point (4), we are reduced to prove that $\varphi^*(1) = [E : K]$. Because of property (G2a) of the Gysin morphism, we reduce to the case where E/K is generated by a single element. In other words, $E = K[t]/(P)$ where P is a polynomial with coefficients in t . Thus we can find a smooth model X (resp. Y) of K/k (resp. E/k) and a finite surjective morphism $f : Y \rightarrow X$ whose generic residual extension is E/K and which factors as:

$$Y \xrightarrow{i} \mathbb{P}_X^1 \xrightarrow{p} X$$

where p is the canonical projection and i is a closed immersion of codimension 1. Thus point (5) now follows from the definition of (D2) and the trace formula (2.3.10.1). \square

3.2.9. Relations:

The relations follow from the preparatory work done in section 2.3, according to the following table:

(R0)	3.2.7(1)	(R1a)	Obvious.
(R1b)	(G2a)	(R1c)	(G2d)
(R2a)	(G3a)+lem. 3.2.7(2)	(R2c)	(G3b)+lem. 3.2.7(2)
(R2c)	(G3b)+lem. 3.2.7(3)	(R3a)	(G1b)
(R3b)	(G2b)	(R3c)	rem. 3.2.5
(R3d)	prop. 2.3.12(2)	(R3e)	prop. 2.3.12(1)

This concludes the proof of 3.2.2.

3.3. The associated cycle module.

3.3.1. Using the obvious extension of the functor (2.3.1.2) to pro-objects, the homotopy coniveau exact couple induces a pro-triangulated exact couple in $\text{pro}-(\mathbf{H}\text{-mod})$ whose graded terms are:

$$(3.3.1.1) \quad \begin{aligned} \underline{\mathbf{H}}(F_p(X))[-p-q] &= \varprojlim_{Z^* \in \mathcal{D}(X)} \underline{\mathbf{H}}(X - Z_p)[-p-q] \\ \underline{\mathbf{H}}(Gr_p(X))[-p-q] &= \varprojlim_{Z^* \in \mathcal{D}(X)} \underline{\mathbf{H}}(X - Z_p/X - Z_{p-1})[-p-q] \end{aligned}$$

Given a smooth closed subscheme $Z \subset X$ of pure codimension p , if we apply [Dég08b, prop. 4.3] to the closed pair (X, Z) according to paragraph 2.3.1, we get a canonical isomorphism:

$$(3.3.1.2) \quad \mathfrak{p}_{X,Z} : \underline{\mathbf{H}}(X/X - Z) \rightarrow \underline{\mathbf{H}}(Z)(p)[2p].$$

Using this isomorphism, we can easily obtain the analog of [Dég08a, 3.11] in the setting of \mathbf{H} -modules:

Proposition 3.3.2. *Consider the above notations. Let $p \in \mathbb{Z}$ be an integer and denote by $X^{(p)}$ the set of codimension p points of X .¹⁵ Then there exists a canonical isomorphism:*

$$\underline{\mathbf{H}}(Gr_p^M(X)) \xrightarrow{\epsilon_p} \prod_{x \in X^{(p)}} \underline{\mathbf{H}}(\kappa(x))(p)[2p].$$

In particular, for any point $x \in X^{(p)}$ we get a canonical projection map:

$$(3.3.2.1) \quad \pi_x : \underline{\mathbf{H}}(Gr_p(X)) \rightarrow \underline{\mathbf{H}}(\kappa(x))(p)[2p].$$

For the proof, we refer the reader to the proof of [Dég08a, 3.11] — the same proof works in our case if we use the purity isomorphism (3.3.1.2).

3.3.3. Let X be a scheme essentially of finite type over k and consider a couple $(x, y) \in X^{(p)} \times X^{(p+1)}$.

Assume that y is a specialization of x . Let Z be the reduced closure of x in X and $\tilde{Z} \xrightarrow{f} Z$ be its normalization. Each point $t \in f^{-1}(y)$ corresponds to a discrete valuation v_t on κ_x with residue

¹⁵ Conventionally, it is empty if $p < 0$ or $p > \dim(X)$.

field κ_t . We denote by $\varphi_t : \kappa_y \rightarrow \kappa_t$ the morphism induced by f . Then, we define the following morphism of pro- \mathbf{H} -modules:

$$(3.3.3.1) \quad d_y^x = \sum_{t \in f^{-1}(y)} \partial_{v_t} \circ \varphi_{t*} : \underline{\mathbf{H}}(\kappa_y)(1)[1] \rightarrow \underline{\mathbf{H}}(\kappa_x)$$

using the notations of paragraph 3.2.4. If y is not a specialization of x , we put: $\partial_y^x = 0$.

Proposition 3.3.4. *Consider the above hypothesis and notations. If X is smooth then the following diagram is commutative:*

$$\begin{array}{ccc} \underline{\mathbf{H}}(Gr_{p+1}(X)) & \xrightarrow{d_{p+1,-p-1}} & \underline{\mathbf{H}}(Gr_p^M(X))[1] \\ \pi_y \downarrow & & \downarrow \pi_x \\ \underline{\mathbf{H}}(\kappa_y)(p+1)[2p+2] & \xrightarrow{d_y^x} & \underline{\mathbf{H}}(\kappa_x)(p)[2p+1] \end{array}$$

where the vertical maps are defined in (3.3.2.1) and the map $d_{p+1,-p-1}$ in the differential associated with the pro-triangulated exact couple (3.3.1.1).

The proof is the same as the one of [Dég08a, 3.13] once we use the following correspondence table for the results used in it:

[Dég08a, 3.13]	Proposition 3.3.4
Proposition 1.36	Lemma 2.3.6
Theorem 1.34, relation (2)	(G2b)
Proposition 2.13	Remark 2.3.8
Proposition 2.9	(G2a)

3.3.5. Consider an orientable homotopy module F_* , a smooth scheme X and an integer $n \in \mathbb{Z}$. Put $\mathbb{F} = H(F_*)$ (using the functor of (1.2.3.1)).

The coniveau spectral sequence (2.4.4.1) associated with $\mathbb{F}(n)[n]$ has the following shape:

$$E_1^{p,q}(X, \mathbb{F}(n)[n]) \Rightarrow H_{\text{Nis}}^{p+q}(X, F_n),$$

where we have used the isomorphism (1.2.4.1) to identify the abutment.

Consider also the commutative diagram (3.1.2.2) and the obvious extension of $\bar{\varphi}_F^0$ (resp. $\bar{\varphi}_F$) to pro-objects:

$$\bar{\varphi}_F^0 : \text{pro-}SH(k)^{op} \rightarrow \mathcal{A}b, \text{ resp. } \bar{\varphi}_F : \text{pro-}(\mathbf{H}\text{-mod})^{op} \rightarrow \mathcal{A}b.$$

Then the previous spectral sequence is defined by the exact couple

$$(\bar{\varphi}_F^0(D_{p,q}), \bar{\varphi}_F^0(E_{p,q})) = (\bar{\varphi}_F(L_{\mathbf{H}}(D_{p,q})), \bar{\varphi}_F(L_{\mathbf{H}}(E_{p,q}))).$$

Thus, proposition 3.3.2 gives a canonical isomorphism:

$$E_1^{p,q}(X, \mathbb{F}(n)[n]) \xrightarrow{\epsilon_p^*} \begin{cases} C^p(X, \hat{F}_*)_n & \text{if } q = 0, \\ 0 & \text{otherwise,} \end{cases}$$

where \hat{F}_* is the cycle premodule associated with F_* . Consider moreover, a couple $(x, y) \in X^{(p)} \times X^{(p+1)}$. Comparing formula (2.1.3.2) with formula (3.3.3.1), proposition 3.3.4 gives the following commutative diagram:

$$\begin{array}{ccc} E_1^{p,0}(X, \mathbb{F}(n)[n]) & \xrightarrow{d_1^{p,0}} & E_1^{p+1,0}(X, \mathbb{F}(n)[n]) \\ \uparrow & & \uparrow \\ \hat{F}_*(\kappa(x)) & \xrightarrow{\partial_y^x} & \hat{F}_*(\kappa(y)) \end{array}$$

where the vertical maps are the canonical injections. In particular:

Corollary 3.3.6. *Consider the previous notations.*

- (1) *The cycle premodule \hat{F}_* satisfies properties (\mathbf{FD}_X) and (\mathbf{C}_X) .*

(2) *There is a canonical isomorphism of complexes:*

$$E_1^{*,0}(X, \mathbb{F}(n)[n]) \simeq C^*(X; \hat{F}_*)_n.$$

(3) *For any integer $p \in \mathbb{Z}$, there is a canonical isomorphism:*

$$H_{\text{Nis}}^p(X, F_n) \simeq A^p(X; \hat{F}_*)_n$$

where the right hand side is the p -th cohomology of the complex $C^*(X; \hat{F}_*)_n$ (notation of paragraph 2.1.8).

(4) *The cycle premodule \hat{F}_* is a cycle module.*

The last property follows from lemma 2.1.6 and point (1) above. It implies in particular that the functor $\tilde{\rho}'$ of corollary 3.2.3 induces a canonical functor

$$\tilde{\rho} : \Pi_*^{\eta=0}(k) \rightarrow \mathcal{M}Cycl(k)$$

establishing the first part of Theorem 1.3.6.

3.4. The remaining isomorphism. It remains to construct the natural isomorphism which appears in the statement of Theorem 1.3.6. Point (3) of the preceding corollary in the case $p = 0$ gives an isomorphism of \mathbb{Z} -graded abelian groups:

$$\epsilon_X : F_*(X) \rightarrow A^0(X; \hat{F}_*).$$

According to the definition of the functor A^0 of (1.3.5.1), the right hand side is the sections over X of the homotopy module $(\gamma_* \circ A^0 \circ \tilde{\rho})(F_*)$. We prove that ϵ_X is natural in X . Explicitly, for any morphism $f : Y \rightarrow X$ of smooth schemes, we have to prove the following diagram commutes:

$$(3.4.0.1) \quad \begin{array}{ccc} F_*(X) & \xrightarrow{\epsilon_X} & A^0(X; \hat{F}_*) \\ f^* \downarrow & & \downarrow f^* \\ F_*(Y) & \xrightarrow{\epsilon_Y} & A^0(Y; \hat{F}_*) \end{array}$$

where the vertical map on the right hand side refers to the functoriality defined by Rost.

To prove this, we can assume by additivity that X is connected, with function field E . According to our definition,

$$\hat{F}_*(E) = \varinjlim_{j_U : U \subset X} F_*(U)$$

where the colimit runs over the non empty open subschemes of X . In particular, the colimit of the morphism j_U^* induces a canonical map $\rho_X : F_*(X) \rightarrow \hat{F}_*(E)$. By definition of the coniveau spectral sequence, the isomorphism ϵ_X is induced by the exact sequence:

$$0 \rightarrow F_*(X) \xrightarrow{\rho_X} \hat{F}_*(E) \xrightarrow{d_X^0} C^1(X; \hat{F}_*)_n$$

where d_X^0 is the differential (2.1.4.1) associated with the cycle module \hat{F}_* .

(1) *The case of a flat morphism:* Consider a flat morphism $f : Y \rightarrow X$ of connected smooth schemes and let $\varphi : E \rightarrow L$ be the induced morphism on function fields. According to the definition of (D1) in paragraph 3.2.4, the following square is commutative:

$$\begin{array}{ccc} F_*(X) & \xrightarrow{\rho_X} & \hat{F}_*(E) \\ f^* \downarrow & & \downarrow \varphi^* \\ F_*(Y) & \xrightarrow{\rho_Y} & \hat{F}_*(L) \end{array}$$

Thus, the commutativity of (3.4.0.1) in this case follows from the definition of flat pullbacks on A^0 (see [Ros96, 12.2, 3.4]).

(2) *The general case:* The morphism f can be written as the composite

$$Y \xrightarrow{\gamma} Y \times X \xrightarrow{p} X$$

where γ is the graph¹⁶ of f and p is the canonical projection. To prove the commutativity of (3.4.0.1) in the case of f , we are reduced to the cases of p and γ . The case of the smooth morphism p follows from point (1). Thus we are reduced to the case where $f = i : Z \rightarrow X$ is a closed immersion between smooth schemes.

Consider an open subscheme $U \subset X$ such that the induced open immersion $j_Z : Z \cap U \rightarrow Z$ is dense. Recall that the map $j_Z^* : A^0(Z; \hat{F}_*) \rightarrow A^0(Z \cap U; \hat{F}_*)$ is injective¹⁷. Thus, according to case (1) when $f = j$ and $f = j_Z$, we are reduced to prove the commutativity of (3.4.0.1) when f is the closed immersion $Z \cap U \rightarrow U$.

In particular, we can assume that i is the composition of immersions of a smooth principal divisor. Then we are restricted to the case where Z is a smooth principal divisor.

Because Z is principal, it is parametrized by a regular function $\pi : X \rightarrow \mathbb{A}^1$. One can assume Z is connected. Then, its generic point defines a codimension 1 point of X which corresponds to a discrete valuation v on the function field E of X . We denote by $\kappa(v)$ the residue field of v . According to the computation of $i^* : A^0(X; \hat{F}_*) \rightarrow A^0(Z; \hat{F}_*)$ of [Ros96, (12.4)], the commutativity of (3.4.0.1) in the case $f = i$ is equivalent to the commutativity of the following diagram:

$$\begin{array}{ccc} F_*(X) & \xrightarrow{\rho_X} & \hat{F}_*(E) \\ i^* \downarrow & & \downarrow \partial_v \circ \gamma_\pi \\ F_*(Z) & \xrightarrow{\rho_Z} & \hat{F}_*(\kappa(v)) \end{array}$$

But this follows from point (2) of Proposition 2.3.12 and the definition of data (D3) and (D4) for the cycle module \hat{F}_* (paragraph 3.2.4).

According to the construction of the structural isomorphism

$$A^0(X; \hat{F}_*)_n \rightarrow (A^0(\cdot; \hat{F}_*)_{n+1})_{-1}(X)$$

given in [Dég09, 2.8], it is now clear that $\epsilon : F_* \rightarrow A^0(\cdot; \hat{F}_*)$ is a morphism of homotopy modules. As it is an isomorphism by construction, this concludes the proof of the main theorem 1.3.6.

4. SOME FURTHER COMMENTS

4.1. Monoidal structure.

4.1.1. Recall we have defined a t -structure on the category $DM(k)$ of motivic spectra (*cf.* [Dég09, section 5.2]) called the *homotopy t -structure* whose heart is the category $\Pi_*^{tr}(k)$ of definition 1.3.2. By the very construction, the right adjoint functor γ_* of (2.2.6.1) is t -exact with respect to the homotopy t -structures on $SH(k)$ and $DM(k)$; thus it preserves the object of the hearts. It also follows that its left adjoint γ^* preserves homologically positive objects and (2.2.6.1) induces an adjunction of abelian categories:

$$\gamma_{\leq 0}^* : \Pi_*(k) \rightleftarrows \Pi_*^{tr}(k) : \gamma_*$$

where $\gamma_{\leq 0}^* = t_- \gamma^*$.

According to this definition the functor γ_* is equal to the composite map:

$$\Pi_*^{tr}(k) \xrightarrow{\gamma'_*} \Pi_*^{\eta=0}(k) \rightarrow \Pi_*(k)$$

of the equivalence γ'_* of Theorem 1.3.4 followed by the natural inclusion. Thus, it follows from this later theorem that γ_* is fully faithful.

Recall that the functor γ^* is monoidal. According to the definition of the tensor products on $\Pi_*(k)$ and $\Pi_*^{tr}(k)$, it follows that the functor $\gamma_{\leq 0}^*$ is monoidal. Thus, its right adjoint is weakly monoidal: given any homotopy modules with transfers F_* and G_* , we get a canonical comparison map of homotopy modules:

$$\nu_{F,G} : \gamma_*(F_*) \otimes \gamma_*(G_*) \rightarrow \gamma_*(F_* \otimes^{\text{Htr}} G_*)$$

¹⁶As X/k is separated by our general assumption, γ is a closed immersion.

¹⁷This follows for example from the localization long exact sequence for Chow groups with coefficients (*cf.* [Ros96, 3.10]).

where the tensor product \otimes (resp. \otimes^{Htr}) refers to the natural tensor product on $\Pi_*(k)$ (resp. $\Pi_*^{\text{tr}}(k)$) defined in (1.2.3.3) (resp. [Dég09, 1.17]).

The following result is a corollary of Theorem 1.3.4:

Corollary 4.1.2. *For any homotopy modules with transfers F_* and G_* , the morphism $\nu_{F,G}$ introduced above is an isomorphism.*

Proof. Because γ_* is fully faithful, we can write $F_* = \gamma_{\leq 0}^*(F'_*)$ (resp. $G_* = \gamma_{\leq 0}^*(G'_*)$) with $F'_* = \gamma_*(F_*)$ (resp. $G'_* = \gamma_*(G_*)$). Then we consider the following commutative diagram:

$$\begin{array}{ccc} \gamma_*\gamma_{\leq 0}^*(F'_*) \otimes \gamma_*\gamma_{\leq 0}^*(G'_*) & \xrightarrow{\nu_{F,G}} & \gamma_*(\gamma_{\leq 0}^*(F'_*) \otimes^{\text{Htr}} \gamma_{\leq 0}^*(G'_*)) \xrightarrow{\sim} \gamma_*\gamma_{\leq 0}^*(F'_* \otimes G'_*) \\ \downarrow a \otimes a' & & \downarrow b \\ F'_* \otimes G'_* & \xlongequal{\hspace{10em}} & F'_* \otimes G'_* \end{array}$$

where a (resp. a' , b) stands for the natural transformation $\alpha : 1 \rightarrow \gamma_*\gamma_{\leq 0}^*$ evaluated at F'_* (resp. G'_* , $F'_* \otimes G'_*$). Applying again the fact γ_* is fully faithful, α_X is an isomorphism whenever X belongs to the image of γ_* , which is precisely $\Pi_*^{\eta=0}(k)$. Thus a and a' are isomorphisms. In order, to conclude it is sufficient to prove that b is an isomorphism. This amounts to show that the homotopy module $F_* \otimes G_*$ is orientable, which is obvious from definition 1.2.7. \square

4.1.3. Recall that \underline{K}_*^M , equipped with its canonical structure of a graded sheaf with transfers, is the unit object of the monoidal category $\Pi_*^{\text{tr}}(k)$ (cf [Dég09, 1.15, 3.8]). As an object of $\Pi_*(k)$, it corresponds to $\pi_0(\mathbf{H})_*$ (example 1.2.5). Note that it is a commutative monoid in the monoidal category $\Pi_*(k)$ whose multiplication (resp. unit) map is equal to

$$\begin{aligned} \mu &= \pi_0(\mu_{\mathbf{H}}) : \underline{K}_*^M \otimes \underline{K}_*^M \rightarrow \underline{K}_*^M \\ \text{resp. } u &= \pi_0(u_{\mathbf{H}}) : \pi_0(S^0)_* \rightarrow \underline{K}_*^M \end{aligned}$$

where $\mu_{\mathbf{H}}$ (resp. $u_{\mathbf{H}}$) is the multiplication (resp. unit) of the ringed spectrum \mathbf{H} . In particular, we can consider the category $\underline{K}_*^M\text{-mod}$ of modules over \underline{K}_*^M in the monoidal category $\Pi_*(k)$: it is naturally a Grothendieck abelian monoidal category. As a corollary of the previous result we get the following nice characterization of orientability for homotopy modules:

Corollary 4.1.4. *Consider the above notations.*

- (1) *The multiplication map μ is an isomorphism.*
- (2) *Given a homotopy module F_* , the following conditions are equivalent:*
 - (i) *F_* is an orientable homotopy module (definition 1.2.7).*
 - (ii) *F_* as a Nisnevich sheaf admits a canonical structure of a sheaf with transfers.*
 - (iii) *The map $u \otimes 1_{F_*} : F_* \rightarrow \underline{K}_*^M \otimes F_*$ is an isomorphism.*
 - (iv) *F_* admits a structure of a \underline{K}_*^M -module in $\Pi_*(k)$.*
 - (v) *The spectrum $H(F_*)$ has a structure of a (strict) \mathbf{H} -module.*
 - (vi) *The spectrum $H(F_*)$ is orientable.*
- (3) *The functor $\gamma_{\leq 0}^* : \Pi_*(k) \rightarrow \Pi_*^{\text{tr}}(k)$ can be factorized as:*

$$\Pi_*(k) \xrightarrow{\underline{K}_*^M \otimes ?} \underline{K}_*^M\text{-mod} \xrightarrow{\epsilon} \Pi_*^{\text{tr}}(k)$$

and ϵ is an equivalence of abelian monoidal categories.

Proof. Point (1) follows from Corollary 4.1.2 applied in the case $F = G = \underline{K}_*^M$. The assertion of point (2) follows from our previous results: (i) \Rightarrow (ii): 1.3.4, (ii) \Rightarrow (iii): Point (1) follows from Corollary, (iii) \Rightarrow (iv): obvious, (iv) \Rightarrow (v): because $\underline{K}_*^M = \pi_0(\mathbf{H})_*$ (see also the proof of 3.1.1), (v) \Rightarrow (vi): using the morphism of ringed spectra $\mathbf{MGL} \rightarrow \mathbf{H}$ corresponding to the orientation of \mathbf{H} , (vi) \Rightarrow (i): see remark 1.2.8. According to point (1), $\underline{K}_*^M\text{-mod}$ is a full subcategory of $\Pi_*(k)$; thus point (2) implies point (3). \square

Remark 4.1.5. It is not true in general that the multiplication map $\mu : \mathbf{H} \wedge \mathbf{H} \rightarrow \mathbf{H}$ is an isomorphism according to the non triviality of the Steenrod operations.¹⁸

4.2. Weakly orientable spectra.

4.2.1. Consider a spectrum \mathbb{E} . The Hopf map η obviously acts on the \mathbb{E} -cohomology of a smooth scheme as:

$$\gamma_\eta : \mathbb{E}^{n,p}(X) \rightarrow \mathbb{E}^{n-1,p-1}(X), \rho \mapsto \eta \wedge \rho.$$

The following lemma is obvious:

Lemma 4.2.2. *Given a spectrum \mathbb{E} , the following conditions are equivalent:*

- (i) *For any integer $n \in \mathbb{Z}$, $\pi_n(\mathbb{E})_*$ is an orientable.*
- (ii) *The map $\eta \wedge 1_{\mathbb{E}}$ is zero.*
- (ii') *The map $\underline{\mathrm{Hom}}(\eta, \mathbb{E})$ is zero.*
- (iii) *The operation γ_η on \mathbb{E} -cohomology defined above is zero.*

Definition 4.2.3. A spectrum \mathbb{E} satisfying the equivalent conditions of the preceding lemma will be said to be *weakly orientable*.

Remark 4.2.4. From the second point of remark 1.2.8, if 2 is invertible in E^{**} and (-1) is a sum of squares in k , then \mathbb{E} is weakly orientable.

As a corollary of our main theorem 1.3.4, we get the following way to construct cycle modules:

Proposition 4.2.5. *Let \mathbb{E} be a weakly orientable spectrum.*

Then for any integer $i \in \mathbb{Z}$, there exists a canonical cycle modules whose values on a couple (L, n) of $\tilde{\mathcal{E}}_k$ is the abelian group:

$$\mathbb{E}^{i+n,n}(L) = \varinjlim_{X \in \mathcal{M}^{sm}(E/k)^{op}} \mathbb{E}^{i+n,n}(X),$$

with the notations of paragraph 3.2.1.

The new example here is given by the case where $\mathbb{E} = \mathbf{MGL}$ is Voevodsky's algebraic cobordism spectrum.

4.3. Coniveau spectral sequence. As a corollary of the detailed analysis of Proposition 3.3.4, we get the following statement:

Proposition 4.3.1. *Let \mathbb{E} be a spectrum and $q \in \mathbb{Z}$ be an integer such that the homotopy module $\pi_{-q}(\mathbb{E})$ is orientable.*

Then for any smooth scheme X , there exists a canonical isomorphism of complexes of abelian groups:

$$E_1^{*,q}(X, \mathbb{E}) \simeq C^*(X, \hat{\pi}_{-q}(\mathbb{E})_*)_0$$

where the left hand sides refers to the q -th line of the first page of the coniveau spectral sequence associated with \mathbb{E} — cf. example 2.4.4.

Proof. We can obviously assume $q = 0$. The complex $E_1^{*,0}(X, \mathbb{E})$ is natural in \mathbb{E} . Moreover, for any spectrum \mathbb{F} , one checks using the purity isomorphism in $SH(k)$ ([MV99, §4, 2.23]) and the same argument as in the proof of Proposition 3.3.2 that the induced maps:

$$E_1^{*,0}(X, \mathbb{F}_{\geq 0}) \rightarrow E_1^{*,0}(X, \mathbb{F}), \quad E_1^{*,0}(X, \mathbb{F}) \rightarrow E_1^{*,0}(X, \mathbb{F}_{\leq 0})$$

are isomorphisms. This yields an isomorphism of complexes:

$$E_1^{*,0}(X, \mathbb{E}) \simeq E_1^{*,0}(X, \pi_0(\mathbb{E})).$$

so that we are reduced to the case where E is an orientable homotopy module. This case is precisely Corollary 3.3.6. \square

¹⁸ However, we prove in [CD09, 13.1.6] that $\mu \otimes_{\mathbb{Z}} \mathbb{Q}$ is an isomorphism (in $SH(k)_{\mathbb{Q}}$).

Remark 4.3.2. The previous corollary can be applied to any weakly orientable spectrum \mathbb{E} : we get for any smooth scheme X and any integer $n \in \mathbb{Z}$ a convergent spectral sequence of the form:

$$(4.3.2.1) \quad E_2^{p,q}(X, \mathbb{E}(n)) = A^p(X, \hat{\pi}_{q-n}(\mathbb{E})_*) \Rightarrow \mathbb{E}^{p+q,n}(X).$$

4.3.3. Let \mathbf{MGL} be Voevodsky's algebraic cobordism spectrum. According to a theorem of Morel, there exists a canonical isomorphism of homotopy modules: $\pi_0(\mathbf{MGL})_* \simeq K_*^M$.¹⁹ One deduces from this result that the ring morphism $\varphi : \mathbf{MGL} \rightarrow \mathbf{H}$ corresponding to the orientation of \mathbf{H} given by (2.3.1.1) induces an isomorphism:

$$\pi_0(\mathbf{MGL})_* \rightarrow \pi_0(\mathbf{H})_*.$$

Moreover, given a smooth connected scheme X and an integer $d \in \mathbb{Z}$, the morphism φ induces a morphism of the spectral sequences of type (4.3.2.1) which, on the E_2 -term is equal to:

$$A^p(X, \hat{\pi}_{q-d}(\mathbf{MGL})_*) \rightarrow A^p(X, \hat{\pi}_{q-d}(\mathbf{H})_*).$$

In case d is the dimension of X , the terms $E_2^{p,q}$ of the respective spectral sequences for \mathbf{MGL} and \mathbf{H} are concentrated in the region $0 \leq p \leq d, q \leq d$. Thus we get the following:

Corollary 4.3.4. *Let X be a smooth scheme of pure dimension d . Then the morphism φ induces an isomorphism:*

$$\mathbf{MGL}^{2d,d}(X) \xrightarrow{\varphi_*} \mathbf{H}^{2d,d}(X) = A^d(X, K_*^M)_d = CH^d(X).$$

Obviously, this isomorphism is natural with respect to pullbacks. It is also compatible with pushouts induced by the Gysin morphism of a projective morphism $f : Y \rightarrow X$ of smooth connected schemes.

Remark 4.3.5. If X is projective smooth, the corollary can be reformulated using duality (see [Dég08b, th. 5.23]) by saying that φ induces an isomorphism in homology:

$$\mathbf{MGL}_0(X) \xrightarrow{\sim} CH_0(X).$$

If X is only smooth, under the assumption of resolution of singularities, one can replace the left hand side in the previous isomorphism by the Borel Moore algebraic cobordism:

$$\mathbf{MGL}_0^c(X) \xrightarrow{\sim} CH_0(X).$$

4.4. Cohomology spectral sequence and cycle classes.

4.4.1. Consider a spectrum \mathbb{E} . The truncation functor for the homotopy t -structure gives a canonical (functorial) tower in $SH(k)(k)$:

$$\dots \rightarrow \mathbb{E}_{\geq p} \rightarrow \mathbb{E}_{\geq p+1} \rightarrow \dots$$

called the *Postnikov tower* of \mathbb{E} .

Then for any smooth scheme X and any integer $n \in \mathbb{Z}$, the Postnikov tower of $\mathbb{E}(n)$ gives the following spectral sequence:

$$(4.4.1.1) \quad E_{2,t}^{p,q}(X, \mathbb{E}) = H_{\text{Nis}}^q(X, \pi_{n-p}(\mathbb{E})_*) \Rightarrow \mathbb{E}^{p+q,n}(X)$$

which we simply call the *cohomology spectral sequence*.

Note that, when \mathbb{E} is weakly orientable, Corollary 3.3.6 gives a canonical isomorphism between the E_2 -term of the coniveau spectral sequence (4.3.2.1) and the E_2 -term of the cohomology spectral sequence.

Remark 4.4.2. Using the same argument as in the proof of [Dég09, th. 6.4] and a construction of Gillet and Soulé²⁰, one can show that this isomorphism is compatible with the differential on each E_2 -term and that they induces an isomorphism of spectral sequences. However, in the remainings, we will only need to be able to compute the E_2 -term using the isomorphism of Corollary 3.3.6.

¹⁹Indeed, this follows from the exact sequence (1.2.6.1).

²⁰More precisely, one replaces the use of the shifted filtration of Deligne on complexes by its generalization for spectra given in [GS99].

4.4.3. The advantage of the spectral sequence (4.4.1.1) compared to the coniveau spectral sequence is that it is obviously functorial in X (contravariantly).

Assume that \mathbb{E} has the structure of a weak \mathbf{MGL} -module. Consider smooth schemes X, Y and a projective morphism $f : Y \rightarrow X$ equidimensional of dimension n . Then according to the construction of [Dég08b, def. 5.12], we get a canonical morphism of \mathbf{MGL} -modules:

$$f^* : \mathbf{MGL}(X)(n)[2n] \rightarrow \mathbf{MGL}(Y).$$

Using diagram (2.2.5.4) in the case $R = \mathbf{MGL}$, this map induces after applying the functor $\mathbb{E}^{i,j}$ a canonical pushout:

$$(4.4.3.1) \quad \mathbb{E}^{i,j}(Y) \xrightarrow{f^*} \mathbb{E}^{i-2n,j-n}(X).$$

Lemma 4.4.4. *Consider the notations above. Then the Gysin map f^* induces a morphism of spectral sequences:*

$$E_{2,t}^{p,q}(\mathbf{MGL}(Y), \mathbb{E}) = H_{\text{Nis}}^q(Y, \pi_{-p}(\mathbb{E})_0) \rightarrow H_{\text{Nis}}^{q-n}(Y, \pi_{-p}(\mathbb{E})_{-n}) = E_{2,t}^{p,q}(\mathbf{MGL}(X)(n)[2n], \mathbb{E})$$

which converges to the morphism (4.4.3.1).

Using the cohomology spectral sequence, we get the following proposition which gives mild conditions on a spectrum \mathbb{E} for the existence of cycle classes in \mathbb{E} -cohomology satisfying the usual properties:

Proposition 4.4.5. *Let \mathbb{E} be a (weak) ring spectrum such that:*

- (a) *The homotopy module $\pi_0(\mathbb{E})_*$ is orientable.*
- (b) *For any function field K/k , $\mathbb{E}^{n,m}(K) = 0$ if $n < 0$ and $m < 0$.*

Then the following conditions hold:

- (1) *The spectrum \mathbb{E} admits an orientation whose associated formal group law is additive.*
- (2) *For any smooth scheme X and any integer $n \geq 0$, there exists a canonical morphism of abelian groups:*

$$\sigma_X : CH^n(X) \rightarrow \mathbb{E}^{2n,n}(X)$$

which is natural with respect to pullbacks, projective pushforwards and compatible with products.

Proof. According to assumption (a) and Corollary 4.1.4, the unit map $S^0 \rightarrow \mathbb{E}$ induces a canonical map

$$\underline{K}_*^M = \underline{K}_*^M \otimes \pi_0(S^0)_* \rightarrow \underline{K}_*^M \otimes \pi_0(\mathbb{E})_* = \pi_0(\mathbb{E})_*$$

which is a morphism of monoids in $\Pi_*^{\neq 0}(k)$. In particular, according to Corollary 3.3.6, we get for any smooth scheme X and any integer $n \geq 0$ a canonical morphism:

$$(4.4.5.1) \quad CH^n(X) = A^n(X, K_*)_n \rightarrow A^n(X, \hat{\pi}_0(\mathbb{E})_*)_n = H_{\text{Nis}}^n(X, \pi_0(\mathbb{E})_n)$$

compatible with pullbacks, pushouts and products.

Applying again Corollary 3.3.6, assertion (b) implies that the term $E_{2,t}^{p,q}(X, \mathbb{E}(n))$ of the cohomology spectral sequence is zero if $p > \min(q, n)$. Thus, we get a canonical composite map:

$$(4.4.5.2) \quad E_{2,t}^{n,n}(X, \mathbb{E}(n)) \xrightarrow{(1)} E_{\infty,t}^{n,n}(X, \mathbb{E}(n)) \xrightarrow{(2)} \mathbb{E}^{2n,n}(X)$$

where the map (1) is obtain as the sequence of epimorphism deduced from the spectral sequence (4.4.1.1) and the map (2) is the edge morphism (which is a monomorphism). This composite map is compatible with products and pushouts according to paragraph 4.4.3. The fact it is compatible with products follows from the construction of products on the spectral sequence of the type (4.4.1.1) (see [McC01]).

The composition of (4.4.5.1) and (4.4.5.2) gives the map of the point (2). In the case $n = 1$, we get an orientation of \mathbb{E} whose associated formal group law is additive because σ_X is a morphism of groups. \square

Remark 4.4.6. According to a result announced by Morel and Voevodsky, at least over a field of characteristic 0, the map induced by the morphism φ of paragraph 4.3.3 induces an isomorphism of ring spectra

$$\mathbf{MGL}/\{a_{ij}, (i, j) \in \mathbb{N}^2\} \rightarrow \mathbf{H}$$

where $a_{ij} : S^{2(i+j), i+j} \rightarrow \mathbf{MGL}$ denotes the coefficients of the formal group law of \mathbf{MGL} equipped with its obvious orientation.

The orientation of the first point of the previous proposition corresponds to a morphism of ring spectra

$$\psi : \mathbf{MGL} \rightarrow \mathbb{E}$$

such that $\psi \circ a_{ij}$ is zero. In particular, ψ induces a canonical morphism of ring spectra:

$$\sigma : \mathbf{H} \rightarrow \mathbb{E}.$$

which gives back the cycle class of point (2).

REFERENCES

- [BO74] S. Bloch and A. Ogus, *Gersten's conjecture and the homology of schemes*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 181–201 (1975). MR MR0412191 (54 #318)
- [CD09] D.-C. Cisinski and F. Déglise, *Triangulated categories of mixed motives*, arXiv:0912.2110, 2009.
- [CTHK97] J.-L. Colliot-Thélène, R. T. Hoobler, and B. Kahn, *The Bloch-Ogus-Gabber theorem*, Algebraic K-theory (Toronto, ON, 1996), Fields Inst. Commun., vol. 16, Amer. Math. Soc., Providence, RI, 1997, pp. 31–94. MR MR1466971 (98j:14021)
- [Dég08a] F. Déglise, *Around the gysin triangle I*, arXiv:0804.2415v1, 2008.
- [Dég08b] ———, *Around the gysin triangle II*, Doc. Math. **13** (2008), 613–675.
- [Dég08c] ———, *Motifs génériques*, Rend. Semin. Mat. Univ. Padova **119** (2008), 173–244.
- [Dég09] ———, *Modules homotopiques*, arXiv:0904.4747v1, 2009.
- [GS99] H. Gillet and C. Soulé, *Filtrations on higher algebraic K-theory*, Algebraic K-theory (Seattle, WA, 1997), Proc. Sympos. Pure Math., vol. 67, Amer. Math. Soc., Providence, RI, 1999, pp. 89–148. MR MR1743238 (2001i:19005)
- [Jar00] J. F. Jardine, *Motivic symmetric spectra*, Doc. Math. **5** (2000), 445–553 (electronic).
- [McC01] J. McCleary, *A user's guide to spectral sequences*, second ed., Cambridge Studies in Advanced Mathematics, vol. 58, Cambridge University Press, Cambridge, 2001. MR MR1793722 (2002c:55027)
- [ML98] Saunders Mac Lane, *Categories for the working mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998. MR MR1712872 (2001j:18001)
- [Mor04] F. Morel, *An introduction to \mathbb{A}^1 -homotopy theory*, Contemporary developments in algebraic K-theory, ICTP Lect. Notes, XV, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004, pp. 357–441 (electronic). MR MR2175638 (2006m:19007)
- [Mor06] ———, *\mathbb{A}^1 -algebraic topology over a field.*, <http://www.mathematik.uni-muenchen.de/~morel/preprint.html>, 2006.
- [MV99] F. Morel and V. Voevodsky, *\mathbb{A}^1 -homotopy theory of schemes*, Inst. Hautes Études Sci. Publ. Math. **90** (1999), 45–143 (2001).
- [Ros96] M. Rost, *Chow groups with coefficients*, Doc. Math. J. (1996), 319–393.
- [SS00] S. Schwede and B.E. Shipley, *Algebras and modules in monoidal model categories*, Proc. London Math. Soc. (3) **80** (2000), no. 2, 491–511.
- [SV00] Andrei Suslin and Vladimir Voevodsky, *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, The arithmetic and geometry of algebraic cycles (Banff, AB, 1998), NATO Sci. Ser. C Math. Phys. Sci., vol. 548, Kluwer Acad. Publ., Dordrecht, 2000, pp. 117–189. MR MR1744945 (2001g:14031)
- [Voe98] V. Voevodsky, *\mathbb{A}^1 -homotopy theory*, Proceedings of the International Congress of Mathematicians, Vol. I (Berlin, 1998), no. Extra Vol. I, 1998, pp. 579–604 (electronic).

LAGA, CNRS (UMR 7539), UNIVERSITÉ PARIS 13, AVENUE JEAN-BAPTISTE CLÉMENT, 93430 VILLETANEUSE, FRANCE

E-mail address: deglise@math.univ-paris13.fr

URL: <http://www.math.univ-paris13.fr/~deglise/>