

Observation of negative line tensions from Plateau border regions in dry foam films

J.-C. GÉMINARD[†]§, A. ŻYWOCIŃSKI[‡], F. CAILLIER[†] and P. OSWALD[†]

[†]Laboratoire de Physique, Unité Mixte de Recherche associée
au CNRS 5672, École Normale Supérieure de Lyon, 46 Allée d'Italie,
69364 Lyon Cedex 07, France

[‡]Institute of Physical Chemistry, Polish Academy of Sciences, PL-01-224
Warsaw 42, Poland

[Received in final form 10 October 2003 and accepted 30 October 2003]

ABSTRACT

We measured the angles produced in the Plateau border region of 'dry' soap films. In a simple experimental geometry, we demonstrated that a negative line tension can be attributed to these regions. This result has important consequences for the theoretical description of foams approaching the dry limit.

§ 1. INTRODUCTION

Foams are collections of gas bubbles trapped in a complex network of thin liquid or solid films and they have a great deal of practical importance in industrial processes as well as in everyday life (food, shaving foam, insulators, etc.) (Weaire and Hutzler 1999). Establishing the relations between the physical properties of the films and gas, and those of the resulting material remains a fascinating and yet puzzling problem that has motivated numerous experimental and theoretical studies. Interestingly, understanding foam physics exhibits the same types of challenges as those faced when studying granular materials. In these materials, the relations between the individual properties of the solid grains and their collective behaviour are still only partially understood (Jaeger *et al.* 1996, Kadanoff 1999, Duran 2000, Herrman *et al.* 2001). In foams, the problem is further complicated by the possibility that the gas bubbles might deform, which necessitates complete understanding of the local geometry in these systems.

Idealization of liquid foams commonly consists in considering the films as surfaces of negligible thickness, the liquid being entirely contained in curved Plateau (1873) borders of triangular section at the junctions (figure 1). This limit, convenient for a theoretical study, is that of the so-called 'dry foam'; the water content approaches zero and one is left with a skeleton of interconnected thin films. In two dimensions, the decoration theorem states that the structure of a two-dimensional foam of finite liquid fraction ϕ_l corresponds to that of the appropriate dry skeleton

§Author for correspondence. E-mail: Jean-Christophe.Geminard@ens-lyon.fr.




Figure 1. Cross-section of the Plateau border joining three planar soap films. Within the triangle outlined by broken lines, the surface free energy, associated with the air–water interface (black lines) is less than of the skeleton (white lines) which corresponds to the dry limit, $r \rightarrow 0$.

(i.e. a limiting structure for zero liquid fraction in which the vertices shrink to points); in other words, from the dry limit, the increase in ϕ_1 leaves the overall foam geometry unchanged. The liquid accumulates in the Plateau borders and *decorates* the initial point-like vertices of the dry skeleton (Weaire 1999). In three dimensions, there is no equivalent for the decoration theorem. In the following, we demonstrate theoretically that a negative line tension is associated with the accumulation of water in the Plateau border regions. By direct observation of an annular Plateau border, we give the first experimental evidence that such a line tension plays a role in the structure of simple three-dimensional foams at finite liquid fractions.

§ 2. THEORETICAL BACKGROUND

An increase in the amount of water in the Plateau border leads to a decrease in the total surface free energy and, as a consequence, the corresponding line tension is negative. This can be easily demonstrated by considering the case of three planar soap films, assumed to make 120° angles between them and connected by a rectilinear Plateau border (figure 1) (Krotov and Rusanov 1998). In this case, the cross-section of the air–water interface in the Plateau border region is a circle (radius r). Relative to the dry limit ($r \rightarrow 0$), the variation in the surface free energy per unit length can be written

$$\Delta\varepsilon_s = \gamma(\pi - 2 \times 3^{1/2})r \approx -0.32\gamma r, \quad (1)$$

where γ denotes the surface free energy of the air–water interface. The decoration energy $\Delta\varepsilon_s$ is negative and proportional to the radius r of curvature. Thus, the surface free energy per unit length of the Plateau border decreases when the liquid fraction increases. One can determine the corresponding line tension T , by considering a length l of the Plateau border and by calculating the variation in the




Figure 2. Sketch of the experimental situation. A circular soap film is suspended by two catenoidal films attached to two horizontal annular frames. The radius R of the Plateau border that connects the three films is altered by changing either the diameter D of the frames or the distance H between them. The upper catenoidal film makes the angle θ and the lower catenoidal film makes the angle α with the horizontal plane. Experimentally, accurate values of the angles θ and α are obtained by carefully detecting the profile $\rho(z)$.

associated surface free energy at constant water volume V . One obtains the negative line tension

$$T = \left. \frac{\partial l(\Delta \varepsilon_s)}{\partial l} \right|_V = \frac{\Delta \varepsilon_s}{2}. \tag{2}$$

In the dry limit ($r \rightarrow 0$), three soap films are predicted to make 120° angles between them at the vertices (Almgren and Taylor 1976). In the following, we shall show both theoretically and experimentally that, because of the line tension T , the angles can differ significantly from 120° .

§ 3. EXPERIMENTAL SITUATION AND RESULTS

One can easily produce a catenoidal film by stretching soap between two horizontal annular frames. Cutting the initial catenoid in the horizontal plane with the help of a wet spatula, one can add to the system a horizontal circular soap film hanging between two catenoidal films (figure 2). After a short transient (about 5 min), during which drainage occurs, the system equilibrates. Kept in contact with a humid atmosphere in order to avoid excessive drying, the equilibrated system can be observed for several minutes during which our measurements do not exhibit any systematic evolution with time. The radius R of the Plateau border, which surrounds the horizontal film, is altered by changing either the diameter D of the annular frames (usually 1 cm) or the distance H (up to about $0.4 D$) between them[†]. The volume of water in the Plateau border and consequently the radius r depend on the preparation of the system.

Using a charge-coupled device (CCD) camera (Panasonic, WV-CL700), we acquired images of the system from the side along the horizontal axis (figure 3). These images were subsequently digitized and analysed with the help of a personal

[†]For a larger distance H between the frames, the system is unstable and the collapse of the structure leads to the formation of one planar film on each of the frames and of a droplet. This collapse, similar to that of a simple catenoid (Cryer and Steen 1992, Chen and Steen 1997), as well as the response of the structure to a mechanical perturbation will be the subject of further publications.




Figure 3. Picture of the experimental situation. Three soap films are stretched with the help of two horizontal annular frames. The central part of the structure is made of horizontal circular film held by two catenoidal films. In this side view, the horizontal film is hidden by the thick annular Plateau border (the angle between the two catenoidal films equals $120.3 \pm 0.1^\circ$).

computer equipped with a frame grabber board (Data Translation DT2255) and data analysis software (Wavemetrics, Igor Pro 4). We carefully analysed and then excluded the deformations of the images induced by the optical system; we imaged a grid in the same optical conditions as the soap films and found that the only deformation is due to the small anisotropy of the CCD, which we measured to be $1.71 \pm 0.01\%$. The measurements consisted in detecting the profile $\rho(z)$ of the films, above and below the Plateau border. In order to achieve the required accuracy, the numerical procedure scanned first the image from top to bottom, detecting the outline of the catenoidal films to within 1 pixel; the position $\rho(z)$ was then detected within about 0.1 pixel by interpolating several intensity profiles obtained along lines normal to the initial outline with a Gaussian function. In order to exclude the possible residual tilt of the image (always less than 1°) and to determine accurately the axis of revolution, the profile $\rho(z)$ was detected on both the right- and left-hand sides of the image. Both the lower and the upper experimental profiles were then simultaneously interpolated with two different catenoidal profiles (hyperbolic cosines), allowing the free rotation of the image. We determined three quantities in the plane of the horizontal film, where the two catenoidal profiles intersect: the angles θ and α that the upper and lower films respectively make with the horizontal plane and the radius R of the Plateau border (figure 2). We measured the radius r by direct interpolation of the air–water interface observed on additional images taken at higher magnification.

For large radii r , the weight of water in the Plateau border is not necessarily negligible and the experimental situation needs further analysis before we can draw conclusions. (Note that the upper and lower films are not symmetric with respect to the horizontal film in figure 3.) Taking into account the line tension T and the weight $\rho s g$ of water per unit length, one can write the two relations imposed by the local mechanical equilibrium of the Plateau border in the horizontal plane and along the vertical as

$$\cos \theta + \cos \alpha = 1 + \frac{T}{2\gamma R}, \quad (3)$$

$$\sin \theta + \sin \alpha = \frac{\rho s g}{2\gamma}, \quad (4)$$

where γ denotes the surface free energy of the air–water interface, ρ the density of water, g the acceleration due to gravity, and $s = (3^{1/2} - \pi/2)r^2$ the surface area of the Plateau border cross-section. We point out that equations (3) and (4) apply to the

skeleton and must be evaluated in the plane of the horizontal film. Summing the squares of equations (3) and (4), one obtains that the angle $\beta = \theta - \alpha$ satisfies

$$\cos \beta = -\frac{1}{2} + \frac{1}{8} \left(\frac{s}{l_c} \right)^2 + \frac{1}{2} \frac{T}{\gamma R} + \frac{1}{8} \left(\frac{T}{\gamma R} \right)^2, \tag{5}$$

where $l_c = (\gamma/\rho g)^{1/2}$ is the gravitational capillary length.

According to equation (5), in the dry limit ($r \rightarrow 0$), the angle $\beta \rightarrow 120^\circ$. The weight of water tends to decrease the angle β when the characteristic size r of the Plateau border compares with the capillary length l_c . By contrast, the negative line tension T tends to increase the angle β when the radius R of curvature approaches the typical size r of the Plateau border. From equation (5), one can deduce that the correction to β due to gravity is negligible in comparison with that due to the line tension only in the limit $l_c \gg (Rr^3)^{1/4}$. This condition is usually not satisfied in our experimental conditions. Thus, in order to compare accurately the experimental results with the theory, one must consider the relevant quantity $\Sigma = \cos \theta + \cos \alpha$, which is not altered by gravity (equation 3). Moreover, one can better appreciate the magnitude of the line tension effects on the geometry of the structure by considering the angle $\beta^* \equiv 2 \cos^{-1}(\Sigma/2)$ rather than Σ . We expect, from equation (3) in the limit $T \ll \gamma R$, that

$$\beta^* = \frac{2\pi}{3} + \left(1 - \frac{\pi}{2 \times 3^{1/2}} \right) \frac{r}{R} + O \left[\left(1 - \frac{\pi}{2 \times 3^{1/2}} \right)^2 \frac{r^2}{R^2} \right]. \tag{6}$$

Because of the small value $1 - \pi/(2 \times 3^{1/2}) \approx 0.1$ of the prefactor, the nonlinear terms in equation (6) are negligible in any experimental situation. The angle β^* corresponds to the value of the angle $\beta \equiv \theta - \alpha$ expected from measurements in absence of gravity. In the dry limit ($r \rightarrow 0$), $\beta^* = 120^\circ$. Because of the negative line tension T , the angle β^* is about 121° for $R/r = 5$ and remains larger than 120.2° for $R/r = 20$. The agreement in figure 4 between our experimental measurements and equation (6) demonstrates clearly that the negative line tension T , defined by equation (2), is associated with the Plateau border region.

We point out that the cross-section of the air–water interface in the Plateau border region is no longer circular when the Plateau border is curved, because of the additional curvature of about $1/R$ in the horizontal plane. Moreover, because of the line tension T itself, the angles between the films are not 120° . As a consequence, the line tension T differs slightly from the value given by equation (2), which is exact only in the limit of a rectilinear Plateau border. Writing the Lagrange equation and minimizing the total surface free energy of the system at constant water volume, we computed the exact shape of the air–water interface in the Plateau border region as well as that of the films in our experimental geometry (figure 2). We found that the discrepancy between the line tension T and the approximation given by equation (2) agree quantitatively to within about $2.2r/R\%$. Thus, equation (2) provides a good approximation of the line tension T in most practical cases.

§ 4. CONCLUSION

In conclusion, negative line tension is associated with finite water volumes in the Plateau borders joining soap films. This latter line tension affects the overall geometry in three dimensions; as observed experimentally, the angles that the films make at the Plateau border junction can differ significantly from 120° . This effect,




Figure 4. Experimental angle β^* versus experimental ratio r/R : (■), experimental data; (—), equation (6). The angles are known to within 0.1° whereas the radius r is measured to within 1 pixel (to be compared with $R \approx 150$ pixels).

which is not due to gravity, is often neglected in the theoretical description of foams, usually studied in the dry limit and could be important when the volume fraction of water is increased. Kern and Weaire (2003) suggested recently, that negative decoration energy and line tension must be introduced into the modelling of foams approaching the dry limit. Our experimental results reinforce their theoretical approach which consisted in considering a dry skeleton, whose properties are evaluated in the presence of a negative tension.

ACKNOWLEDGEMENTS

The authors would like to thank D. Weaire, N. Kern, F. Graner and V. Bergeron for stimulating discussions on foam physics. One of us (A. Ż.) would like to thank the Centre National de la Recherche Scientifique (France), the Ministère des Affaires Étrangères (France, Polonium 055844XG), and the Committee of Scientific Research (Poland, KBN 2P03B00923) for financial support.

REFERENCES

- ALMGREN, F. J., and TAYLOR, J., 1976, *Sci. Am.*, **235**, 82.
 CHEN, Y.-J., and STEEN, P. H., 1997, *J. Fluid Mech.*, **341**, 245.
 CRYER, S. A., and STEEN P. H., 1992, *J. Colloid. Interface. Sci.*, **154**, 276.
 DURAN, J., 2000, *Sands, Powders, and Grains: An Introduction to the Physics of Granular Materials* (New York: Springer).
 HERRMANN, H. J., LUDING, S., and CAFIERO, R., 2001, *Physica A*, **295**, 93.
 JAEGER, H. M., NAGEL, S. R., and BEHRINGER, R. P., 1996, *Rev. mod. Phys.*, **68**, 1259.
 KADANOFF, L. P., 1999, *Rev. mod. Phys.*, **71**, 435.
 KROTOV, V. V., and RUSANOV, A. I., 1998, *Mendeleev Commun.*, **5**, 177.
 KERN, N., and WEAIRE, D., 2003, *Phil. Mag.*, **83**, 2973.
 PLATEAU, J., 1873, *Statique Expérimentale et Théorique des Liquides Soumis aux Seules Forces Moléculaires* (Paris: Gauthier-Villars).
 WEAIRE, D., 1999, *Phil. Mag. Lett.*, **79**, 491.
 WEAIRE, D., and HUTZLER, S., 1999, *The Physics of Foams* (Oxford: Clarendon).