# Categories for Me (memorandum)

Olivier . Laurent @ens-lyon.fr

October 18, 2013

# Contents

Categories	1
1.1 Constructions	3
1.2 Morphisms	3
1.3 Functors	5
1.4 Objects	9
Monoidal Categories         2.1 Monoids	<b>11</b> 15
Monads	17
Adjunctions	20
Closed Categories	23
2-Categories	<b>25</b>
	1.1 Constructions   1.2 Morphisms   1.3 Functors   1.4 Objects   Objects

# 1 Categories

# **Definition 1** (Category)

A category  $\mathbb{C}$  is given by a class of objects  $obj(\mathbb{C})$  and, for each pair of objects A and B in  $obj(\mathbb{C})$ , a class of morphisms (or arrows)  $\mathbb{C}(A, B)$  from A to B together with:

• *identities*:  $id_A \in \mathbb{C}(A, A)$  for each object A

$$A \xrightarrow{id_A} A$$

• composition:  $\mathbb{C}(A, B) \times \mathbb{C}(B, C) \to \mathbb{C}(A, C)$ , denoted by  $(f, g) \mapsto f$ ; g:



such that the following diagrams commute:



We can "summarize" these four diagrams into:



# **Example 1** (Category Set) The *category of sets* Set is given by:

- objects are sets
- morphisms are functions
- identities are identity functions
- composition is composition of functions

# **Definition 2** (Sub-Category)

A category  $\mathbb{D}$  is a *sub-category* of the category  $\mathbb{C}$  if its objects are objects of  $\mathbb{C}$   $(obj(\mathbb{D}) \subseteq obj(\mathbb{C}))$ , its morphisms are morphisms of  $\mathbb{C}$   $(\mathbb{D}(A, B) \subseteq \mathbb{C}(A, B))$ , its identities are the identities of  $\mathbb{C}$  $(id_A^{\mathbb{D}} = id_A^{\mathbb{C}})$  and its composition is the composition of  $\mathbb{C}$   $(f ; \mathbb{D} g = f ; \mathbb{C} g)$ .  $\mathbb{D}$  is a *full sub-category* of  $\mathbb{C}$  if, whenever A and B are objects of  $\mathbb{D}$ ,  $\mathbb{D}(A, B) = \mathbb{C}(A, B)$ .  $\mathbb{D}$  is a *wide sub-category* of  $\mathbb{C}$  if  $obj(\mathbb{D}) = obj(\mathbb{C})$ .

A full sub-category is characterized by its class of objects.

**Example 2** (Full Wide Sub-Category) The unique full wide sub-category of a category is itself.

# 1.1 Constructions

### **Definition 3** (Dual Category)

The *dual* (or *opposite*)  $\mathbb{C}^{op}$  of a category  $\mathbb{C}$  is the category with:

- objects of  $\mathbb{C}^{op}$  are objects of  $\mathbb{C}$
- morphisms of  $\mathbb{C}^{op}$  from A to B are morphisms of  $\mathbb{C}$  from B to A
- identities of  $\mathbb{C}^{op}$  are identities of  $\mathbb{C}$
- composition of f and g in  $\mathbb{C}^{op}$  is g; f in  $\mathbb{C}$

# **Definition 4** (Unit Category)

The *unit category*  $\mathbb{T}$  is given by:

- a unique object  $\star$
- a unique morphism u from  $\star$  to  $\star$
- $id_{\star} = u$
- u; u = u

### **Definition 5** (Product Category)

The *product*  $\mathbb{C} \times \mathbb{D}$  of two categories  $\mathbb{C}$  and  $\mathbb{D}$  is the category with:

- objects are pairs of objects of  $\mathbb C$  and objects of  $\mathbb D$
- morphisms from (A, A') to (B, B') are pairs of morphisms of C from A to B and morphisms of D from A' to B'
- identity on (A, A') is the pair  $(id_A, id_{A'})$
- composition of (f, f') and (g, g') is (f; f', g; g')

## 1.2 Morphisms

### **Definition 6** (Monomorphism)

A monomorphism f from the object A to the object B (denoted  $f : A \hookrightarrow B$ ) is a morphism from A to B such that for any two morphisms g and h from some object C to A, we have:

$$g; f = h; f \Longrightarrow g = h$$

### **Definition 7** (Epimorphism)

An *epimorphism* f from the object A to the object B (denoted  $f : A \rightarrow B$ ) is a morphism from A to B such that for any two morphisms g and h from B to some object C, we have:

$$f; g = f; h \Longrightarrow g = h$$

It is thus a monomorphism in  $\mathbb{C}^{op}$ .

### **Definition 8** (Idempotent)

A morphism f from the object A to itself is an *idempotent* if f; f = f. This can be written:

$$A \xrightarrow{f} A \xrightarrow{f} A$$

### **Definition 9** (Retract)

An object A is a *retract* of an object B (denoted  $A \triangleleft B$ ) if there exist two morphisms  $s \in \mathbb{C}(A, B)$ and  $r \in \mathbb{C}(B, A)$  such that  $s; r = id_A$ . This can be written:

This can be written:



s is then called a *section* of r, and r is called a *retraction* of s. (s, r) is called a *section-retraction* pair.

If (s, r) is a section-retraction pair, s is a monomorphism and r is an epimorphism. Such monomorphisms and epimorphisms coming from a section-retraction pair are called *split monomorphisms* and *split epimorphisms*. r; s is an idempotent. Such idempotents coming from a section-retraction pair are called *split idempotents*.

Proof page 32

### **Definition 10** (Isomorphism)

An *isomorphism* f from the object A to the object B is a morphism from A to B such that there exists a morphism g from B to A (called the *inverse* of f) such that the following diagrams commute:



We can "summarize" these two diagrams into:



**Property 1** (Retracts and Isomorphisms) *We have:* 

- If there exists an isomorphism between A and B (denoted  $A \simeq B$ ) then both  $A \triangleleft B$  and  $B \triangleleft A$ .
- If  $f \in \mathbb{C}(A, B)$  is both a section and a retraction then it is an isomorphism.

Proof page 32

In particular an isomorphism is both a monomorphism and an epimorphism (the converse does not hold in general).

Proof page 32

### **Definition 11** (Essentially Wide Sub-Category)

 $\mathbb{D}$  is an *essentially wide sub-category* of  $\mathbb{C}$  if it is a sub-category such that, for each object A of  $\mathbb{C}$ , there is an object A' of  $\mathbb{D}$  such that  $A' \simeq A$ .

# 1.3 Functors

### **Definition 12** (Functor)

A *functor* F between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is:

- a function from the objects of  $\mathbb C$  to the objects  $\mathbb D$
- for each A and B, a function from  $\mathbb{C}(A, B)$  to  $\mathbb{D}(FA, FB)$

such that the following diagrams in  $\mathbb{D}$  commute:



A functor from a category to itself is called an *endofunctor*.

### **Example 3** (Constant Functor)

If  $\mathbb{C}$  and  $\mathbb{D}$  are two categories and D is an object of  $\mathbb{D}$ , the *constant functor*  $C_D$  from  $\mathbb{C}$  to  $\mathbb{D}$  is defined by:

- for any  $A \in obj(\mathbb{C}), C_D A = D$
- for any  $f \in \mathbb{C}(A, B), C_D f = id_D$

The constant functor  $C_{\star}$  is the unique functor from any category  $\mathbb{C}$  to  $\mathbb{T}$ .

Proof page 32

# **Example 4** (Inclusion Functor)

If  $\mathbb{D}$  is a sub-category of  $\mathbb{C}$ , the *inclusion functor* I from  $\mathbb{D}$  to  $\mathbb{C}$  is defined by:

- for each  $A \in obj(\mathbb{D})$ , IA = A
- if A and B are in  $obj(\mathbb{D})$  and  $f \in \mathbb{D}(A, B)$ , If = f

We denote by  $Id_{\mathbb{C}}$  the *identity endofunctor* of  $\mathbb{C}$  which is the inclusion functor of  $\mathbb{C}$  into itself.

Proof page 32

# **Example 5** (Category Cat)

The *category of categories* Cat is given by:

- objects are (small) categories
- morphisms are functors
- identities are identity functors

• composition is composition of functors: if F is a functor from  $\mathbb{C}$  to  $\mathbb{D}$  and G is a functor from  $\mathbb{D}$  to  $\mathbb{E}$ , their composition F; G (or GF) is the functor from  $\mathbb{C}$  to  $\mathbb{E}$  which maps the object A to G(FA) and the morphism f to G(Ff).

If F is an endofunctor of a category  $\mathbb{C}$ , we use the notations  $F^2$  for  $F; F = FF, F^3$  for  $F; F; F = FFF, \ldots$ 

Proof page 32

### Property 2 (Preservation of Retracts)

Functors preserve retracts and isomorphisms: if F is a functor,

- $A \triangleleft B \Longrightarrow FA \triangleleft FB$
- $A \simeq B \Longrightarrow FA \simeq FB$

# **Definition 13** (Bi-Functor)

A *bi-functor* from two categories  $\mathbb{C}$  and  $\mathbb{D}$  to a category  $\mathbb{E}$  is a functor from  $\mathbb{C} \times \mathbb{D}$  to  $\mathbb{E}$ . More concretely, if it is given by:

- a function from  $obj(\mathbb{C}) \times obj(\mathbb{D})$  to  $obj(\mathbb{E})$
- for each A and B in  $obj(\mathbb{C})$  and A' and B' in  $obj(\mathbb{D})$ , a function from  $\mathbb{C}(A, B) \times \mathbb{D}(A', B')$  to  $\mathbb{E}(FAA', FBB')$

such that the following diagrams in  $\mathbb{E}$  commute:

One often uses the notations FAf for  $Fid_Af$  and FfA for  $Ffid_A$ , if A is an object.

### Example 6 (Homset Functor)

The *homset functor*  $\mathbb{C}(\_,\_)$  of a category  $\mathbb{C}$  is the bi-functor from  $\mathbb{C}^{op}$  and  $\mathbb{C}$  to Set given by:

- $\mathbb{C}(\underline{A},\underline{A}) = \mathbb{C}(A,B)$
- $\mathbb{C}(\underline{a},\underline{b})(f,g)h = f; h; g \text{ (for } f \in \mathbb{C}(A',A), g \in \mathbb{C}(B,B') \text{ and } h \in \mathbb{C}(A,B))$

### **Example 7** (Fixed Component Bi-Functor)

If F is a bi-functor from  $\mathbb{C}$  and  $\mathbb{D}$  to  $\mathbb{E}$  and if A is an object of C, we can define a functor  $F_A$  from  $\mathbb{D}$  to  $\mathbb{E}$  by:

- for any object B of  $\mathbb{D}$ ,  $F_A B = F A B$
- for any morphism  $g \in \mathbb{D}(B, B'), F_A g = Fid_A^{\mathbb{C}}g$

### **Definition 14** (Full and Faithful Functors)

A functor F between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is *full* if, for any pair (A, B) of objects of  $\mathbb{C}$ , F is surjective from  $\mathbb{C}(A, B)$  to  $\mathbb{D}(FA, FB)$ .

A functor F between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is *faithful* if, for any pair (A, B) of objects of  $\mathbb{C}$ , F is injective from  $\mathbb{C}(A, B)$  to  $\mathbb{D}(FA, FB)$ .

### **Definition 15** (Essentially Surjective Functor)

A functor F between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is *essentially surjective* if, for each object A' of  $\mathbb{D}$ , there exists an object A of  $\mathbb{C}$  such that A' is isomorphic to FA.

### **Example 8** (Inclusion Functor (bis))

If  $\mathbb{D}$  is a sub-category of  $\mathbb{C}$ , the inclusion functor is faithful. It is full if and only if  $\mathbb{D}$  is a full sub-category of  $\mathbb{C}$ . It is essentially surjective if and only if  $\mathbb{D}$  is an essentially wide sub-category of  $\mathbb{C}$ .

## Example 9 (Projection Functor)

Let  $\mathbb{C}$  and  $\mathbb{D}$  be two categories, the *projection functor* P from  $\mathbb{C} \times \mathbb{D}$  to  $\mathbb{C}$  is defined by:

- for each  $(A, B) \in obj(\mathbb{C} \times \mathbb{D}), P(A, B) = A \in obj(\mathbb{C})$
- if A and A' are objects in  $\mathbb{C}$ , B and B' are objects in  $\mathbb{D}$ , and  $(f,g) \in \mathbb{C} \times \mathbb{D}((A,B), (A',B'))$ ,  $P(f,g) = f \in \mathbb{C}(A,A')$

It is a full functor if  $\mathbb{D}$  has at least one morphism between any two objects.

Proof page 33

### **Definition 16** (Algebra)

An *algebra* for the endofunctor F is a pair  $(A, h_A)$  where:

- A is an object
- $h_A$  is a morphism from FA to A

### **Definition 17** (Algebra Morphism)

An *algebra morphism* f from  $(A, h_A)$  to  $(B, h_B)$  is a morphism from A to B such that the following diagram commutes:



If F is a functor, its *category of algebras* Alg(F) has objects the algebras of F and morphisms the algebra morphisms between them.

### **Definition 18** (Natural Transformation)

A transformation  $\alpha$  between two functions F and G from the objects of a category  $\mathbb{C}$  to the objects of a category  $\mathbb{D}$  (in particular between two functors from  $\mathbb{C}$  to  $\mathbb{D}$ ) is a family  $(\alpha_A)_{A \in obj(\mathbb{C})}$  of morphisms from FA to GA.

A transformation  $\alpha$  between two functors F and G is *natural* if the following diagram in  $\mathbb{D}$  commutes for all  $f \in \mathbb{C}(A, B)$ :

$$\begin{array}{c} FA \xrightarrow{Ff} FB \\ \alpha_A \\ \downarrow \\ GA \xrightarrow{Gf} GB \end{array}$$

It is represented:



A *natural isomorphism* is a natural transformation where each element is an isomorphism.

### **Example 10** (Identity Natural Transformation)

If F is a functor between the categories  $\mathbb{C}$  and  $\mathbb{D}$ ,  $(id_{FA})_{A \in obj(\mathbb{C})}$  is a natural isomorphism from F to itself.

#### Proof page 33

### **Definition 19** (Vertical Composition)

Let F, G and H be three functors between the same two categories  $\mathbb{C}$  and  $\mathbb{D}$ , if  $\alpha$  is a natural transformation for F to G and  $\beta$  is a natural transformation from G to H, the *vertical composition*  $\alpha$ ;<sup>1</sup> $\beta$  is the natural transformation from F to H defined by  $(\alpha; {}^{1}\beta)_{A} = \alpha_{A}; \beta_{A}$ .



PROOF PAGE 33

### **Definition 20** (Horizontal Composition)

Let  $\mathbb{C}$ ,  $\mathbb{D}$  and  $\mathbb{E}$  be three categories, F and F' be two functors from  $\mathbb{C}$  to  $\mathbb{D}$  and G and G' be two functors from  $\mathbb{D}$  to  $\mathbb{E}$ , if  $\alpha$  is a natural transformation for F to F' and  $\beta$  is a natural transformation from G to G', the *horizontal composition*  $\alpha$ ;  $^{0}\beta$  is the natural transformation from F; G to F'; G' defined by  $(\alpha; ^{0}\beta)_{A} = G\alpha_{A}; \beta_{F'A} = \beta_{FA}; G'\alpha_{A}$ .



Proof page 33

### **Example 11** (Category of Functors)

Let  $\mathbb{C}$  and  $\mathbb{D}$  be two categories, the *category of functors*  $\mathbb{F}unc(\mathbb{C}, \mathbb{D})$  is given by:

- objects are functors between  $\mathbb{C}$  and  $\mathbb{D}$
- morphisms are natural transformations
- identities are the identity natural transformations
- composition is the vertical composition of natural transformations

# 1.4 Objects

**Definition 21** (Terminal Object)

A terminal object in a category  $\mathbb{C}$  is an object  $\top$  such that, for any object A of  $\mathbb{C}$ , there exists a unique morphism  $t_A$  from A to  $\top$ .

If  $\mathbb{C}$  is a category with a terminal object  $\top$ , a *point* of an object A of  $\mathbb{C}$  is a morphism from  $\top$  to A.

### **Definition 22** (Initial Object)

An *initial object* in a category  $\mathbb{C}$  is an object  $\perp$  such that, for any object A of  $\mathbb{C}$ , there exists a unique morphism  $i_A$  from  $\perp$  to A.

It is thus a terminal object in  $\mathbb{C}^{op}$ .

A zero object is an object 0 which is both initial and terminal. If 0 is a zero object in the category  $\mathbb{C}$  and A and B are two objects of  $\mathbb{C}$ , the zero morphism  $z_{A,B}$  is:

$$A \xrightarrow{t_A} 0 \xrightarrow{i_A} B$$

### **Definition 23** (Product)

A *product* of two objects A and B in a category  $\mathbb{C}$  is a triple  $(A \times B, \pi_A, \pi_B)$  where:

- $A \times B$  is an object of  $\mathbb{C}$
- $\pi_A$  is a morphism from  $A \times B$  to A
- $\pi_B$  is a morphism from  $A \times B$  to B

such that, for any triple (C, f, g), where C is an object of  $\mathbb{C}$ , f is a morphism from C to A and g is a morphism from C to B, there exists a unique morphism  $\langle f, g \rangle$  from C to  $A \times B$  such that  $\langle f, g \rangle$ ;  $\pi_A = f$  and  $\langle f, g \rangle$ ;  $\pi_B = g$ .

This can be written:



If  $(A \times A, \pi_A^l, \pi_A^r)$  is a product of A and A in  $\mathbb{C}$ , the *diagonal morphism*  $\Delta_A$  is  $\langle id_A, id_A \rangle$  from A to  $A \times A$ . It a section of both projections  $\pi_A^l$  and  $\pi_A^r$ .

A category equipped with a product for each pair of objects and which has a terminal object is called a *cartesian category*. In such a category, one can form all products of finite families of objects. If  $\mathbb{C}$  is a cartesian category,  $\times$  defines a bi-functor from  $\mathbb{C}$  and  $\mathbb{C}$  to  $\mathbb{C}$ , and  $\Delta$  is a natural transformation from  $Id_{\mathbb{C}}$  to  $-\times$ .

**PROOF PAGE 33** 

### **Definition 24** (Co-Product)

A *co-product* of two objects A and B in a category  $\mathbb{C}$  is a triple  $(A + B, \iota_A, \iota_B)$  where:

- A + B is an object of  $\mathbb{C}$
- $\iota_A$  is a morphism from A to A + B
- $\iota_B$  is a morphism from B to A + B

such that, for any triple (C, f, g), where C is an object of  $\mathbb{C}$ , f is a morphism from A to C and g is a morphism from B to C, there exists a unique morphism [f, g] from A + B to C such that  $\iota_A$ ; [f, g] = f and  $\iota_B$ ; [f, g] = g.



It is thus a product in  $\mathbb{C}^{op}$ .

If  $(A + A, \iota_A^l, \iota_A^r)$  is a co-product of A and A in  $\mathbb{C}$ , the *co-diagonal morphism*  $\nabla_A$  is  $[id_A, id_A]$  from A + A to A.

# Example 12 (Products and Co-Products in Set)

If A and B are two sets, the cartesian product  $A \times B$  (with the projection functions) defines a product of A and B in Set. The singleton set  $\{\star\}$  is terminal in Set. With this structure, Set is a cartesian category.

The disjoint union  $A \uplus B$  (with the injection functions) is a co-product in Set. The empty set  $\emptyset$  is an initial object in Set.

### Proof page 33

# Example 13 (Products in Cat)

If  $\mathbb{C}$  and  $\mathbb{D}$  are two categories, the product category  $\mathbb{C} \times \mathbb{D}$  (with the projection functors) defines a product of  $\mathbb{C}$  and  $\mathbb{D}$  in  $\mathbb{C}$ at. The unit category  $\mathbb{T}$  is terminal in  $\mathbb{C}$ at. With this structure,  $\mathbb{C}$ at is a cartesian category.

### Proof page 34

### **Example 14** (Co-Products in Cat)

If  $\mathbb{C}$  and  $\mathbb{D}$  are two categories, the category  $\mathbb{C} + \mathbb{D}$  is given by:

- objects are in the disjoint union  $obj(\mathbb{C}) \uplus obj(\mathbb{D})$
- morphisms from (0, A) to (0, B) are C(A, B), morphisms from (1, A') to (1, B') are D(A', B') (and there is no morphism from (i, A) to (j, B') if i ≠ j)
- composition and identities come from those of  $\mathbb C$  and  $\mathbb D$

Up to the identification of  $obj(\mathbb{C})$  and  $obj(\mathbb{D})$  with their disjoint copies in  $obj(\mathbb{C}) \uplus obj(\mathbb{D})$ , one can consider the inclusion functors as functors from  $\mathbb{C}$  to  $\mathbb{C} + \mathbb{D}$  and from  $\mathbb{D}$  to  $\mathbb{C} + \mathbb{D}$ . The category  $\mathbb{C} + \mathbb{D}$  with these two functors defines a co-product of  $\mathbb{C}$  and  $\mathbb{D}$  in  $\mathbb{C}$ at.

The *empty category*  $\perp$  with no object and no morphism is initial in Cat.

### **Definition 25** (Bi-Product)

Let  $\mathbb{C}$  be a category with a zero object 0 and A and B two objects of  $\mathbb{C}$ , a *bi-product* of A and B is a 5-tuple  $(A \oplus B, \iota_A, \iota_B, \pi_A, \pi_B)$  where:

- $(A \oplus B, \pi_A, \pi_B)$  is a product of A and B in  $\mathbb{C}$
- $(A \oplus B, \iota_A, \iota_B)$  is a co-product of A and B in  $\mathbb{C}$

and such that:

$$\iota_A ; \pi_A = id_A$$
  
 $\iota_B ; \pi_B = id_B$   
 $\iota_A ; \pi_B = z_{A,B}$   
 $\iota_B ; \pi_A = z_{B,A}$ 

### **Definition 26** (Equalizer)

An *equalizer* of two morphisms f and g between the same two objects A and B in a category  $\mathbb{C}$  is a pair (E, e) where E is an object of  $\mathbb{C}$  and e is a morphism from E to A such that e; f = e; hand, for any pair (E', e'), where E' is an object of  $\mathbb{C}$  and e' is a morphism from E' to A such that e'; f = e'; g, there exists a unique morphism h from E' to E such that e' = h; e. This can be written:



If (E, e) is an equalizer, e is a monomorphism. Such monomorphisms coming from an equalizer are called *regular monomorphisms*.

Proof page 34

# 2 Monoidal Categories

**Definition 27** (Monoidal Category) A *monoidal category* is a 6-tuple  $(\mathbb{C}, \otimes, \mathbf{1}, a, u^l, u^r)$  where:

- $\bullet \ \otimes$  is a bi-functor from  $\mathbb C$  and  $\mathbb C$  to  $\mathbb C$
- 1 is an object of  $\mathbb{C}$
- *a* is a natural isomorphism from  $(\_ \otimes \_') \otimes \_''$  to  $\_ \otimes (\_' \otimes \_'')$
- $u^l$  is a natural isomorphism from  $Id_{\mathbb{C}}$  to  $_{-}\otimes 1$
- $u^r$  is a natural isomorphism from  $Id_{\mathbb{C}}$  to  $1\otimes$  \_

such that the following diagrams commute:



A monoidal category is *strict* if the natural isomorphisms a,  $u^l$  and  $u^r$  are the identity natural isomorphism.

A symmetric monoidal category is a 7-tuple  $(\mathbb{C}, \otimes, 1, a, u^l, u^r, s)$  where:

- $(\mathbb{C}, \otimes, 1, a, u^l, u^r)$  is a monoidal category
- s is a natural isomorphism from  $\_ \otimes \_'$  to  $\_' \otimes \_$

such that the following diagrams commute:

$$\begin{array}{c} A\otimes B \xrightarrow{s_{A,B}} B\otimes A \\ & & \downarrow \\ & & \downarrow \\ & & \downarrow \\ & & A\otimes B \end{array}$$

From this definition, it is possible to deduce that, in any monoidal category,  $u_1^r = u_1^l$ . PROOF PAGE 35 From this definition, it is possible to deduce that, in any symmetric monoidal category:



### Proof page 35

If  $(\mathbb{C}, \otimes, 1, a, u^l, u^r)$  is a monoidal category (resp. a symmetric monoidal category) then  $(\mathbb{C}^{op}, \otimes, 1, a^{-1}, u^{l-1}, u^{r-1})$  as well.

# Example 15 (Cartesian Category)

A cartesian category  $\mathbb{C}$  is a symmetric monoidal category  $(\mathbb{C}, \times, \top)$  with the natural isomorphisms:

- $a_{A,B,C} = \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle$
- $u_A^l = \langle id_A, t_A \rangle$
- $u_A^r = \langle t_A, id_A \rangle$
- $s_{A,B} = \langle \pi_B, \pi_A \rangle$

### Proof page 36

### **Definition 28** (Monoidal Functor)

A monoidal functor between two monoidal categories  $(\mathbb{C}, \otimes, 1)$  and  $(\mathbb{D}, \boxtimes, I)$  is a triple (F, m, n) where:

- F is a functor from  $\mathbb{C}$  to  $\mathbb{D}$
- *m* is a natural transformation from  $F_{-} \boxtimes F_{-}'$  to  $F(-\otimes -')$
- n is a morphism from I to F1

such that the following diagrams in  $\mathbb{D}$  commute:

$$\begin{array}{c|c} (FA \boxtimes FB) \boxtimes FC \xrightarrow{a_{FA,FB,FC}} FA \boxtimes (FB \boxtimes FC) \\ \hline m_{A,B} \boxtimes FC & & & & & & \\ F(A \otimes B) \boxtimes FC & & & & & & \\ m_{A \otimes B,C} & & & & & & \\ F((A \otimes B) \otimes C) \xrightarrow{m_{A,B,C}} F(A \otimes (B \otimes C)) \\ \hline \end{array}$$

$$FA \xrightarrow{u_{FA}^{l}} FA \boxtimes I$$

$$Fa \boxtimes FA \boxtimes FA$$

$$Fu_{A}^{l} \xrightarrow{FA \boxtimes F1}$$

$$F(A \otimes 1)$$

$$FA \boxtimes FA \boxtimes FA$$

$$FA \boxtimes FA \boxtimes FA$$

$$Fa \boxtimes FA$$

If  $\mathbb{C}$  and  $\mathbb{D}$  are symmetric monoidal, a *symmetric monoidal functor* is a monoidal functor such that the following diagram in  $\mathbb{D}$  commutes:

$$\begin{array}{c|c} FA \boxtimes FB \xrightarrow{s_{FA,FB}} FB \boxtimes FA \\ \hline m_{A,B} \\ \downarrow \\ F(A \otimes B) \xrightarrow{Fs_{A,B}} F(B \otimes A) \end{array}$$

Let (F, m, n) be a monoidal functor, F is *strong* if  $m_{A,B}$  and n are isomorphisms and F is *strict* if they are equalities.

#### **Definition 29** (Co-Monoidal Functor)

A co-monoidal functor between two monoidal categories  $(\mathbb{C}, \otimes, 1)$  and  $(\mathbb{D}, \boxtimes, \mathbb{I})$  is a triple (F, m, n) which is a monoidal functor between  $(\mathbb{C}^{op}, \otimes, 1)$  and  $(\mathbb{D}^{op}, \boxtimes, \mathbb{I})$ , thus: *m* natural transformation from  $F(\_\otimes\_')$  to  $F\_\boxtimes F\_'$  and *n* morphism from F1 to  $\mathbb{I}$ . We thus have the following commutative diagrams:



**Definition 30** (Monoidal Natural Transformation)

A monoidal natural transformation  $\alpha$  between two monoidal functors F and G between the same two monoidal categories  $(\mathbb{C}, \otimes, 1)$  and  $(\mathbb{D}, \boxtimes, I)$  is a natural transformation such that the following diagrams in  $\mathbb{D}$  commute:

# 2.1 Monoids

### **Definition 31** (Monoid)

A *monoid* in a monoidal category  $(\mathbb{C}, \otimes, 1)$  is a triple  $(A, c_A, w_A)$  where:

- A is an object
- $c_A$  is a morphism from  $A \otimes A$  to A
- $w_A$  is a morphism from 1 to A

that is:

$$A \otimes A \xrightarrow{c_A} A \xleftarrow{w_A} 1$$

such that the following diagrams commute:



If  $\mathbb{C}$  is symmetric monoidal, a monoid is *symmetric* if the following diagram commutes:



### **Definition 32** (Monoidal Morphism)

A monoidal morphism f between two monoids  $(A, c_A, w_A)$  and  $(B, c_B, w_B)$  in a monoidal category is a morphism from A to B such that the following diagrams commute:



Monoids of a monoidal category  $(\mathbb{C}, \otimes, 1)$  and monoidal morphisms between them define a category  $Mon(\mathbb{C})$  called the *category of monoids* of  $\mathbb{C}$ .

### Definition 33 (Co-Monoid)

A co-monoid in  $\mathbb{C}$  is a monoid in  $\mathbb{C}^{op}$ . It is thus a triple  $(A, d_A, e_A)$  with  $d_A$  morphism from A to

 $A \otimes A$  and  $e_A$  morphism from A to 1 such that:





### **Definition 34** (Co-Monoidal Morphism)

A co-monoidal morphism f between two co-monoids  $(A, d_A, e_A)$  and  $(B, d_B, e_B)$  in a monoidal category is a morphism from A to B such that the following diagrams commute:



Co-monoids of a monoidal category  $(\mathbb{C}, \otimes, 1)$  and co-monoidal morphisms between them define a category  $coMon(\mathbb{C})$  called the *category of co-monoids* of  $\mathbb{C}$ .

### Example 16 (Co-Monoids and Cartesian Categories)

In a cartesian category  $\mathbb{C}$ , each object A comes with a canonical structure of symmetric co-monoid  $(A, \Delta_A, t_A)$ . Since any morphism of  $\mathbb{C}$  is co-monoidal for these co-monoid structures, one can see  $\mathbb{C}$  as a full sub-category of coMon( $\mathbb{C}$ ).

Conversely, let  $\mathbb{C}$  be a monoidal category and  $\mathbb{M}$  be a sub-category of  $\mathsf{coMon}(\mathbb{C})$  such that:

- the forgetful functor U from M to C which maps triples  $(A, d_A, e_A)$  to A is full and injective on objects
- if A and B are in the image of U then  $A \otimes B$  as well
- 1 is in the image of U
- the following diagram commutes:



• 
$$e_1 = id_1$$

then UM is a cartesian category with  $\otimes$  as product and 1 as terminal object.

### **Property 3** (Preservation of Monoids)

If (F, m, n) is a monoidal functor from  $(\mathbb{C}, \otimes, 1)$  to  $(\mathbb{D}, \boxtimes, \mathbb{I})$  and  $(A, c_A, w_A)$  is a monoid in  $(\mathbb{C}, \otimes, 1)$ , then  $(FA, m_{A,A}; Fc_A, n; Fw_A)$  is a monoid in  $(\mathbb{D}, \boxtimes, \mathbb{I})$ . We say that monoidal functors preserve monoids.

 $FA \boxtimes FA \xrightarrow{m_{A,A}} F(A \otimes A) \xrightarrow{Fc_A} FA \xleftarrow{Fw_A} F1 \xleftarrow{n} I$ 

Similarly, symmetric monoidal functors preserve symmetric monoids, and co-monoidal functors preserve co-monoids.

Proof page 39

# 3 Monads

### **Definition 35** (Monad)

A *monad* on a category  $\mathbb{C}$  is a triple  $(T, \eta, \mu)$  where:

- T is an endofunctor of  $\mathbb{C}$
- $\eta$  is a natural transformation from  $Id_{\mathbb{C}}$  to T
- $\mu$  is a natural transformation from  $T^2$  to T

$$T^2 \xrightarrow{\mu} T \xleftarrow{\eta} Id_{\mathbb{C}}$$

such that the following diagrams commute:

A co-monad on  $\mathbb{C}$  is a monad on  $\mathbb{C}^{op}$ , that is a triple  $(T, \varepsilon, \delta)$   $(T \text{ endofunctor of } \mathbb{C}, \varepsilon \text{ natural transformation from } T \text{ to } Id_{\mathbb{C}}$  and  $\delta$  natural transformation from T to  $T^2$ ) such that:

# **Definition 36** (Kleisli Triple)

A *Kleisli triple* on a category  $\mathbb{C}$  is a triple  $(T, \eta, (\_)^{\dagger})$  where:

- T is a function from  $obj(\mathbb{C})$  to  $obj(\mathbb{C})$
- $\eta$  is a transformation from  $Id_{\mathbb{C}}$  to T
- $(_)^{\dagger}$  is a function from  $\mathbb{C}(A, TB)$  to  $\mathbb{C}(TA, TB)$

such that the following diagrams commute:



The notions of monad and Kleisli triple are equivalent through:

$$(T, \eta, \mu) \mapsto (T, \eta, T_{-}; \mu)$$
$$(T, \eta, (_{-})^{\dagger}) \mapsto (T, \eta, id_{T_{-}}^{\dagger})$$

# Definition 37 (Strong Monad)

A strong monad on a monoidal category  $\mathbb{C}$  is a monad equipped with  $\tau$  where:

•  $\tau$  is a natural transformation from  $_{-}\otimes T_{-}'$  to  $T(_{-}\otimes _{-}')$ 

such that the following diagrams commute:

$$1 \otimes TA \xrightarrow{\tau_{1,A}} T(1 \otimes A) \qquad (A \otimes B) \otimes TC \xrightarrow{\tau_{A \otimes B,C}} T((A \otimes B) \otimes C)$$

$$\downarrow^{Tu_{A}} \downarrow^{Tu_{A}^{r}} \qquad a_{A,B,TC} \downarrow \qquad \downarrow^{Ta_{A,B,C}}$$

$$TA \qquad A \otimes (B \otimes TC) \xrightarrow{\tau_{A \otimes T_{B,C}}} A \otimes T(B \otimes C) \xrightarrow{\tau_{A,B \otimes C}} T(A \otimes (B \otimes C))$$

**Definition 38** (Commutative Monad)

A *commutative monad* on a symmetric monoidal category  $\mathbb{C}$  is a strong monad such that, if:

$$\tau'_{A,B} = TA \otimes B \xrightarrow{s_{TA,B}} B \otimes TA \xrightarrow{\tau_{B,A}} T(B \otimes A) \xrightarrow{Ts_{B,A}} T(A \otimes B)$$

then the following diagram commutes:



### **Definition 39** (Monoidal Monad)

A monad  $(T, \eta, \mu)$  on a monoidal category  $\mathbb{C}$  is *monoidal* if T is a monoidal functor, and  $\eta$  and  $\mu$  are monoidal natural transformations.

If  $\mathbb{C}$  is symmetric monoidal, the monad is *symmetric monoidal* if, moreover, T is a symmetric monoidal functor.

# Property 4 (Monoidal and Commutative Monads)

Let  $\mathbb{C}$  be a symmetric monoidal category and T be a strong monad on  $\mathbb{C}$ :

• T equipped with either:

$$TA \otimes TB \xrightarrow{\tau_{TA,B}} T(TA \otimes B) \xrightarrow{T\tau'_{A,B}} T^2(A \otimes B) \xrightarrow{\mu_{A \otimes B}} T(A \otimes B)$$

or

$$TA \otimes TB \xrightarrow{\tau'_{A,TB}} T(A \otimes TB) \xrightarrow{T\tau_{A,B}} T^2(A \otimes B) \xrightarrow{\mu_{A \otimes B}} T(A \otimes B)$$

and  $\eta_1: 1 \to T1$  is a monoidal functor

- in both cases,  $\eta$  and  $\mu$  are monoidal natural transformations
- T is a symmetric monoidal functor  $\iff$  T is a commutative monad

### **Definition 40** (Algebra)

An *algebra* for the monad T is a pair  $(A, h_A)$  which is an algebra for the functor T such that the following diagrams commute:



### Example 17 (Free Algebra)

For any object A,  $(TA, \mu_A)$  is an algebra called the *free algebra* generated by A.

### **Definition 41** (Eilenberg-Moore Category)

If T is a monad on the category  $\mathbb{C}$ , its *category of algebras* is the full sub-category of the category of algebras of the functor T whose objects are the algebras of the monad T. It is also called the *Eilenberg-Moore category* of T and denoted  $\mathbb{C}^T$ .

### **Definition 42** (Kleisli Category)

If T is a monad on the category  $\mathbb{C}$ , the *Kleisli category*  $\mathbb{C}_T$  has objects the objects of  $\mathbb{C}$  and for morphisms:  $\mathbb{C}_{\mathbb{T}}(A, B) = \mathbb{C}(A, TB)$ . The identities are  $\eta_A \in \mathbb{C}(A, TA)$ , and the composition of  $f \in \mathbb{C}(A, TB)$  and  $g \in \mathbb{C}(B, TC)$  is  $f ; Tg ; \mu_C \in \mathbb{C}(A, TC)$ .

$$A \xrightarrow{f} TB \xrightarrow{Tg} TTC \xrightarrow{\mu_C} TC$$
$$\xrightarrow{f; \mathbb{C}_{Tg}} TC$$

### **Definition 43** (Distributive Law)

If  $(T, \eta^T, \mu^T)$  and  $(S, \eta^S, \mu^S)$  are two monads on the category  $\mathbb{C}$ , a *distributive law* of T over S is a natural transformation l from ST to TS such that the following diagrams commute:







Let  $(T, \eta^T, \mu^T)$  and  $(S, \eta^S, \mu^S)$  be two monads on the category  $\mathbb{C}$ , and l be a distributive law of T over S, TS equipped with

$$A \xrightarrow{\eta_A^S} SA \xrightarrow{\eta_{SA}^T} TSA \qquad \text{and} \qquad TSTSA \xrightarrow{Tl_{SA}} TTSSA \xrightarrow{\mu_{SSA}^T} TSSA \xrightarrow{T\mu_A^S} TSA$$

is a monad on  $\mathbb{C}$ .

# 4 Adjunctions

**Definition 44** (Adjunction) An *adjunction*  $F \dashv G$  between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is a triple  $(F, G, \varphi)$  where:

- F is a functor from  $\mathbb{C}$  to  $\mathbb{D}$
- G is a functor from  $\mathbb{D}$  to  $\mathbb{C}$

•  $\varphi$  is a natural isomorphism from the functor  $\mathbb{D}(F_{-}, \underline{\ }')$  to the functor  $\mathbb{C}(\underline{\ }, G_{-}')$  (both from  $\mathbb{C}^{op} \times \mathbb{D}$  to Set).

$$\mathbb{C} \underbrace{\stackrel{F}{\underbrace{\qquad}}}_{G} \mathbb{D} \underbrace{\qquad \qquad \underbrace{FA \longrightarrow B'}_{A \longrightarrow GB'} \varphi}_{}$$

Equivalently, an *adjunction*  $F \dashv G$  between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is a quadruple  $(F, G, \eta, \varepsilon)$  where:

- F is a functor from  $\mathbb{C}$  to  $\mathbb{D}$
- G is a functor from  $\mathbb{D}$  to  $\mathbb{C}$
- $\eta$  is a natural transformation from  $Id_{\mathbb{C}}$  to GF
- $\varepsilon$  is a natural transformation from FG to  $Id_{\mathbb{D}}$

such that the following diagrams commute:



If  $F \dashv G$  is an adjunction, F is called a *left adjoint* and G is called a *right adjoint*. The diagram underlying the naturality of  $\varphi$  is, in  $\mathbb{C}$ :



The equivalence between the two definitions is given by:

$$\varphi_{A,A'}(f) = A \xrightarrow{\eta_A} GFA \xrightarrow{Gf} GA'$$
$$\eta_A = A \xrightarrow{\varphi_{A,FA}(id_{FA})} GFA$$
$$\varepsilon_{A'} = FGA' \xrightarrow{\varphi_{GA',A'}^{-1}(id_{GA'})} A'$$

**Example 19** (Category of Adjunctions) The *category of adjunctions* Adj is given by:

- objects are (small) categories
- morphisms in  $Adj(\mathbb{C}, \mathbb{D})$  are adjunctions between  $\mathbb{C}$  and  $\mathbb{D}$
- identities are identity adjunctions (*Id*, *Id*, *id*)

• composition is composition of adjunctions: if  $(F, G, \varphi)$  is an adjunction between  $\mathbb{C}$  and  $\mathbb{D}$  and  $(F', G', \varphi')$  is an adjunction between  $\mathbb{D}$  and  $\mathbb{E}$  then  $(F; F', G'; G, \varphi'_{F_{\neg,\neg'}}; \varphi_{\neg,G'_{\neg'}})$  is an adjunction between  $\mathbb{C}$  and  $\mathbb{E}$ .



### **Definition 45** (Monoidal Adjunction)

An adjunction  $(F, G, \eta, \varepsilon)$  between two monoidal categories  $\mathbb{C}$  and  $\mathbb{D}$  is *monoidal* if F and G are monoidal functors and  $\eta$  and  $\varepsilon$  are monoidal natural transformations.

If  $\mathbb{C}$  and  $\mathbb{D}$  are symmetric monoidal, the adjunction is *symmetric monoidal* if, moreover, F and G are symmetric monoidal functors.

In a monoidal adjunction, F is strong.

### **Property 5** (Monad of an Adjunction)

If  $(F, G, \eta, \varepsilon)$  is an adjunction,  $(GF, \eta, G\varepsilon_{F_{-}})$  is a monad called the monad of the adjunction. Similarly,  $(FG, \varepsilon, F\eta_{G_{-}})$  is a co-monad.

If the adjunction is monoidal, the monad is monoidal. If the adjunction is symmetric monoidal, the monad is symmetric monoidal.

# Example 20 (Eilenberg-Moore Adjunction)

Let T be a monad on  $\mathbb{C}$ , let F be the free-algebra functor from  $\mathbb{C}$  to  $\mathbb{C}^T$  associating  $(TA, \mu_A)$  with A, and associating  $Tf \in \mathbb{C}^{\mathbb{T}}((TA, \mu_A), (TB, \mu_B))$  with  $f \in \mathbb{C}(A, B)$ .

Let U be the forgetful functor from  $\mathbb{C}^T$  to  $\mathbb{C}$  associating A with the algebra  $(A, h_A)$  and such that Uf = f.



F is a left adjoint to U and the monad associated with this adjunction is T.

### Example 21 (Kleisli Adjunction)

Let T be a monad on  $\mathbb{C}$ , let E be the embedding functor from  $\mathbb{C}$  to  $\mathbb{C}_T$  associating A with A (EA = A), and associating  $\eta_A$ ;  $Tf \in \mathbb{C}_{\mathbb{T}}(A, B)$  with  $f \in \mathbb{C}(A, B)$ .

Let T' be the functor from  $\mathbb{C}_T$  to  $\mathbb{C}$  defined by T'A = TA and T'f = Tf;  $\mu_B$  for  $f \in \mathbb{C}_{\mathbb{T}}(A, B)$ .



E is a left adjoint to T' and the monad associated with this adjunction is T.

**Example 22** (Category of Adjunctions of a Monad)

Let T be a monad on a category  $\mathbb{C}$ , the category T-Adj of adjunctions of the monad T is given by:

• objects are tuples  $(\mathbb{D}, F, G, \eta, \varepsilon)$  where  $(F, G, \eta, \varepsilon)$  is an adjunction between  $\mathbb{C}$  and  $\mathbb{D}$  which induces the monad T on  $\mathbb{C}$  (Property 5)

• morphisms between  $(\mathbb{D}, F, G, \eta, \varepsilon)$  and  $(\mathbb{D}', F', G', \eta', \varepsilon')$  are functors L from  $\mathbb{D}$  to  $\mathbb{D}'$  such that the following diagram commutes:



and  $L\varepsilon = \varepsilon'_L$ .

The Kleisli adjunction is the initial object of T-Adj. The Eilenberg-Moore adjunction is the terminal object of T-Adj.

### **Definition 46** (Equivalence of Categories)

A functor F between two categories  $\mathbb{C}$  and  $\mathbb{D}$  is an *equivalence of categories* if one of the two following equivalent properties is true:

- There exists an adjunction  $(G, F, \eta, \varepsilon)$  between  $\mathbb{D}$  and  $\mathbb{C}$  such that  $\eta$  and  $\varepsilon$  are natural isomorphisms.
- F is full, faithful and essentially surjective.

Property 6 (Strict Monoidal Categories)

Every monoidal category is equivalent to a strict monoidal category.

**Property 7** (Kleisli Category and Free Algebras)

If T is a monad on the category  $\mathbb{C}$ , the category  $\mathbb{C}_T$  is equivalent to the full-subcategory of  $\mathbb{C}^T$  consisting of free algebras.

# 5 Closed Categories

**Definition 47** (Symmetric Monoidal Closed Category)

A symmetric monoidal category  $(\mathbb{C}, \otimes, 1, a, u^l, u^r, s)$  is *closed* if, for any object A of  $\mathbb{C}$ , the functor  $\mathbb{C} \otimes A$  has a right adjoint (noted  $A \rightarrow \mathbb{C}$ ).

$$\frac{C \otimes A \longrightarrow B}{C \longrightarrow A \multimap B} \ curry$$

In a symmetric monoidal closed category, if f is a morphism from  $C \otimes A$  to B, we denote by curry(f) the induced morphism from C to  $A \multimap B$ . We define  $ev_{A,B}$  as  $curry^{-1}(id_{A \multimap B}) \in \mathbb{C}((A \multimap B) \otimes A, B)$ .

### **Definition 48** (Exponential Object)

If A and B are two objects of a symmetric monoidal category  $\mathbb{C}$ , an *exponential object* of A and B is a pair  $(B^A, ev_{A,B})$  where  $B^A$  is an object of  $\mathbb{C}$  and  $ev_{A,B} \in \mathbb{C}(B^A \otimes A, B)$  such that, for any morphism  $f \in \mathbb{C}(C \otimes A, B)$ , there exists a unique morphism  $\lambda f \in \mathbb{C}(C, B^A)$  such that  $f = (\lambda f \otimes id_A)$ ;  $ev_{A,B}$ .

This can be written:



The notions of symmetric monoidal closed category and exponential object are related by the fact that a symmetric monoidal category is closed if and only if each pair of objects has an associated exponential object.

# Definition 49 (Dual Object)

In a A symmetric monoidal category  $(\mathbb{C}, \otimes, 1, a, u^l, u^r, s)$ , a *dual* of an object A is an object  $A^{\perp}$  with two morphisms  $\eta \in \mathbb{C}(1, A \otimes A^{\perp})$  and  $\varepsilon \in \mathbb{C}(A^{\perp} \otimes A, 1)$  such that the following diagrams commute:



**Definition 50** (Compact Closed Category) A symmetric monoidal category is *compact closed* if each object has a dual object.

**Example 23** (Closure of Compact Closed Categories) A compact closed category is a symmetric monoidal closed category with  $A \multimap \_ = A^{\perp} \otimes \_$ .

Remember (Example 15) that a cartesian category has a canonical symmetric monoidal structure.

### Definition 51

Cartesian Closed Category A cartesian category is *cartesian closed* if, as a symmetric monoidal category, it is closed.

### **Definition 52** (\*-Autonomous Category)

A symmetric monoidal closed category  $\mathbb{C}$  is \*-*autonomous* if it contains a *dualizing object*, that is an object  $\bot$  such that, for each object A of  $\mathbb{C}$ , the following morphism is an isomorphism between A and  $(A \multimap \bot) \multimap \bot$ :

$$curry\left( A \otimes (A \multimap \bot) \xrightarrow{s_{A,A \multimap \bot}} (A \multimap \bot) \otimes A \xrightarrow{ev_{A,\bot}} \bot \right)$$

Example 24 (Compact Closed and \*-Autonomous Categories)

Any compact closed category is \*-autonomous with  $1^{\perp}$  as dualizing object. Any \*-autonomous category such that  $(A \otimes B) \multimap \bot \simeq (B \multimap \bot) \otimes (A \multimap \bot)$  is compact closed with  $A \multimap \bot$  as dual of A.

# 6 2-Categories

**Definition 53** (2-Category) A 2-category  $\mathbb{C}$  is given by:

- a class of objects  $obj(\mathbb{C})$
- for any two objects A and B, a class of 1-morphisms  $\mathbb{C}(A, B)$
- for any two object A and B and any two morphisms f and g in  $\mathbb{C}(A, B)$ , a class of 2-morphisms (or 2-cells)  $\mathbb{C}^2(f, g)$
- for any object A, a *1-identity* morphism  $id_A$  in  $\mathbb{C}(A, A)$
- for any 1-morphism f, a 2-*identity* morphism  $id_f^1$  in  $\mathbb{C}^2(f, f)$
- for any two morphisms  $f \in \mathbb{C}(A, B)$  and  $g \in \mathbb{C}(B, C)$ , a composition  $f ; g \in \mathbb{C}(A, C)$
- for any two 2-morphisms  $\alpha \in \mathbb{C}^2(f,g)$  and  $\beta \in \mathbb{C}^2(g,h)$ , a *vertical composition*  $\alpha$ ;  $\beta \in \mathbb{C}^2(f,h)$
- for any two 2-morphisms  $\alpha \in \mathbb{C}^2(f,g)$  and  $\beta \in \mathbb{C}^2(f',g')$  with f and g in  $\mathbb{C}(A,B)$  and f' and g' in  $\mathbb{C}(B,C)$ , an *horizontal composition*  $\alpha ; {}^0 \beta \in \mathbb{C}^2(f;f',g;g')$

such that:

- $obj(\mathbb{C})$  with 1-morphisms, 1-identities, and composition is a category
- for any two objects A and B,  $\mathbb{C}(A, B)$  with  $\mathbb{C}^2(A, B)$  for morphisms, 2-identities between morphisms of  $\mathbb{C}(A, B)$  for identities, and vertical composition for composition is a category
- obj(ℂ) with 2-morphisms for morphisms, 2-identities between 1-identities as identities, and horizontal composition for composition is a category

and given any four 2-morphisms of the following shape:



we have:



and we also have:



### Example 25 (2-Category Cat)

(Small) Categories with functors for 1-morphisms, natural transformations for 2-morphisms, identity functors for 1-identities, identity natural transformations for 2-identities, composition of functors for composition, vertical composition of natural transformations for vertical composition, and horizontal composition of natural transformations for horizontal composition is a 2-category.

**Example 26** (Monoidal Categories)

A 2-category with one object is the same thing as a strict monoidal category.

**Property 8** (Monoidal Structures in 2-Categories) Each object A of a 2-category  $\mathbb{C}$  defines a strict monoidal category:

- objects are 1-morphisms in  $\mathbb{C}(A, A)$
- morphisms are 2-morphisms between them
- *identities are id*<sup>1</sup>
- composition is vertical composition

- tensor product on objects is composition of 1-morphisms
- tensor product on morphisms is horizontal composition of 2-morphisms
- unit of the tensor is  $id_A$

# Example 27 (Monads as Monoids)

Let  $\mathbb{C}$  be a category, since it is an object in the 2-category  $\mathbb{C}at$ ,  $\mathbb{F}unc(\mathbb{C}, \mathbb{C})$  has a strict monoidal category structure given by Property 8. A monad is exactly a monoid in this monoidal category.

# Index

\*-autonomous category, 24

+, 9 0, 9 ⊥⊥, 10 ⊥, 9 ⊣, **20** ι, 9  $\mathbb{C}^T$ , **19**  $\mathbb{C}_T$ , **19** T, 3 ;, **2 ⊸**, **23 ⊕**, **11** ⊗, 11  $\pi, 9$  $\simeq$ , **4** ev., 23 ×, 3, 9 T, 9 ⊲, **4** ℂ<sup>op</sup>, **3** *curry*, **23** 1-identity, 24 1-morphism, 24 2-category, 24 2-identity, 25 2-morphism, 24

# 1, **11**

Adj, 21 adjunction, 20 Alg, 7 algebra morphism, 7 algebra of a functor, 7, 18 algebra of a monad, 18

bi-functor, 6 bi-product, 11

cartesian category, **9**, 13, 15 cartesian closed category, **24** Cat, **5**, 10, 26 category, **1**  category of adjunctions, 21 category of algebras of a functor, 7 category of algebras of a monad, 19 category of categories, 5 category of co-monoids, 15 category of functors, 8 category of monoids, 15 category of sets, 2 co-diagonal, 10 co-monad, 17, 21 co-monoid, 15 co-monoidal functor, 14 co-monoidal morphism, 15 co-product, 9 commutative monad, 18 coMon, 15 compact closed category, 24 composition, 2 constant functor, 5

diagonal, 9 distributive law, 19 dual category, 3 dual object, 23, 24 dualizing object, 24

Eilenberg-Moore category, **19**, 21 empty category, endofunctor, epimorphism, equalizer, equivalence of categories, essentially surjective functor, **7**, 22 essentially wide sub-category, **5**, 7 exponential object,

faithful functor, **6**, 22 free algebra, **19**, 22 full functor, **6**, 22 full sub-category, **2**, 7 Func, **8**, 26 functor, **5**, 26

homset functor, 6

horizontal composition, 8, 25, 26

### Id, 5

id, 1
idempotent, 4
identity, 1
identity endofunctor, 5
identity functor, 26
inclusion functor, 5, 7
initial object, 9
inverse, 4
isomorphism, 4, 6, 7

Kleisli category, **19**, 22 Kleisli triple, **17** 

left adjoint, 20

# $\mathrm{Mon},\,\mathbf{15}$

monad, 16, 21
monad of an adjunction, 21
monoid, 14
monoidal adjunction, 21
monoidal category, 11, 22
monoidal functor, 13, 16
monoidal monad, 18, 21
monoidal morphism, 15
monoidal natural transformation, 14
monomorphism, 3
morphism, 1

natural isomorphism, 8 natural transformation, 7, 26

obj, **1** object, **1** opposite, **3** point, **9** product, **9** product category, **3**, 10 projection functor, **7**, 10 regular monomorphism, **11** 

retraction, **4** right adjoint, **20**  section, 4 section-retraction pair, 4 Set, 2, 10 split epimorphism, 4 split idempotent, 4 split monomorphism, 4 strict monoidal category, 12, 26 strict monoidal functor, 14 strong monad, 17 strong monoidal functor, 14sub-category, 2, 5, 7 symmetric monoid, 15 symmetric monoidal adjunction, 21 symmetric monoidal category, 12, 23, 24 symmetric monoidal closed category, 23 symmetric monoidal functor, 14 symmetric monoidal monad, 18

terminal object, 9 transformation, 7 unit category, **3**, 10 vertical composition, **8**, 8, **25**, 26

zero morphism, **9** zero object, **9**, 11

wide sub-category, 2

# **Additional Properties**

# **Cartesian Product**

We consider a category  $\mathbb{C}$ , two objects A and B of  $\mathbb{C}$  and a product  $(A \times B, \pi_A, \pi_B)$  of A and B in  $\mathbb{C}$ .

Fact 1 (Pair of Projections)  $\langle \pi_A, \pi_B \rangle = i d_{A \times B}.$ 

PROOF:  $\langle \pi_A, \pi_B \rangle$ ;  $\pi_A = \pi_A = id_{A \times B}$ ;  $\pi_A$  and  $\langle \pi_A, \pi_B \rangle$ ;  $\pi_B = \pi_B = id_{A \times B}$ ;  $\pi_B$  thus, by uniqueness of the pair, we have  $\langle \pi_A, \pi_B \rangle = id_{A \times B}$ .

Fact 2 (Composition with Pair)

Let C and D be two objects of  $\mathbb{C}$ , if  $f \in \mathbb{C}(C, A)$ ,  $g \in \mathbb{C}(C, B)$  and  $h \in \mathbb{C}(D, C)$  then  $h ; \langle f, g \rangle = \langle h ; f, h ; g \rangle$ .

PROOF: We have  $h; \langle f, g \rangle; \pi_A = h; f = \langle h; f, h; g \rangle; \pi_A$  and  $h; \langle f, g \rangle; \pi_B = h; g = \langle h; f, h; g \rangle; \pi_B$ , thus  $h; \langle f, g \rangle = \langle h; f, h; g \rangle$  by uniqueness of the pair.  $\Box$ 

# Monoidal Categories

We consider a monoidal category  $(\mathbb{C}, \otimes, 1, a, u^l, u^r)$ .

**Fact 3** (Equality up to  $\_ \otimes 1$  and  $1 \otimes \_$ ) Let A and B be two objects of  $\mathbb{C}$  and f and g be two morphisms of  $\mathbb{C}$  from A to B,  $f \otimes 1 = g \otimes 1 \iff f = g \iff 1 \otimes f = 1 \otimes g$ .

**PROOF:** We have f = g implies both  $f \otimes 1 = g \otimes 1$  and  $1 \otimes f = 1 \otimes g$ .

Now assume  $f \otimes 1 = g \otimes 1$ , the following diagram commutes:



since the two squares commute by naturality of  $u^l$ . We conclude f = g because  $u^l_B$  is an isomorphism.

Similarly, we obtain the implication  $1 \otimes f = 1 \otimes g \Longrightarrow f = g$  by naturality of  $u^r$ .  $\Box$ 

Fact 4 (Unit of Unit)

Let A be an object of  $\mathbb{C}$ ,  $u_{1\otimes A}^r = 1 \otimes u_A^r : 1 \otimes A \to 1 \otimes (1 \otimes A).$ 

**PROOF:** By naturality of  $u^r$ , we have:



thus, since  $u_A^r$  is an isomorphism,  $u_{1\otimes A}^r = 1 \otimes u_A^r$ .

**Fact 5** (Associativity of Unit) Let A and B be two objects of  $\mathbb{C}$ , the following diagram commutes:



PROOF: Thanks to Fact 3, it is sufficient to prove the commutation of the following diagram (since *a* is an isomorphism):



which commutes by:

- (a) naturality of a
- (b) triangle of monoidal categories
- (c) triangle of monoidal categories
- (d) naturality of a
- (e) pentagon of monoidal categories



# Proofs

# **Definition 9**

- If g; s = h; s then  $g = g; id_A = g; s; r = h; s; r = h; id_A = h$ .
- If r; g = r; h then  $g = id_A; g = s; r; g = s; r; h = id_A; h = h$ .
- $r; s; r; s = r; id_A; s = r; s$

# **Property 1**

- Let f from A to B be an isomorphism and  $f^{-1}$  be its inverse, we have  $f; f^{-1} = id_A$  and  $f^{-1}; f = id_B$ .
- There exist  $g \in \mathbb{C}(B, A)$  such that  $f; g = id_A$  and  $h \in \mathbb{C}(B, A)$  such that  $h; f = id_B$  thus  $h = h; id_A = h; f; g = id_B; g = g$  and we conclude that g = h is an inverse of f.

# **Comment Page 4**

We give a direct proof: let f be an isomorphism from A to B,  $f^{-1}$  be its inverse, if g and g' are morphisms from A' to A then g; f = g'; f implies g = g;  $id_A = g$ ;  $f^{-1} = g'$ ; f;  $f^{-1} = g'$ ;  $id_A = g'$ . If h and h' are morphisms from B to B' then f; h = f; h' implies  $h = f^{-1}$ ; f;  $h = f^{-1}$ ; f; h' = h'. In the following category:

$$id_A \bigcirc A \xrightarrow{f} B \bigcirc id_B$$

with  $id_A$ ; f = f and f;  $id_B = id_B$ , f is both a monomorphism and an epimorphism but it is not an isomorphism since there is no morphism from B to A.

# Example 3

Let A be an object of  $\mathbb{C}$ ,  $C_D id_A = id_D = id_{C_D A}$ , and if  $f \in \mathbb{C}(A, B)$  and  $g \in \mathbb{C}(B, C)$  then  $C_D(f;g) = id_D = id_D$ ;  $id_D = C_D f$ ;  $C_D g$ .

A functor F from  $\mathbb{C}$  to  $\mathbb{T}$  must satisfy  $FA = \star$  for any object A of  $\mathbb{C}$  since  $\star$  is the unique object of  $\mathbb{T}$ . We must then have  $Ff \in \mathbb{T}(\star, \star) = \{id_{\star}\}$ , so  $F = C_{\star}$ .

# Example 4

We have  $Iid_A = id_A = id_{IA}$  and I(f;g) = f; g = If; Ig.

# Example 5

If  $\mathbb{C}$  and  $\mathbb{D}$  are two (small) categories and F is a functor from  $\mathbb{C}$  to  $\mathbb{D}$ , let A be an object of  $\mathbb{C}$ , we have  $(Id_{\mathbb{C}}; F)A = FId_{\mathbb{C}}A = FA = Id_{\mathbb{D}}FA = (F; Id_{\mathbb{D}})A$  and if  $f \in \mathbb{C}(A, B)$  then  $(Id_{\mathbb{C}}; F)f = FId_{\mathbb{C}}f = Ff = Id_{\mathbb{D}}Ff = (F; Id_{\mathbb{D}})f$ .

If  $\mathbb{C}$ ,  $\mathbb{D}$  and  $\mathbb{E}$  are three (small) categories, F is a functor from  $\mathbb{C}$  to  $\mathbb{D}$  and G is a functor from  $\mathbb{D}$  to  $\mathbb{E}$ , let A be an object of  $\mathbb{C}$ , we have ((F;G);H)A = H(F;G)A = HGFA = (G;H)FA = (F;(G;H))A and if  $f \in \mathbb{C}(A, B)$  then ((F;G);H)f = H(F;G)f = HGFf = (G;H)Ff = (F;(G;H))f.

# Example 9

If  $(A, B) \in obj(\mathbb{C} \times \mathbb{D})$ ,  $Pid_{(A,B)} = P(id_A, id_B) = id_A = id_{P(A,B)}$ . If  $(f,g) \in \mathbb{C} \times \mathbb{D}((A,B), (A',B'))$  and  $(f',g') \in \mathbb{C} \times \mathbb{D}((A',B'), (A'',B''))$ , P((f,g); (f',g')) = P(f;f',g;g') = f;f' = (P(f,g)); (P(f',g')). If  $\mathbb{D}$  has at least one membrism between any two objects let P and P' be two objects of  $\mathbb{D}$  and

If  $\mathbb{D}$  has at least one morphism between any two objects, let B and B' be two objects of  $\mathbb{D}$  and  $g \in \mathbb{D}(B, B')$ , for any  $f \in \mathbb{C}(A, A') = \mathbb{C}(P(A, B), P(A', B'))$ , we have P(f, g) = f.

# Example 10

If A is an object of  $\mathbb{C}$ ,  $id_{FA} \in \mathbb{D}(FA, FA)$  is an isomorphism (it is its own inverse). If  $f \in \mathbb{C}(A, B)$ , Ff;  $id_{FA} = Ff = id_{FA}$ ; Ff.

# **Definition 19**

If  $f \in \mathbb{C}(A, B)$ , Ff;  $(\alpha; {}^{1}\beta)_{B} = Ff$ ;  $\alpha_{B}; \beta_{B} = \alpha_{A}; Gf; \beta_{B} = \alpha_{A}; \beta_{A}; Hf = (\alpha; {}^{1}\beta)_{A}; Hf$ .

# **Definition 20**

Since  $\beta$  is a natural transformation from G to G', we have  $G\alpha_A$ ;  $\beta_{F'A} = \beta_{FA}$ ;  $G'\alpha_A$ . If  $f \in \mathbb{C}(A, B)$ , (F; G)f;  $(\alpha; {}^0\beta)_B = GFf$ ;  $G\alpha_B$ ;  $\beta_{F'B} = G(Ff; \alpha_B)$ ;  $\beta_{F'B} = G(\alpha_A; F'f)$ ;  $\beta_{F'B} = G\alpha_A$ ; GF'f;  $\beta_{F'B} = G\alpha_A$ ;  $\beta_{F'A}$ ;  $G'F'f = (\alpha; {}^0\beta)_A$ ; (F'; G')f.

# **Comment Page 9**

For any two objects A and B, we have a product  $A \times B$ . If  $f \in \mathbb{C}(A, B)$  and  $f' \in \mathbb{C}(A', B')$ , we define  $f \times f' = \langle \pi_A; f, \pi_{A'}; f' \rangle \in \mathbb{C}(A \times A', B \times B')$ . We have  $id_A \times id_{A'} = \langle \pi_A; id_A, \pi_{A'}; id_{A'} \rangle = \langle \pi_A, \pi_{A'} \rangle = id_{A \times A'}$  (using Fact 1). If  $f \in \mathbb{C}(A, B)$ ,  $g \in \mathbb{C}(B, C)$ ,  $f' \in \mathbb{C}(A', B')$  and  $g' \in \mathbb{C}(B', C')$ , we have, using Fact 2,  $(f \times f')$ ;  $(g \times g') = \langle \pi_A; f, \pi_{A'}; f' \rangle; \langle \pi_B; g, \pi_{B'}; g' \rangle = \langle \langle \pi_A; f, \pi_{A'}; f' \rangle; \pi_B; g, \langle \pi_A; f, \pi_{A'}; f' \rangle; \pi_{B'}; g' \rangle = \langle \pi_A; f; g, \pi_{A'}; f'; g' \rangle = (f; g) \times (f'; g')$ If  $f \in \mathbb{C}(A, B)$ , using Fact 2,  $f; \Delta_B = f; \langle id_B, id_B \rangle = \langle f; id_B, f; id_B \rangle = \langle f, f \rangle = \langle id_A; f, id_A; f \rangle = \langle \langle id_A, id_A \rangle; \pi_A^r; f \rangle = \langle id_A, id_A \rangle; \langle \pi_A^r; f, \pi_A^r; f \rangle = \Delta_A; (f \times f).$ 

# Example 12

If  $f: C \to A$  and  $g: C \to B$ , we define:

$$\langle f, g \rangle : C \to A \times B$$
  
 $x \mapsto (f(x), g(x))$ 

For all  $x \in C$ , we have  $\pi_1 \circ \langle f, g \rangle(x) = f(x)$  and  $\pi_2 \circ \langle f, g \rangle(x) = g(x)$ . Let  $h: C \to A \times B$  be such that any  $x \in C$ ,  $\pi_1 \circ h(x) = f(x)$  and  $\pi_2 \circ h(x) = g(x)$  then  $h(x) = (f(x), g(x)) = \langle f, g \rangle(x)$  that is  $h = \langle f, g \rangle$ .

For any set C, there is a unique function from C to  $\{\star\}$  defined by:

$$t_C: C \to \{\star\}$$
$$x \mapsto \star$$

If  $f : A \to C$  and  $g : B \to C$ , we define:

$$\begin{split} [f,g] &: A \uplus B \to C \\ & (0,a) \mapsto f(a) & \text{if } a \in A \\ & (1,b) \mapsto g(b) & \text{if } b \in B \end{split}$$

For any  $a \in A$ ,  $[f,g] \circ \iota_1(a) = f(a)$  and for any  $b \in B$ ,  $[f,g] \circ \iota_2(b) = g(b)$ . Let  $h: A \uplus B \to C$ be such that for any  $a \in A$ ,  $h \circ \iota_1(a) = f(a)$  and for any  $b \in B$ ,  $h \circ \iota_2(b) = g(b)$ , we have for any  $z \in A \uplus B$ , h(z) = [f,g](z) that is h = [f,g].

For any set C, there is a unique function from  $\emptyset$  to C which is the empty function.

### Example 13

If  $F : \mathbb{E} \to \mathbb{C}$  and  $G : \mathbb{E} \to \mathbb{D}$  are two functors, we define:

 $\begin{array}{ll} \langle F,G\rangle:\mathbb{E}\to\mathbb{C}\times\mathbb{D}\\ E\mapsto(FE,GE) & \text{ for objects of }\mathbb{E}\\ f\mapsto(Ff,Gf) & \text{ for morphisms of }\mathbb{E} \end{array}$ 

For any object E of  $\mathbb{E}$ , we have  $P_{\mathbb{C}}\langle F, G \rangle E = FE$  and  $P_{\mathbb{D}}\langle F, G \rangle E = GE$ . For any morphism f of  $\mathbb{E}$ , we have  $P_{\mathbb{C}}\langle F, G \rangle f = Ff$  and  $P_{\mathbb{D}}\langle F, G \rangle f = Gf$ . Let H be a functor from  $\mathbb{E}$  to  $\mathbb{C} \times \mathbb{D}$  such that  $P_{\mathbb{C}}HE = FE$ ,  $P_{\mathbb{D}}HE = GE$ ,  $P_{\mathbb{C}}Hf = Ff$  and  $P_{\mathbb{D}}Hf = Gf$  for any object E and any morphism f of  $\mathbb{E}$ , then  $HE = (FE, GE) = \langle F, G \rangle E$  and  $Hf = (Ff, Gf) = \langle F, G \rangle f$  that is  $H = \langle F, G \rangle$ . Let  $\mathbb{E}$  be a category, the unique functor  $T_{\mathbb{E}}$  from  $\mathbb{E}$  to  $\mathbb{T}$  is defined by  $T_{\mathbb{E}}E = \star$  for any object E of  $\mathbb{E}$  and  $T_{\mathbb{E}}f = id_{\star}$  for any morphism f of  $\mathbb{E}$ .

# Example 14

If  $F : \mathbb{C} \to \mathbb{E}$  and  $G : \mathbb{D} \to \mathbb{E}$  are two functors, we define:

$$\begin{split} [F,G] &: \mathbb{C} + \mathbb{D} \to \mathbb{E} \\ & (0,C) \mapsto FC & \text{if } C \in obj(\mathbb{C}) \\ & (1,D) \mapsto GD & \text{if } D \in obj(\mathbb{D}) \\ & f \mapsto Ff & \text{if } f \text{ morphism for } \mathbb{C} \\ & g \mapsto Gg & \text{if } g \text{ morphism for } \mathbb{D} \end{split}$$

For any  $C \in obj(\mathbb{C})$ ,  $[F,G]I_{\mathbb{C}}C = FC$  and for any  $B \in obj(\mathbb{D})$ ,  $[F,G]I_{\mathbb{D}}D = GD$ . For any fmorphism in  $\mathbb{C}$ ,  $[F,G]I_{\mathbb{C}}f = Ff$  and for any g morphism in  $\mathbb{D}$ ,  $[F,G]I_{\mathbb{D}}g = Gg$ . Let  $H : \mathbb{C} + \mathbb{D} \to \mathbb{E}$ be a functor such that for any  $C \in obj(\mathbb{C})$ ,  $HI_{\mathbb{C}}C = FC$ , for any  $B \in obj(\mathbb{D})$ ,  $HI_{\mathbb{D}}D = GD$ , for any f morphism in  $\mathbb{C}$ ,  $HI_{\mathbb{C}}f = Ff$  and for any g morphism in  $\mathbb{D}$ ,  $HI_{\mathbb{D}}g = Gg$ , we have for any object A and for any morphism h of  $\mathbb{C} + \mathbb{D}$ , HA = [F,G]A and Hh = [F,G]h, that is H = [F,G]. Let  $\mathbb{E}$  be a category, the empty functor is the unique functor from  $\bot$  to  $\mathbb{E}$ .

### **Definition 26**

Let (E, e) be an equalizer of  $f \in \mathbb{C}(A, B)$  and  $g \in \mathbb{C}(A, B)$ , if f' and g' are in  $\mathbb{C}(D, E)$  such that f' ; e = g' ; e then f' ; e ; f = g' ; e ; f = g' ; e ; g thus there exists a unique  $h \in \mathbb{C}(D, E)$  such that f' ; e = g' ; e = h ; e so that f' = h = g'.

# Comment Page 12

The following diagram commutes:



by:

- (a) triangle of monoidal categories
- (b) **Fact** 4
- (c) Fact 5

We thus have  $u_1^l \otimes 1 = u_1^r \otimes 1$  since  $a_{1,1,1}$  is an isomorphism, and finally  $u_1^l = u_1^r$  by Fact 3.

# Comment Page 13

Thanks to Fact 3, it is sufficient to prove the commutation of the following diagram (since s and a are isomorphisms):



which commutes by:

- (a) triangle of monoidal categories
- (b) naturality of s
- (c) naturality of  $u^r$

- (d) Fact 5
- (e) Fact 5
- (f) hexagon of symmetric monoidal categories

# Example 15

× is a bi-functor from  $\mathbb{C}$  and  $\mathbb{C}$  to  $\mathbb{C}$  (see page 9). We consider three morphisms  $f \in \mathbb{C}(A, A'), g \in \mathbb{C}(B, B')$  and  $h \in \mathbb{C}(C, C')$ . We have:

• using Fact 2 and the definition of the bi-functor  $\times$ :

$$(f \times g) \times h; \langle \pi_{A' \times B'}; \pi_{A'}, \langle \pi_{A' \times B'}; \pi_{B'}, \pi_{C'} \rangle \rangle$$
  
=  $\langle (f \times g) \times h; \pi_{A' \times B'}; \pi_{A'}, (f \times g) \times h; \langle \pi_{A' \times B'}; \pi_{B'}, \pi_{C'} \rangle \rangle$   
=  $\langle \pi_{A \times B}; \pi_{A}; f, (f \times g) \times h; \langle \pi_{A' \times B'}; \pi_{B'}, \pi_{C'} \rangle \rangle$   
=  $\langle \pi_{A \times B}; \pi_{A}; f, \langle (f \times g) \times h; \pi_{A' \times B'}; \pi_{B'}, (f \times g) \times h; \pi_{C'} \rangle \rangle$   
=  $\langle \pi_{A \times B}; \pi_{A}; f, \langle \pi_{A \times B}; \pi_{B}; g, \pi_{C}; h \rangle \rangle$ 

and

$$\begin{split} \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; f \times (g \times h) \\ &= \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \langle \pi_A ; f, \pi_{B \times C} ; g \times h \rangle \\ &= \langle \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_A ; f, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_{B \times C} ; g \times h \rangle \\ &= \langle \pi_{A \times B} ; \pi_A ; f, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle ; g \times h \rangle \\ &= \langle \pi_{A \times B} ; \pi_A ; f, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle ; \langle \pi_B ; g, \pi_C ; h \rangle \rangle \\ &= \langle \pi_{A \times B} ; \pi_A ; f, \langle \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle ; \pi_B ; g, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle ; \pi_C ; h \rangle \rangle \\ &= \langle \pi_{A \times B} ; \pi_A ; f, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle ; \pi_B ; g, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle ; \pi_C ; h \rangle \rangle$$

Moreover, with Fact 1 and Fact 2:

$$\begin{split} \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle \\ &= \langle \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_{B \times C} ; \pi_C \rangle \\ &= \langle \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_C \rangle \\ &= \langle \langle \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_A, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_{B \times C} ; \pi_B \rangle, \pi_C \rangle \\ &= \langle \langle \pi_{A \times B} ; \pi_A, \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_A, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_{B \times C} ; \pi_B \rangle, \pi_C \rangle \\ &= \langle \pi_{A \times B} ; \pi_A, \pi_{A \times B} ; \pi_B \rangle, \pi_C \rangle \\ &= \langle \pi_{A \times B} ; \langle \pi_A, \pi_B \rangle, \pi_C \rangle \\ &= \langle \pi_{A \times B} ; \langle \pi_A, \pi_B \rangle, \pi_C \rangle \\ &= \langle \pi_{A \times B}, \pi_C \rangle \end{split}$$

 $= id_{(A \times B) \times C}$ 

and

$$\begin{aligned} \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle &; \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \\ &= \langle \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle ; \pi_{A \times B} ; \pi_A, \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \\ &= \langle \pi_A, \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \\ &= \langle \pi_A, \langle \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle ; \pi_{A \times B} ; \pi_B, \langle \langle \pi_A, \pi_{B \times C} ; \pi_B \rangle, \pi_{B \times C} ; \pi_C \rangle ; \pi_C \rangle \rangle \\ &= \langle \pi_A, \langle \pi_{B \times C} ; \pi_B, \pi_{B \times C} ; \pi_C \rangle \\ &= \langle \pi_A, \pi_{B \times C} ; \pi_B, \pi_{B \times C} ; \pi_C \rangle \rangle \\ &= \langle \pi_A, \pi_{B \times C} ; \langle \pi_B, \pi_C \rangle \rangle \\ &= \langle \pi_A, \pi_{B \times C} ; \langle \pi_B, \pi_C \rangle \rangle \\ &= \langle \pi_A, \pi_{B \times C} \rangle \\ &= id_{A \times (B \times C)} \end{aligned}$$

• We first prove that  $\pi_A \in \mathbb{C}(A \times \top, A)$  is the inverse of  $\langle id_A, t_A \rangle \in \mathbb{C}(A, A \times \top)$  using Fact 1 and Fact 2:

$$\langle id_A, t_A \rangle; \pi_A = id_A$$

and

$$\begin{aligned} \pi_A ; \langle id_A, t_A \rangle &= \langle \pi_A ; id_A, \pi_A ; t_A \rangle \\ &= \langle \pi_A, \pi_\top \rangle \\ &= id_{A \times \top} \end{aligned}$$

We also have:

$$\begin{split} \langle id_A, t_A \rangle \, ; \, f \times id_{\top} &= \langle id_A, t_A \rangle \, ; \, \langle \pi_A \; ; \, f, \pi_{\top} \; ; \; id_{\top} \rangle \\ &= \langle \langle id_A, t_A \rangle \; ; \, \pi_A \; ; \, f, \langle id_A, t_A \rangle \; ; \, \pi_{\top} \; ; \; id_{\top} \rangle \\ &= \langle f, t_A \rangle \\ &= \langle f \; ; \; id_{A'}, f \; ; \; t_{A'} \rangle \\ &= f \; ; \; \langle id_{A'}, t_{A'} \rangle \end{split}$$

- The results for  $\langle t_A, id_A \rangle$  are very similar.
- Using Fact 2:

$$f \times g; \langle \pi_{B'}, \pi_{A'} \rangle = \langle f \times g; \pi_{B'}, f \times g; \pi_{A'} \rangle$$
$$= \langle \pi_B; g, \pi_A; f \rangle$$

and

$$\begin{aligned} \langle \pi_B, \pi_A \rangle \, ; \, g \times f &= \langle \pi_B, \pi_A \rangle \, ; \, \langle \pi_B \, ; \, g, \pi_A \, ; \, f \rangle \\ &= \langle \langle \pi_B, \pi_A \rangle \, ; \, \pi_B \, ; \, g, \langle \pi_B, \pi_A \rangle \, ; \, \pi_A \, ; \, f \rangle \\ &= \langle \pi_B \, ; \, g, \pi_A \, ; \, f \rangle \end{aligned}$$

Moreover, with Fact 1 and Fact 2:

$$\begin{aligned} \langle \pi_B, \pi_A \rangle ; \langle \pi_A, \pi_B \rangle &= \langle \langle \pi_B, \pi_A \rangle ; \pi_A, \langle \pi_B, \pi_A \rangle ; \pi_B \rangle \\ &= \langle \pi_A, \pi_B \rangle \\ &= id_{A \times B} \end{aligned}$$

We now have to prove to the three additional commutative diagrams of symmetric monoidal categories.

• Pentagon of monoidal categories:

$$\langle \pi_{(A\times B)\times C} ; \pi_{A\times B}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle ; \langle \pi_{A\times B} ; \pi_{A}, \langle \pi_{A\times B} ; \pi_{B}, \pi_{C\times D} \rangle \rangle$$

$$= \langle \langle \pi_{(A\times B)\times C} ; \pi_{A\times B}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle ; \pi_{A\times B} ; \pi_{A}, \langle \pi_{(A\times B)\times C} ; \pi_{A\times B}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle ; \langle \pi_{A\times B} ; \pi_{A}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle ; \langle \pi_{A\times B} ; \pi_{A}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle ; \langle \pi_{A\times B} ; \pi_{A}, \langle \pi_{(A\times B)\times C} ; \pi_{A\times B} ; \pi_{A}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle ; \pi_{A\times B} ; \pi_{B}, \langle \pi_{(A\times B)\times C} ; \pi_{A\times B}, \langle \pi_{(A\times B)\times C} ; \pi_{A\times B}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle$$

$$= \langle \pi_{(A\times B)\times C} ; \pi_{A\times B} ; \pi_{A}, \langle \pi_{(A\times B)\times C} ; \pi_{A\times B} ; \pi_{B}, \langle \pi_{(A\times B)\times C} ; \pi_{C}, \pi_{D} \rangle \rangle \rangle$$

and

 $\langle \pi_{A\times B} ; \pi_A, \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \langle \pi_{A\times (B\times C)} ; \pi_A, \langle \pi_{A\times (B\times C)} ; \pi_{B\times C}, \pi_D \rangle \rangle ; id_A \times \langle \pi_{B\times C} ; \pi_B, \langle \pi_{B\times C} ; \pi_C, \pi_D \rangle \rangle$ 

- $= \langle \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \pi_{A \times (B \times C)} ; \pi_A, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \langle \pi_{A \times (B \times C)} ; \pi_B \rangle \rangle \times id_D ; \langle \pi_{A \times (B \times C)} ; \pi_B \rangle \rangle \times id_D ; \langle \pi_{A \times (B \times C)} ; \pi_B \rangle \rangle \rangle \times id_D ; \langle \pi_{A \times (B \times C)} ; \pi_B \rangle \rangle \rangle \times id_D ; \langle \pi_{A \times (B \times C)} ; \pi_B \rangle \rangle \rangle \rangle = 0$
- $= \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_A, \langle \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \pi_{A \times (B \times C)} ; \pi_{B \times C}, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \pi_{A \times (B \times C)} ; \pi_{B \times C}, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \pi_{A \times (B \times C)} ; \pi_{B \times C}, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \pi_{A \times (B \times C)} ; \pi_{B \times C}, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle \times id_D ; \pi_{A \times (B \times C)} ; \pi_{B \times C}, \langle \pi_{A \times B} ; \pi_A, \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle$
- $= \langle \pi_{(A \times B) \times C} \ ; \ \pi_{A \times B} \ ; \ \pi_{A}, \langle \pi_{(A \times B) \times C} \ ; \ \langle \pi_{A \times B} \ ; \ \pi_{B}, \ \pi_{C} \rangle, \ \pi_{D} \rangle \rangle \ ; \ \langle \pi_{A}, \ \pi_{(B \times C) \times D} \ ; \ \langle \pi_{B \times C} \ ; \ \pi_{B}, \ \langle \pi_{B \times C} \ ; \ \pi_{C}, \ \pi_{D} \rangle \rangle \rangle$
- $= \left\langle \left\langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{B}, \pi_{C} \right\rangle, \pi_{D} \right\rangle \right\rangle; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A \times B} ; \pi_{A}, \left\langle \pi_{(A \times B) \times C} ; \left\langle \pi_{A} \right\rangle \right\rangle \right\rangle \right\rangle\right\rangle\right\rangle$
- $= \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_A, \langle \pi_{(A \times B) \times C} ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{B \times C} ; \pi_B, \langle \pi_{B \times C} ; \pi_C, \pi_D \rangle \rangle \rangle$
- $= \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_A, \langle \langle \pi_{(A \times B) \times C} ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \pi_{B \times C} ; \pi_B, \langle \pi_{(A \times B) \times C} ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \langle \pi_A ; \pi_A ; \pi_A ; \pi_A \rangle ; \langle \pi_A ; \pi_A ; \pi_A ; \pi_A ; \pi_A ; \pi_A \rangle ; \langle \pi_A ; \pi_A$
- $= \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_A, \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_B, \langle \langle \pi_{(A \times B) \times C} ; \langle \pi_{A \times B} ; \pi_B, \pi_C \rangle, \pi_D \rangle ; \pi_{B \times C} ; \pi_C, \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_B, \pi_C \rangle \rangle$
- $= \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_A, \langle \pi_{(A \times B) \times C} ; \pi_{A \times B} ; \pi_B, \langle \pi_{(A \times B) \times C} ; \pi_C, \pi_D \rangle \rangle \rangle$
- Triangle of monoidal categories:

$$\begin{split} \langle id_A, t_A \rangle &\times id_B ; \langle \pi_{A \times \top} ; \pi_A, \langle \pi_{A \times \top} ; \pi_{\top}, \pi_B \rangle \rangle \\ &= \langle \langle id_A, t_A \rangle \times id_B ; \pi_{A \times \top} ; \pi_A, \langle id_A, t_A \rangle \times id_B ; \langle \pi_{A \times \top} ; \pi_{\top}, \pi_B \rangle \rangle \\ &= \langle \pi_A, \langle \langle id_A, t_A \rangle \times id_B ; \pi_{A \times \top} ; \pi_{\top}, \langle id_A, t_A \rangle \times id_B ; \pi_B \rangle \rangle \\ &= \langle \pi_A, \langle \pi_A ; t_A, \pi_B \rangle \rangle \\ &= \langle \pi_A, \langle t_{A \times B}, \pi_B \rangle \rangle \\ &= \langle \pi_A, \langle \pi_B ; t_B, \pi_B \rangle \rangle \\ &= \langle \pi_A, \pi_B ; \langle t_B, id_B \rangle \rangle \\ &= id_A \times \langle t_B, id_B \rangle \end{split}$$

• Hexagon of symmetric monoidal categories:

$$\begin{aligned} \langle \pi_{A\times B} ; \pi_A, \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle \rangle ; \langle \pi_{B\times C}, \pi_A \rangle ; \langle \pi_{B\times C} ; \pi_B, \langle \pi_{B\times C} ; \pi_C, \pi_A \rangle \rangle \\ &= \langle \langle \pi_{A\times B} ; \pi_A, \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_{B\times C}, \langle \pi_{A\times B} ; \pi_A, \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle \rangle ; \pi_A \rangle ; \langle \pi_{B\times C} ; \pi_B, \langle \pi_{B\times C} ; \pi_C, \pi_A \rangle \rangle \\ &= \langle \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle, \pi_{A\times B} ; \pi_A \rangle ; \langle \pi_{B\times C} ; \pi_B, \langle \pi_{B\times C} ; \pi_C, \pi_A \rangle \rangle \\ &= \langle \pi_{A\times B} ; \pi_B, \langle \langle \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle, \pi_{A\times B} ; \pi_A \rangle ; \pi_{B\times C} ; \pi_C, \pi_A \rangle \rangle \\ &= \langle \pi_{A\times B} ; \pi_B, \langle \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle, \pi_{A\times B} ; \pi_A \rangle ; \pi_{B\times C} ; \pi_C, \langle \langle \pi_{A\times B} ; \pi_B, \pi_C \rangle, \pi_{A\times B} ; \pi_A \rangle ; \pi_A \rangle \rangle$$

and

$$\begin{split} \langle \pi_B, \pi_A \rangle &\times id_C ; \langle \pi_{B \times A} ; \pi_B, \langle \pi_{B \times A} ; \pi_A, \pi_C \rangle \rangle ; id_B \times \langle \pi_C, \pi_A \rangle \\ &= \langle \langle \pi_B, \pi_A \rangle \times id_C ; \pi_{B \times A} ; \pi_B, \langle \pi_B, \pi_A \rangle \times id_C ; \langle \pi_{B \times A} ; \pi_A, \pi_C \rangle \rangle ; id_B \times \langle \pi_C, \pi_A \rangle \\ &= \langle \pi_{A \times B} ; \pi_B, \langle \langle \pi_B, \pi_A \rangle \times id_C ; \pi_{B \times A} ; \pi_A, \langle \pi_B, \pi_A \rangle \times id_C ; \pi_C \rangle \rangle ; id_B \times \langle \pi_C, \pi_A \rangle \\ &= \langle \pi_{A \times B} ; \pi_B, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle \rangle ; \langle \pi_B, \pi_{A \times C} ; \langle \pi_C, \pi_A \rangle \rangle \\ &= \langle \langle \pi_{A \times B} ; \pi_B, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle \rangle ; \pi_B, \langle \pi_{A \times B} ; \pi_B, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle \rangle ; \pi_{A \times C} ; \langle \pi_C, \pi_A \rangle \rangle \\ &= \langle \pi_{A \times B} ; \pi_B, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle ; \langle \pi_C, \pi_A \rangle \rangle \\ &= \langle \pi_{A \times B} ; \pi_B, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle ; \pi_C, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle ; \pi_A \rangle \rangle \\ &= \langle \pi_{A \times B} ; \pi_B, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle ; \pi_C, \langle \pi_{A \times B} ; \pi_A, \pi_C \rangle ; \pi_A \rangle \rangle \\ &= \langle \pi_{A \times B} ; \pi_B, \langle \pi_A \times B ; \pi_A, \pi_C \rangle ; \pi_C, \langle \pi_A \times B ; \pi_A, \pi_C \rangle ; \pi_A \rangle \rangle$$

# **Property 3**

The diagram:



commutes by:

- (a) functoriality of  $\boxtimes$
- (b) hexagon of monoidal functors
- (c) functoriality of  $\boxtimes$
- (d) naturality of m
- (e) pentagon of monoids
- (f) naturality of m

The diagram:



commutes by:

- (a) square of monoidal functors
- (b) naturality of m
- (c) triangle of monoids

The diagram:



commutes by:

- (a) square of monoidal functors
- (b) naturality of m
- (c) triangle of monoids

In the case of a symmetric monoidal functor and a symmetric monoid, the diagram:





- (a) square of symmetric monoidal functors
- (b) triangle of symmetric monoids