Subject 2: Linear Sequent Calculus

to be returned on Friday, October 2nd

In the whole subject, exchange rules can be left implicit.

MLL

Formulas are given by:

$$A ::= X \mid X^{\perp} \mid A \otimes A \mid A \mathcal{R} A \mid 1 \mid \bot$$

where X ranges over the elements of a given countable set \mathcal{V} of variables.

We consider the following rules for the one-sided multiplicative linear sequent calculus MLL:

$$\frac{-\Gamma}{\vdash A, A^{\perp}} ax \qquad \frac{\vdash \Gamma, A \vdash \Delta, A^{\perp}}{\vdash \Gamma, \Delta} cut \qquad \frac{\vdash \Gamma}{\vdash \sigma(\Gamma)} ex$$

$$\frac{\vdash \Gamma, A \vdash \Delta, B}{\vdash \Gamma, \Delta, A \otimes B} \otimes \qquad \frac{\vdash \Gamma, A, B}{\vdash \Gamma, A ? B} ? ? \qquad \frac{\vdash \Gamma}{\vdash \Gamma, \bot} \bot$$

Question 1. For each sequent below, if it is provable give a proof in one-sided MLL, and if it is not provable try to give a short justification.

a.
$$\vdash \bot, X^{\bot} \Im X$$

b.
$$\vdash X \otimes Y, Y^{\perp} ? ? (1 \otimes X^{\perp})$$

$$\mathbf{c.} \vdash X \otimes X, X^{\perp} \otimes X^{\perp}$$

d.
$$\vdash X^{\perp}, X \otimes (X^{\perp} \Im X)$$

e.
$$\vdash Y \ \mathcal{F} \ X^{\perp}, \perp \otimes \perp, X, Y^{\perp}$$

$$\mathbf{f.} \; \vdash X \otimes (Y \, \mathfrak{P} \, X^{\perp}), X^{\perp}, X$$

$$\mathbf{g.} \ \vdash X^{\perp} \ \Im \ (Y^{\perp} \otimes Z^{\perp}), (X \otimes Y) \ \Im \ Z$$

$$\mathbf{h.} \; \vdash X^{\perp} \otimes X, (X \; \mathfrak{P} \; X^{\perp}) \otimes (X \; \mathfrak{P} \; X^{\perp})$$

i.
$$\vdash X \, \mathfrak{P} \, X^{\perp}, (X^{\perp} \otimes X) \, \mathfrak{P} \, (X^{\perp} \otimes X)$$

Question 2. Prove the following facts about MLL:

- **a.** $\vdash A \otimes B$ is provable if and only if both $\vdash A$ and $\vdash B$ are provable.
- **b.** If $\vdash 1, A \otimes B$ is provable then either $\vdash A$ or $\vdash B$ is provable.
- **c.** If $\vdash A \otimes B, C \otimes D$ is provable then $\vdash A$ or $\vdash B$ or $\vdash C$ or $\vdash D$ is provable.

- **d.** Let $\sharp_1(\Gamma)$ be the number of occurrences of 1 in Γ , $\sharp_{\otimes}(\Gamma)$ be the number of occurrences of \otimes in Γ , $\sharp_{\mathcal{V}}(\Gamma)$ be the number of occurrences of elements of \mathcal{V} not below a $_{-}^{\perp}$ in Γ , prove that $\sharp_1(\Gamma) + \sharp_{\mathcal{V}}(\Gamma) = 1 + \sharp_{\otimes}(\Gamma)$ for all Γ such that $\vdash \Gamma$ is provable.
- **e.** Assuming that $\vdash A$ is provable, show $\vdash A$, A is not provable.

Question 3. Define the formulas and rules of two-sided MLL (do not forget to be careful with negation).

MALL

Formulas are given by:

$$A ::= X \mid X^{\perp} \mid A \otimes A \mid A \Re A \mid 1 \mid \bot \mid A \& A \mid A \oplus A \mid \top \mid 0$$

where X ranges over the elements of a given countable set $\mathcal V$ of variables.

We consider the following rules for the one-sided multiplicative-additive linear sequent calculus MALL:

Question 4. For each sequent below, if it is provable give a proof in one-sided MALL, and if it is not provable try to give a short justification.

a.
$$\vdash X, X \oplus X^{\perp}$$

b.
$$\vdash X^{\perp}, X \oplus X$$

c.
$$\vdash X^{\perp}, X \& X$$

d.
$$\vdash X^{\perp}, X \& X^{\perp}$$

e.
$$\vdash X^{\perp}, X \& (X \oplus Y)$$

$$\mathbf{f.} \vdash X. X^{\perp} \oplus (X^{\perp} \& Y^{\perp})$$

$$\mathbf{g.} \vdash 0, \top \Im X$$

h.
$$\vdash X^{\perp} \& (Y^{\perp} \oplus Z^{\perp}), (X \oplus Y) \& (X \oplus Z)$$

i.
$$\vdash X^{\perp} \Re (Y^{\perp} \oplus Z^{\perp}), (X \otimes Y) \& (X \otimes Z)$$

$$\mathbf{j.} \vdash X^{\perp} \, \mathfrak{P} \, (Y^{\perp} \, \& \, Z^{\perp}), (X \otimes Y) \oplus (X \otimes Z)$$

$$\mathbf{k.} \vdash (X^{\perp} \Im Y^{\perp}) \oplus (X^{\perp} \Im Z^{\perp}), X \otimes (Y \& Z)$$

1.
$$\vdash (X^{\perp} ? Y^{\perp}) & (X^{\perp} ? Z^{\perp}), X \otimes (Y \oplus Z)$$