

BOUNDING LINEAR HEAD REDUCTION AND VISIBLE INTERACTION THROUGH SKELETONS

PIERRE CLAIRAMBAULT

CNRS, ENS de Lyon, Inria, UCBL, Université de Lyon
e-mail address: pierre.clairambault@ens-lyon.fr

ABSTRACT. In this paper, we study the complexity of execution in higher-order programming languages. Our study has two facets: on the one hand we give an upper bound to the length of interactions between bounded P-visible strategies in Hyland-Ong game semantics. This result covers models of programming languages with access to computational effects like non-determinism, state or control operators, but its semantic formulation causes a loose connection to syntax. On the other hand we give a syntactic counterpart of our semantic study: a non-elementary upper bound to the length of the linear head reduction sequence (a low-level notion of reduction, close to the actual implementation of the reduction of higher-order programs by abstract machines) of simply-typed λ -terms. In both cases our upper bounds are proved optimal by giving matching lower bounds.

These two results, although different in scope, are proved using the same method: we introduce a simple reduction on finite trees of natural numbers, hereby called *interaction skeletons*. We study this reduction and give upper bounds to its complexity. We then apply this study by giving two simulation results: a semantic one measuring progress in game-theoretic interaction via interaction skeletons, and a syntactic one establishing a correspondence between linear head reduction of terms satisfying a locality condition called *local scope* and the reduction of interaction skeletons. This result is then generalized to arbitrary terms by a local scopization transformation.

1. INTRODUCTION

In the last two decades there has been a significant interest in the study of *quantitative* or *intensional* aspects of higher-order programs; in particular, the study of their *complexity* has generated a lot of effort. In the context of the λ -calculus, the first result that comes to mind is the work by Schwichtenberg [19], later improved by Beckmann [4], establishing upper bounds to the length of β -reduction sequences for simply-typed λ -terms. In the related line of work of *implicit complexity*, type systems have been developed to characterize extensionally certain classes of functions, such as polynomial [13] or elementary [11] time. Such systems rely on a soundness theorem establishing that well-typed terms normalize in a

1998 ACM Subject Classification: F.3.2.

Key words and phrases: Linear head reduction, complexity, game semantics.

This work presents the combined results of two conference papers [6, 7].

This work was partially supported by the ERC Advanced Grant ECSYM and by the French ANR grant ELICA, ANR-14-CE25-0005.

certain restricted time, which is itself established using syntactic methods that are specific to the system being studied. This calls for the development of syntax-independent tools to study precisely the execution time of higher-order programs. The present paper is a contribution towards that goal.

In order to develop tools for estimating the complexity of programs that are at the same time syntax-independent but still precise and computationally relevant, game semantics is a good place to start. Indeed Hyland-Ong game semantics, originally introduced to solve the full abstraction problem for PCF [15], have since proved powerful and flexible enough to provide fully abstract models for various computational features of programming languages: non-determinism, state, exceptions, control operators, concurrency... At the same time game semantics are computationally informative: moves in an interaction exactly correspond to computation steps as performed by some abstract machines, as was formally proved by Danos, Herbelin and Regnier [9] for the simply-typed λ -calculus. Accordingly, the first contribution of this paper is an upper bound to the length of interactions between bounded P-visible strategies in Hyland-Ong game semantics. Bounded P-visible strategies allow computational effects such as non-determinism, control operators or ground state but disallow, for instance, higher-order references using which a fixpoint combinator can be defined. In this context, we give a non-elementary upper bound to the length of interactions (so implicitly, computation), for which we prove optimality by providing a matching lower bound. This is done by extracting from interaction sequences a finite tree of natural numbers called an *interaction skeleton*. We then prove a simulation result, showing that progress in the game theoretic interaction is witnessed by a simple reduction on the corresponding interaction skeletons. This reduction is then analyzed independently of its connection with game semantics, and this analysis informs a bound on the length of P-visible interactions.

This syntax-independent approach advocated above yields a result with a broader scope, however there is a price to pay: the innocent strategies of game semantics are representations of η -long β -normal λ -terms, so their interactions are only easily linked to syntactic reduction of terms of the form $M N_1 \dots N_p$ where M and all the N_i s are β -normal η -long normal forms; we will call such terms *game situations* in this paper. In order to compensate for this structural restriction, we investigate the direct relationship between interaction skeletons and linear head reduction, independently of game semantics. We define a set of terms called *generalized game situations*, that are both η -long in a sense adapted to linear head reduction, and satisfy a key locality constraint called *local scope*. Generalized game situations are preserved by linear head reduction. Moreover they can be simulated by interaction skeletons, from which we deduce optimal upper bounds to their complexity. These bounds are then generalized to arbitrary simply-typed λ -terms by means of a syntactic λ -lifting [16] transformation putting arbitrary terms to locally scoped form.

Related works. There are multiple approaches to the complexity analysis of higher-order programs, but they seem to separate into two major families. On the one hand, Beckmann [4], extending earlier work by Schwichtenberg [19], gave exact bounds to the maximal length of β -reduction on simply-typed λ -terms. His analysis uses very basic information on the terms (their length, or height, and order), but gives bounds that are in general very rough. On the other hand other groups, including Dal Lago and Laurent [17], De Carvalho [12], or Bernardet and Lengrand [5], use semantic structures (respectively, game semantics, relational semantics, or non-idempotent intersection types) to capture abstractly the precise complexity of particular terms. Their bounds are much more precise on particular terms, but require information on the terms whose extraction is in general as long to obtain as

actual execution. The present contribution belongs to the first family. However, unlike Beckmann and Schwichtenberg our core tools are syntax-independent. Moreover we focus on *linear head reduction*, the notion of reduction implemented by several call-by-name abstract machines [10], closer to the actual execution of functional programming languages.

Outline. In Section 2, we introduce a few basic notions or notations useful to the rest of the paper. In Section 3 we present the game semantics framework from which interaction skeletons were originally extracted, and prove our semantic simulation result. Section 4 is a largely standalone section in which we study interaction skeletons and their reduction, and prove our main complexity result. Finally, Section 5 focuses on the syntactic implications of our study of skeletons and details their relationship with linear head reduction.

This paper is organized around the notion of interaction skeleton and their reduction, with two largely independent applications to game semantics and to the complexity of linear head reduction. We chose to present first the game-theoretic development, since it motivates the definition of interaction skeletons. However, our intention is that the paper should be accessible to semantically-minded as well as more syntactically-minded readers. In particular, readers not interested in game semantics should be able to skip Section 3 and still have everything needed to understand Sections 4 and 5.

2. PRELIMINARIES

2.1. Syntax and dynamics of the λ -calculus. In this paper, we consider the simply-typed λ -calculus Λ built from one base type o . Its **types** and **terms** are:

$$\begin{aligned} A, B &::= o \mid A \rightarrow B \\ M, N &::= \lambda x^A. M \mid M N \mid x \mid *_A \end{aligned}$$

subject to the usual typing rules defining the typing relation $\Gamma \vdash M : A$ (for definiteness, contexts Γ are considered to be sets of pairs $x : A$, where x is a variable name and A is a simple type). All the terms in this paper are considered well-typed, although we will not always make it explicit. Note that we work here with the simply-typed λ -calculus *à la Church*, *i.e.* the variables are explicitly annotated with types (although we often omit the annotations for the sake of readability). For each type A , there is a constant $*_A : A$ of type A . We will often omit the index and write $*$. As usual, we write $\text{fv}(M)$ for the set of *free variables* of a term M , *i.e.* variables appearing in the term but not bound by a λ . If A is a type, we write id_A for the term $\vdash \lambda x^A. x : A \rightarrow A$. Terms are assumed to obey Barendregt's convention, and are considered up to α -equivalence. Note that our design choices – only one atom, each type is inhabited – merely make the presentation simpler and are not strictly required for our results to hold.

If $\Gamma, x : A \vdash M : B$ and $\Gamma \vdash N : A$, we write $M[N/x]$ for the **substitution** of x by N in M , *i.e.* M where all occurrences of x have been replaced by N . Although this paper focuses on linear head reduction (to be defined later), we will occasionally need **β -reduction**. It is the usual notion of reduction in the λ -calculus, defined by the context-closure of:

$$(\lambda x. M) N \rightarrow_{\beta} M[N/x]$$

Likewise we consider **η -expansion** to be the context closure of:

$$M \rightarrow_{\eta} \lambda x^A. M x$$

valid whenever M has type $A \rightarrow B$ for some types A, B and $x \notin \text{fv}(M)$.

The **level** of a type is defined by $\text{lv}(o) = 0$ and $\text{lv}(A \rightarrow B) = \max(\text{lv}(A) + 1, \text{lv}(B))$. Likewise, the **level** $\text{lv}(M)$ of a term M is the level of its type. Finally, the **order** $\text{ord}(M)$ of a term M is the maximal $\text{lv}(N)$, for all subterms N of M . Within a term $\Gamma \vdash M : A$ such that $(x : B) \in \Gamma$, we write $\text{lv}_M(x) = \text{lv}(B)$. The term M will generally be implicit from the context, so we will just write $\text{lv}(x)$.

2.2. Growth rates of functions. We recall some standard notations for comparing growth rates of functions. For functions $f, g : \mathbb{N} \rightarrow \mathbb{N}$, we write $f(n) = \Theta(g(n))$ when there exists reals $c_1, c_2 > 0$ and $N \in \mathbb{N}$ such that for all $n \geq N$, $c_1 g(n) \leq f(n) \leq c_2 g(n)$. This is generalized to functions of multiple variables $f, g : \mathbb{N}^p \rightarrow \mathbb{N}$ by setting that $f(n_1, \dots, n_p) = \Theta(g(n_1, \dots, n_p))$ iff there are $c_1, c_2 > 0$ and $N \in \mathbb{N}$ such that for all $n_i \geq N$ we have $c_1 g(n_1, \dots, n_p) \leq f(n_1, \dots, n_p) \leq c_2 g(n_1, \dots, n_p)$. If $h : \mathbb{N} \rightarrow \mathbb{N}$ is another function, we write $f(n_1, \dots, n_p) = h(\Theta(g(n_1, \dots, n_p)))$ iff there is a function $\phi : \mathbb{N}^p \rightarrow \mathbb{N}$ such that $f(n_1, \dots, n_p) = h(\phi(n_1, \dots, n_p))$ and $\phi(n_1, \dots, n_p) = \Theta(g(n_1, \dots, n_p))$.

3. VISIBLE POINTER STRUCTURES AND INTERACTION SKELETONS

As we said before, the general purpose of this paper is to develop syntax-independent tools to reason about termination and complexity of programming languages. *Game semantics* provide such a framework: in this setting, programs are identified with the set of their possible interactions with the execution environment, presented as a *strategy*. All the syntactic information is forgotten but at the same time no dynamic information is lost, since the strategy exactly describes the behaviour of the term within any possible evaluation context.

More importantly, game semantics can be seen both as a *denotational semantics* and as an *operational semantics* in the sense that the interaction process at the heart of games is operationally informative and strongly related to the actual evaluation process as implemented, for instance, by abstract machines. This important intuition was made formal for the first time, to the author's knowledge, by Danos, Herbelin and Regnier in [9]. There, they showed that assuming one has a simply-typed λ -term $M N_1 \dots N_p$ where M and the N_i s are both η -long and β -normal, then there is a step-by-step correspondence between:

- The *linear head reduction sequence* of $M N_1 \dots N_p$,
- The *game-theoretic interaction* between strategies $\llbracket M \rrbracket$ and $\prod_{1 \leq i \leq p} \llbracket N_i \rrbracket$.

In this paper, we will refer to simply-typed λ -terms of that particular shape as **game situations**, because it is in those situations that the connection between game semantics and execution of programs is the most direct. Indeed (innocent) game semantics can be seen as a reduction-free way of composing Böhm trees, *i.e.* of computing game situations.

Game situations also provide the starting point of our present contributions. Indeed, the connection above reduces the termination and complexity analysis of the execution of a game situation $M N_1 \dots N_p$ to the syntax-independent analysis of the game-theoretic witness of this execution, *i.e.* the game-theoretic interaction between the strategies $\llbracket M \rrbracket$ and $\prod_{1 \leq i \leq p} \llbracket N_i \rrbracket$. More precisely, it turns out that from this interaction one just has to keep the structure of pointers in order to get a precise estimate of the complexity of execution.

In this section, we will start by recalling a few basic definitions of Hyland-Ong game semantics. We will show that the mechanism of composition gives rise to structure called *visible pointer structures*. We will then show the main result of this section, that visible pointer structures can be simulated by a simple rewriting system called *interaction skeletons*.

3.1. Brief reminder of Hyland-Ong games. We start this section by recalling some of the basic definitions of Hyland-Ong games. The presentation of game semantics will be intentionally brief and informal, firstly because it is only there to provide context and is not a prerequisite to understand the paper, and secondly because good introductions can be easily found in the literature, see *e.g.* [14].

We are interested in games with two participants: Opponent (O, the *environment*) and Player (P, the *program*). They play on directed graphs called *arenas*, which are semantic versions of *types*. Formally, an **arena** is a structure $A = (M_A, \lambda_A, I_A, \vdash_A)$ where:

- M_A is a set of **moves**,
- $\lambda_A : M_A \rightarrow \{O, P\}$ is a polarity function indicating whether a move is an Opponent or Player move (*O*-move or *P*-move).
- $I_A \subseteq \lambda_A^{-1}(\{O\})$ is a set of **initial moves**.
- $\vdash_A \subseteq M_A \times M_A$ is a relation called **enabling**, such that if $m \vdash_A n$, $\lambda_A(m) \neq \lambda_A(n)$.

In our complexity analysis we will use notions of *depth*. A move $m \in M_A$ has **depth** $d \in \mathbb{N}$ if there is an enabling sequence:

$$m_0 \vdash_A m_1 \vdash \dots \vdash m_{n-1} = m$$

where $m_0 \in I_A$, and no shortest enabling sequence exists; for instance any initial move has depth 0. Likewise, an arena A has depth d if d is the largest depth of any move in M_A .

We now define plays as **justified sequences** over A : these are sequences s of moves of A , each non-initial move m in s being equipped with a pointer to an earlier move n in s , satisfying $n \vdash_A m$. In other words, a justified sequence s over A is such that each reversed pointer chain $s_{i_0} \leftarrow s_{i_1} \leftarrow \dots \leftarrow s_{i_n}$ is a path on A (viewed as a graph). The role of pointers is to allow *reopenings* or *backtracking* in plays. When writing justified sequences, we will often omit the justification information if this does not cause any ambiguity. The symbol \sqsubseteq will denote the prefix ordering on justified sequences, and $s_1 \sqsubseteq^P s_2$ (resp. $s_1 \sqsubseteq^O s_2$) will mean that s_1 is a *P*-ending (resp. *O*-ending) prefix of s_2 . If s is a justified sequence on A , $|s|$ will denote its length. If s is a justified sequence over A and $\Sigma \subseteq M_A$, then the **restriction** $s \upharpoonright \Sigma$ comprises the moves of s in Σ . Pointers in $s \upharpoonright \Sigma$ are those obtained by following a path of pointers in s :

$$s_{j_1} \rightarrow s_{j_2} \rightarrow \dots \rightarrow s_{j_n}$$

where $s_{j_1}, s_{j_n} \in \Sigma$ but for $2 \leq k \leq n-1$ $s_{j_k} \notin \Sigma$. If $s = s_0 \dots s_n$ is a justified sequence and $i \leq n$, write $s_{\leq i}$ for its subsequence $s_0 \dots s_i$. The **legal plays** over A are the justified sequences s on A satisfying the **alternation** condition, *i.e.* that if $tmn \sqsubseteq s$, then $\lambda_A(m) \neq \lambda_A(n)$. The set of legal plays on A is denoted by \mathcal{L}_A .

Given a justified sequence s on A , it has two subsequences of particular interest: the P-view and O-view. The view for P (resp. O) may be understood as the subsequence of the play where P (resp. O) only sees his own duplications. Practically, the P-view $\lceil s \rceil$ of s is computed recursively by forgetting everything under Opponent's pointers, as follows:

- $\lceil sm \rceil = \lceil s \rceil m$ if $\lambda_A(m) = P$;
- $\lceil sm \rceil = m$ if $m \in I_A$ and m has no justification pointer;
- $\lceil s_1 m s_2 n \rceil = \lceil s_1 \rceil mn$ if $\lambda_A(n) = O$ and n points to m .

The O-view $\lfloor s \rfloor$ of s is defined dually, without the special treatment of initial moves.

In this subsection, we will present several classes of strategies on arena games that are of interest to us in the present paper. A **strategy** σ on A is a set of even-length legal plays

on A , closed under even-length prefix. A strategy from A to B is a strategy $\sigma : A \Rightarrow B$, where $A \Rightarrow B$ is the usual arrow arena defined by

$$\begin{aligned} M_{A \Rightarrow B} &= M_A + M_B \\ \lambda_{A \Rightarrow B} &= [\overline{\lambda_A}, \lambda_B] \\ I_{A \Rightarrow B} &= I_B \\ \vdash_{A \Rightarrow B} &= \vdash_A + \vdash_B + I_B \times I_A \end{aligned}$$

where $\overline{\lambda_A}$ means λ_A with polarity O/P reversed.

3.1.1. Composition. We define composition of strategies by the usual parallel interaction plus hiding mechanism. If A , B and C are arenas, we define the set of **interactions** $I(A, B, C)$ as the set of justified sequences u over A , B and C such that $u \upharpoonright A, B \in \mathcal{L}_{A \Rightarrow B}$, $u \upharpoonright B, C \in \mathcal{L}_{B \Rightarrow C}$ and $u \upharpoonright A, C \in \mathcal{L}_{A \Rightarrow C}$. Then, if $\sigma : A \Rightarrow B$ and $\tau : B \Rightarrow C$, their parallel interaction is $\sigma \parallel \tau = \{u \in I(A, B, C) \mid u \upharpoonright A, B \in \sigma \wedge u \upharpoonright B, C \in \tau\}$. Their composition is $\sigma; \tau = \{u \upharpoonright A, C \mid u \in \sigma \parallel \tau\}$, is associative and admits copycat strategies as identities.

3.1.2. P -visible strategies. A strategy σ is **P -visible** if each of its moves points to the current P -view. Formally, for all $sab \in \sigma$, b points inside $\lceil sa \rceil$. P -visible strategies are stable under composition, and correspond to functional programs with ground type references [1].

3.1.3. Innocent strategies. The class of *innocent* strategies is central in game semantics, because of their correspondence with purely functional programs (or λ -terms) and of their useful definability properties. A strategy σ is **innocent** if

$$sab \in \sigma \wedge t \in \sigma \wedge ta \in \mathcal{L}_A \wedge \lceil sa \rceil = \lceil ta \rceil \implies tab \in \sigma$$

Intuitively, an innocent strategy only takes its P -view into account to determine its next move. Indeed, any innocent strategy is characterized by a set of P -views. This observation is very important since P -views can be seen as abstract representations of branches of η -expanded Böhm trees (*a.k.a.* Nakajima trees [18]): this is the key to the definability process on innocent strategies. Arenas and innocent strategies form a cartesian closed category and are therefore a model of Λ ; let us add for completeness that o is interpreted as the singleton arena and $*_A$ is interpreted as the singleton strategy on $\llbracket A \rrbracket$ containing only the empty play.

3.1.4. Bounded strategies. A strategy σ is **bounded** if it is P -visible and if the length of its P -views is bounded: formally, there exists $N \in \mathbb{N}$ such that for all $s \in \sigma$, $|\lceil s \rceil| \leq N$. Bounded strategies only have finite interactions [8]; this result corresponds loosely to the normalisation result on simply-typed λ -calculus. Syntactically, bounded strategies include the interpretation of all terms of a higher-order programming language with ground type references, arbitrary non-determinism and control operators, but without recursion. This remark is important since it implies that our results will hold for any program written with these constructs, as long as they do not use recursion or a fixed point operator.

Our complexity results, in this game-theoretic part of this paper, will be expressed as a function of the **size** of the involved bounded strategies, given by:

$$|\sigma| = \frac{\max_{s \in \sigma} |\lceil s \rceil|}{2}$$

When σ is an innocent strategy coming from a Böhm tree, then $|\sigma|$ is proportional to the *height* of this Böhm tree, as defined in the syntactic part of this paper.

3.2. Visible pointer structures.

3.2.1. *Introduction of visible pointer structures.* We note here that the notion of P -view (and of its size) only takes into account the structure of pointers within the plays, and completely ignores the actual labels of the moves. In fact the underlying *pointer structure* of a play will be all we need to study the asymptotic complexity of execution.

Definition 1. The **pure arena** I_∞ is defined by:

$$\begin{aligned} M_{I_\infty} &= \mathbb{N} \\ \lambda_{I_\infty} &= \begin{cases} 2n \mapsto O \\ 2n+1 \mapsto P \end{cases} \\ \vdash_{I_\infty} &= \{(n, n+1) \mid n \in \mathbb{N}\} \\ I_{I_\infty} &= \{0\} \end{aligned}$$

A **pointer structure** is a legal play $s \in \mathcal{L}_{I_\infty}$ with at most one initial move. Note that for any arena A , a legal play $s \in \mathcal{L}_A$ with only one initial move can always be mapped to its pointer structure $\bar{s} \in \mathcal{L}_{I_\infty}$ by sending each move s_i to its *depth*, *i.e.* the number of pointers to be taken in s before reaching the initial move.

The **depth** of a pointer structure s is the largest depth of $s_i \in \mathbb{N}$, for $0 \leq i \leq |s| - 1$.

Pointer structures retain some information about the control flow of the execution, but on the other hand forget most of the typing information. Our results will rely on the crucial assumption that the strategies involved act in a *visible* way. This is necessary, because non-visible strategies are able to express programs with general (higher-order) references within which a fixpoint operator is definable, so termination is lost.

Definition 2. A **visible pointer structure** s is a pointer structure $s \in \mathcal{L}_{I_\infty}$ such that:

- It is P -visible: for any $s'a \sqsubseteq^P s$, a points within $\ulcorner s \urcorner$.
- It is O -visible: for any $s'a \sqsubseteq^O s$, a points within $\llcorner s' \lrcorner$.

We write \mathcal{V}_d for the set of all visible pointer structures of depth lower than d .

We are interested in the maximal length of a visible pointer structure resulting from the interaction between two bounded strategies. Since the collapse of plays to visible pointer structures forgets the identity of moves, all that remains from bounded strategies in this framework is their *size*. Therefore, for $n, p \in \mathbb{N}$, we define the set $n \star_d p$ of all visible pointer structures possibly resulting from an interaction of strategies of respective sizes n and p , in an ambient arena of depth d . Formally:

$$n \star_d p = \{s \in \mathcal{V}_d \mid \forall s' \sqsubseteq s, |\ulcorner s \urcorner| \leq 2n \wedge |\llcorner s' \lrcorner| \leq 2p + 1\}$$

In [8], we already examined the termination problem for visible pointer structures. We proved that any interaction between bounded strategies is necessarily finite. Therefore since $n \star_d p$, regarded as a tree, is finitely branching, it follows that it is finite. So there is an upper bound $N_d(n, p)$ to the length of any visible pointer structure in $n \star_d p$.

3.2.2. *The visible pointer structure of an interaction.* We take here the time to detail more formally our claim that an estimation of the length of visible pointer structures in $n \star_d p$ is informative of the complexity of interaction between bounded strategies.

Let $\sigma : A \Rightarrow B$ and $\tau : B \Rightarrow C$ be bounded strategies, of respective size n and p . We are interested in the possible length of interactions in $\sigma \parallel \tau$. Of course arbitrary such interactions are not bounded in size, since σ and τ both interact with an external opponent in A and C , whose behavior is not restricted by any size condition. Therefore we restrict our interest to **passive** such interactions, *i.e.* interactions $u \in \sigma \parallel \tau$ such that the unique Opponent move in $u \upharpoonright A, C$ is a unique initial question in C . When σ and τ are both innocent and correspond to λ -terms, this corresponds by the results of [9] to a linear head reduction sequence from a game situation. In particular, the *passivity* condition ensures that the interaction stops if a free variable ever arrives in head position.

Proposition 1. Let $\sigma : A \Rightarrow B$ be a bounded strategy of size p , let $\tau : B \Rightarrow C$ be a bounded strategy of size n , and suppose that B has depth $d - 1$, for $d \geq 2$. Then for all passive $u \in \sigma \parallel \tau$:

$$|u| \leq N_d(n, p) + 1$$

Proof. Since u is passive, it consists in a play $\circ_C u'$ with \circ_C initial in C and $u' \in \mathcal{L}_B$, possibly followed by a trailing move \bullet in A or C . Take any strict prefix $\circ_C u'' \sqsubseteq u$. Then u'' is either in σ or an immediate prefix of a play in σ , so $|\ulcorner u'' \urcorner| \leq 2p$. Accordingly, $|\lfloor \circ_C u'' \rfloor| \leq 2p + 1$. Likewise $\circ_C u''$ is either in τ or an immediate prefix of a play in τ , so $|\ulcorner \circ_C u'' \urcorner| \leq 2n$.

Writing \bar{u} for the **pointer structure** of a play u (*i.e.* the play of I_∞ obtained by sending any move to its depth). Then we have $\overline{\circ_C u'} \in n \star_d p$. Indeed the size of P -views and O -views in a play only depends on pointers, so it is unchanged by forgetting arena labels. Moreover $\circ_C u' \in \mathcal{L}_{B \Rightarrow \{\circ_C\}}$ (where $\{\circ_C\}$ is the singleton arena), which has depth d .

From the above, we deduce that $|\circ_C u'| \leq N_d(n, p)$, so $|u| \leq N_d(n, p) + 1$. \square

So the study of visible pointer structure is sufficient to study the length of interactions in terms of the sizes of the strategies involved.

3.2.3. *Interaction skeletons and simulation of visible pointer structures.* We now introduce *interaction skeletons* (or just *skeletons* for short), the main tool used in this paper in order to study the complexity of execution.

As we mentioned repeatedly, game-theoretic interaction corresponds to linear head reduction — which is itself efficiently implemented by machines with environment such as the Krivine Abstract Machine (KAM). In such machines, game situations produce by reduction situations where the terms interacting are no longer plain closed terms but rather terms-in-environments, also known as *closures*. Following this phenomenon, whereas the measure of the first move s_0 of a visible pointer structure is given by the sizes of the strategies involved (so by a pair of natural numbers), the measure of a later move s_i will be given by a finite tree of natural numbers reminiscent of the structure of *closures*.

We will call a **pointed visible pointer structure** a pair (s, i) where s is a visible pointer structure and $i \leq |s| - 1$ is an arbitrary “starting” move. We adapt the notions of size and depth for them, and introduce a notion of *context*.

Definition 3. Let (s, i) be a pointed visible pointer structure. The **residual size** of s at i , written $\text{rsize}(s, i)$, is defined as follows:

- If s_i is an Opponent move, it is $\max_{s_i \in \ulcorner s_{\leq j} \urcorner} |\ulcorner s_{\leq j} \urcorner| - |\ulcorner s_{\leq i} \urcorner| + 1$

- If s_i is a Player move, it is $\max_{s_i \in \ulcorner s_{\leq j} \urcorner} |\ulcorner s_{\leq j} \urcorner| - |\ulcorner s_{\leq i} \urcorner| + 1$

where $s_i \in \ulcorner s_{\leq j} \urcorner$ means that the computation of $\ulcorner s_{\leq j} \urcorner$ reaches¹ s_i . Dually, we have the notion of **residual co-size** of s at i , written $\text{rcosize}(s, i)$, defined as follows:

- If s_i is an Opponent move, it is $\max_{s_i \in \ulcorner s_{\leq j} \urcorner} |\ulcorner s_{\leq j} \urcorner| - |\ulcorner s_{\leq i} \urcorner| + 1$
- Otherwise, $\max_{s_i \in \ulcorner s_{\leq j} \urcorner} |\ulcorner s_{\leq j} \urcorner| - |\ulcorner s_{\leq i} \urcorner| + 1$

The *residual depth* of s at i is the maximal length of a pointer chain in s starting from s_i .

Definition 4. Let s be a visible pointer structure. We define the **context** of (s, i) as:

- If s_i is an O-move, the set $\{s_{n_1}, \dots, s_{n_p}\}$ of O-moves appearing in $\ulcorner s_{< i} \urcorner$,
- If s_i is a P-move, the set $\{s_{n_1}, \dots, s_{n_p}\}$ of P-moves appearing in $\ulcorner s_{< i} \urcorner$.

In other words it is the set of moves to which s_{i+1} can point whilst abiding to the visibility condition, except s_i . We also need the dual notion of co-context, which contains the moves the other player can point to. The **co-context** of (s, i) is:

- If s_i is an O-move, the set $\{s_{n_1}, \dots, s_{n_p}\}$ of P-moves appearing in $\ulcorner s_{< i} \urcorner$,
- If s_i is a P-move, the set $\{s_{n_1}, \dots, s_{n_p}\}$ of O-moves appearing in $\ulcorner s_{< i} \urcorner$.

Definition 5. A **skeleton** is a finite tree, whose nodes and edges are both labeled by natural numbers. If a_1, \dots, a_p are skeletons and d_1, \dots, d_p are natural numbers, we write:

$$n[\{d_1\}a_1, \dots, \{d_p\}a_p] = \begin{array}{c} n \\ \swarrow \quad \searrow \\ d_1 \quad \quad \quad d_p \\ \swarrow \quad \quad \quad \searrow \\ a_1 \quad \quad \quad \dots \quad \quad \quad a_p \end{array}$$

We now define what it means for (s, i) to respect a skeleton a .

Definition 6 (Trace, co-trace, interaction). The two notions Tr and $coTr$ are defined by mutual recursion, as follows: let $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$ be a skeleton. We say that (s, i) is a **trace** (resp. a **co-trace**) of a , denoted $(s, i) \in Tr(a)$ (resp. $(s, i) \in coTr(a)$) if the following conditions are satisfied:

- $\text{rsize}(s, i) \leq 2n$ (resp. $\text{rcosize}(s, i) \leq 2n + 1$),
- If $\{s_{n_1}, \dots, s_{n_p}\}$ is the context of (s, i) (resp. co-context), then for each $k \in \{1, \dots, p\}$ we have $(s, n_k) \in coTr(a_k)$.
- If $\{s_{n_1}, \dots, s_{n_p}\}$ is the context of (s, i) (resp. co-context), then for each $k \in \{1, \dots, p\}$ the residual depth of s at n_k is less than d_k .

Then, we define an **interaction** of two skeletons a and b at depth d as a pair $(s, i) \in Tr(a) \cap coTr(b)$ where the residual depth of s at i is less than d , which we write $(s, i) \in a \star_d b$.

Notice that we use the same notation \star both for natural numbers and skeletons. This should not generate any confusion, since the definitions above coincide with the previous ones in the special case of “atomic” skeletons: if n and p are natural numbers, then obviously $s \in n \star_d p$ (according to the former definition) if and only if $(s, 0) \in n[\] \star_d p[\]$ (according to the latter). In fact, we will sometimes in the sequel write n for the atomic skeleton with n at the root and without children, *i.e.* $n[\]$.

¹So starting from s_j and following Opponent’s pointers eventually reaches s_i .

3.2.4. *Simulation of visible pointer structures.* We introduce now our main tool, a reduction on skeletons which simulates visible pointer structures: if $n[\{d_1\}a_1, \dots, \{d_p\}a_p]$ and b are skeletons ($n > 0$), we define the non-deterministic reduction relation \rightsquigarrow on triples (a, d, b) , where d is a depth (a natural number) and a and b are skeletons, by the following two cases:

$$\begin{aligned} (n[\{d_1\}a_1, \dots, \{d_p\}a_p], d, b) &\rightsquigarrow (a_i, d_i - 1, (n - 1)[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b]) \\ (n[\{d_1\}a_1, \dots, \{d_p\}a_p], d, b) &\rightsquigarrow (b, d - 1, (n - 1)[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b]) \end{aligned}$$

where $i \in \{1, \dots, p\}$, $d_i > 0$ in the first case and $d > 0$ in the second case.

In order to prove our simulation result, we will make use of the following lemma.

Lemma 1. Let s be a pointed visible pointer structure and $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$ a skeleton such that $(s, i) \in \text{coTr}(a)$. Then if $s_j \rightarrow s_i$, $(s, j) \in \text{Tr}(a)$.

Proof. Let us suppose without loss of generality that s_i is an Opponent move; the other case can be obtained just by switching Player/Opponent and P -views/ O -views everywhere. Then s_j being a Player move, we have to check first that $\text{rsize}(s, j) \leq 2n$, *i.e.*

$$\max_{s_j \in \ulcorner s \leq k \urcorner} |\ulcorner s \leq k \urcorner| - |\ulcorner s \leq j \urcorner| + 1 \leq 2n$$

We use that $\text{rcoSize}(s, i) \leq 2n + 1$, *i.e.*

$$\max_{s_i \in \ulcorner s \leq k \urcorner} |\ulcorner s \leq k \urcorner| - |\ulcorner s \leq i \urcorner| + 1 \leq 2n + 1$$

But $s_j \rightarrow s_i$, hence $|\ulcorner s \leq j \urcorner| = |\ulcorner s \leq i \urcorner| + 1$ and the inequality is obvious. We need now to examine the context of (s, j) . Since s_j is a Player move, it is defined as the set $\{s_{n_1}, \dots, s_{n_p}\}$ of Player moves appearing in $\ulcorner s \leq j \urcorner$, which is also the set of Player moves appearing in $\ulcorner s \leq i \urcorner$ and therefore the co-context of (s, i) . But $(s, i) \in \text{coTr}(a)$, hence for all $k \in \{1, \dots, p\}$ we have $(s, n_k) \in \text{coTr}(a_k)$ which is exactly what we needed. \square

Proposition 2 (Simulation). Let $(s, i) \in a \star_d b$, then if s_{i+1} is defined, there exists $(a, d, b) \rightsquigarrow (a', d', b')$ such that $(s, i + 1) \in a' \star_{d'} b'$.

Proof. Suppose $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$. Let $\{s_{n_1}, \dots, s_{n_p}\}$ be the context of (s, i) . By visibility, s_{i+1} must either point to s_i or to an element of the context. Two cases:

- If $s_{i+1} \rightarrow s_i$, then we claim that $(s, i + 1) \in b \star_{d-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b]$, *i.e.* $(s, i + 1) \in \text{Tr}(b)$, $(s, i + 1) \in \text{coTr}((n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b])$ and the depth of s relative to $i + 1$ is at most $d - 1$. For the first part, we use that $(s, i) \in a \star_d b$: in particular, $(s, i) \in \text{coTr}(b)$ and since $s_{i+1} \rightarrow s_i$ this implies by Lemma 1 that $(s, i + 1) \in \text{Tr}(b)$. For the second part, we must first check that $\text{rcoSize}(s, i + 1) \leq 2(n - 1) + 1$. Let us suppose without loss of generality that s_i is an Opponent move, all the reasoning below can be adapted by switching Player/Opponent and P -views/ O -views everywhere. We want to prove:

$$\text{rcoSize}(s, i + 1) = \max_{s_{i+1} \in \ulcorner s \leq j \urcorner} |\ulcorner s \leq j \urcorner| - |\ulcorner s \leq i+1 \urcorner| + 1 \leq 2(n - 1) + 1$$

But since $(s, i) \in \text{Tr}(a)$, we already know:

$$\text{rsize}(s, i) = \max_{s_i \in \ulcorner s \leq j \urcorner} |\ulcorner s \leq j \urcorner| - |\ulcorner s \leq i \urcorner| + 1 \leq 2n$$

Thus we only need to remark that $|\ulcorner s \leq i+1 \urcorner| = |\ulcorner s \leq i \urcorner| + 1$ since s_{i+1} is a Player move. Now, we must examine the co-context of $(s, i + 1)$, but by definition of P -view it is $\{s_{n_1}, \dots, s_{n_p}, s_i\}$ where $\{s_{n_1}, \dots, s_{n_p}\}$ is the context of (s, i) . Since $(s, i) \in$

$Tr(n[a_1, \dots, a_p])$ we have as required $(s, n_k) \in coTr(a_k)$ for each $k \in \{1, \dots, p\}$ and $(s, i) \in coTr(b)$ because $(s, i) \in a \star_d b$. For the third part, we have to prove that the depth of s relative to $i+1$ is at most $d-1$, but it is obvious since the depth relative to i is at most d and $s_{i+1} \rightarrow s_i$.

- Otherwise, we have $s_{i+1} \rightarrow s_{n_j}$ for $j \in \{1, \dots, p\}$. Then, we claim that $(s, i+1) \in a_j \star_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b]$. We do have $(s, i+1) \in Tr(a_j)$ because $(s, i) \in Tr(n[\{d_1\}a_1, \dots, \{d_p\}a_p])$, thus $(s, n_j) \in coTr(a_j)$ and $(s, i+1) \in Tr(a_j)$ by Lemma 1. It remains to show that $(s, i+1) \in coTr((n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b])$ and that the depth of s relative to $i+1$ is at most d_1-1 , but the proofs are exactly the same as in the previous case. \square

Before going on to the study of skeletons, let us give a last simplification. If $a = n[\{d_1\}t_1, \dots, \{d_p\}t_p]$ and b are skeletons, then $a \cdot_d b$ will denote the skeleton obtained by appending b as a new son of the root of a with label d , i.e. $n[\{d_1\}t_1, \dots, \{d_p\}t_p, \{d\}b]$. Consider the following non-deterministic rewriting rule on skeletons:

$$n[\{d_1\}a_1, \dots, \{d_p\}a_p] \rightsquigarrow a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$$

Both rewriting rules on triples (a, d, b) are actually instances of this reduction, by the isomorphism $(a, d, b) \mapsto a \cdot_d b$. We leave the obvious verification to the reader. Taking this into account, all that remains to study is this reduction on skeletons illustrated in Figures 1 and 2, and analyzed in the next section.

If $\mathcal{N}(a)$ denotes the length of the longest reduction sequence starting from a skeleton a , we have the following property.

Proposition 3. Let $n, p \geq 0$, $d \geq 2$, then $N_d(n, p) \leq \mathcal{N}(n[\{d\}p]) + 1$.

Proof. Obvious from Proposition 2, adding 1 for the initial move which is not accounted for by the reduction on skeletons. \square

We postpone the analysis of the reduction of skeletons to the next section. However with the results proved there, we get the following result:

Theorem 1. For fixed $d \geq 3$, we have $N_d(n, p) = 2^{\Theta(n \log(p))}$.

Proof. Upper bound. Follows from Proposition 3 and Theorem 3.

Lower bound. To construct the example providing the lower bound, it is convenient to consider the extension Λ_\times of Λ with finite product types $\prod_{1 \leq i \leq n} A_i$. Tupling of $\Gamma \vdash M_i : A_i$ for $1 \leq i \leq n$ is written $\langle M_1, \dots, M_n \rangle$. For simplicity we use $\lambda \langle x_1, \dots, x_n \rangle^{\prod_{1 \leq i \leq n} A_i}. M$ as syntactic sugar for $\lambda x^{\prod_{1 \leq i \leq n} A_i}. M[\pi_i x / x_i]$, where $\pi_i : \prod_{1 \leq i \leq n} A_i \rightarrow A_i$ is the corresponding projection. Products are interpreted in the games model following standard lines [15]. In Λ_\times (and Λ), we define higher types for Church integers by setting $A_{-2} = o$ and $A_{n+1} = A_n \rightarrow A_n$. For $n, p \in \mathbb{N}$, we write \underline{n}_p for the Church integer for n of type A_p . For A a type, if $M : A \rightarrow A$ and $N : A$, we write $M^n(N)$ for the n -th iteration of M applied to N . Then for $n, p, d \in \mathbb{N}$, we define:

$$(\lambda \langle x, y, z_1, \dots, z_d \rangle. x^n(y) z_1 \dots z_d) \langle \underline{p}_{d+1}, \underline{2}_d, \underline{2}_{d-1}, \dots, \underline{2}_0 \rangle$$

By elementary calculations on λ -terms, we know that this is β -equivalent to $\underline{2}_{d+1}^n$. So, taking a maximal (necessarily passive) interaction:

$$u \in \llbracket \langle \underline{p}_{d+1}, \underline{2}_d, \underline{2}_{d-1}, \dots, \underline{2}_0, \text{id}_o \rangle \rrbracket \parallel \llbracket \lambda \langle x, y, z_1, \dots, z_d, i \rangle. x^n(y) z_1 \dots z_d i \rrbracket$$

we know that u must have length at least 2^{p^n} . Inspecting these strategies, we see that the left hand side one has size $p + d + 2$ and the right hand side one has size $n + d + 3$, and that they interact in an arena $\llbracket \prod_{i=0}^{d+2} A_{d+1-i} \rrbracket$ of depth $d + 3$. It follows that the underlying pointer structure of u (of length at least 2^{p^n}) is in $(n + d + 3) \star_{d+4} (p + d + 2)$. Therefore,

$$N_{d+4}(n + d + 3, p + d + 2) \geq 2^{p^n}_{d+1}$$

So for $d \geq 4, n \geq d + 3, p \geq d + 2$ we have $N_d(n, p) \geq 2^{(n-d-3)\log(p-d-2)}_{d-2}$, providing the lower bound. Although this example only proves the other bound for $d \geq 4$ it also holds for $d = 3$: this is proved using the same reasoning on the maximal interaction between strategies $\llbracket p_0 \rrbracket$ and $\llbracket \lambda x. x^n(\text{id}_o) \rrbracket$, having length at least p^n . \square

The strength of this result is that being a theorem about interactions of strategies in game semantics, its scope includes any programming language whose terms can be interpreted as bounded strategies. Its weakness however, is that it only applies syntactically to game situations. In order to increase the generality of our study and give exact bounds to the linear head reduction of arbitrary simply-typed λ -terms, we will in Section 5 detail a direct connection between linear head reduction and reduction of skeletons. Before that, as announced we focus Section 4 on the analysis of skeletons.

4. SKELETONS AND THEIR COMPLEXITY ANALYSIS

In the previous section we proved a simulation result of *plays* in the sense of Hyland-Ong games into a reduction on simple combinatorial objects, *interaction skeletons*. In the present section we investigate the properties of this reduction independently of game semantics or syntax, proving among other things Theorem 3 used in the previous section.

As announced in the introduction, the rest of this paper – starting from here – is essentially self-contained. We start this section by recalling the definition of interaction skeletons and investigating their basic properties. Then, we will prove our main result about the length of their reduction.

4.1. Skeletons and their basic properties.

4.1.1. *Skeletons and their dynamics.* **Interaction skeletons**, or *skeletons* for short, are finite trees of natural numbers, whose nodes and edges are labeled by natural numbers. To denote these finite trees, we use the notation illustrated below:

$$n[\{d_1\}a_1, \dots, \{d_p\}a_p] = \begin{array}{c} n \\ \begin{array}{ccc} / & & \backslash \\ d_1 & & d_p \\ \backslash & & / \\ a_1 & \dots & a_p \end{array} \end{array}$$

Each natural number n can be seen as an atomic skeleton $n\llbracket$ without subtrees, still denoted by n . That should never cause any confusion. Given a skeleton a , we define:

- Its **order** $\text{ord}(a)$, the maximal edge label in a ,
- Its **maximum** $\text{max}(a)$, the maximal node label in a ,
- Its **depth** $\text{depth}(a)$, the maximal depth of a node in a , the root having depth 1.




Figure 1: Rewriting rule on skeletons




Figure 2: Example reduction sequence on interaction skeletons

We will also use the notation $a \cdot_d b$ for the skeleton a with a new child b added to the root of a , with edge label d . Formally if $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$:

$$a \cdot_d b = n[\{d_1\}a_1, \dots, \{d_p\}a_p, \{d\}b]$$

With that in place, we define the reduction on skeletons by:

$$n[\{d_1\}a_1, \dots, \{d_p\}a_p] \rightsquigarrow a_i \cdot_{d_{i-1}} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$$

which is allowed whenever $n, d_i \geq 1$ so that no index ever becomes negative. The reduction is illustrated in Figure 1, and performed on an example in Figure 2 (where the node selected for the next reduction is highlighted). It is important to insist that this reduction is only ever performed in head position: n needs to be the actual root of the tree for the reduction to be allowed. We do not know which properties of this reduction are preserved in the generalized system where reduction can occur everywhere.

Let us write $\mathcal{N}(a)$ for the **norm** of a skeleton a , *i.e.* the length of its longest reduction sequence. We will show later that it is always finite; in the meantime for definiteness we define it as a member of $\mathbb{N} \cup \{+\infty\}$.

4.1.2. Embedding lemma. The norm of a skeleton is unchanged by permutation of subtrees, or merging of identical subtrees, and is only increased by an increase of labels. If $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$ and $a' = n'[\{d'_1\}a'_1, \dots, \{d'_{p'}\}a'_{p'}]$, we say that a **embeds** in a' , written $a \hookrightarrow a'$, if $n \leq n'$ and for any $i \in \{1, \dots, p\}$ there exists $j \in \{1, \dots, p'\}$ such that $d_i \leq d'_j$ and $a_i \hookrightarrow a'_j$. Then we have:

Lemma 2 (Embedding lemma). For any skeletons a and a' with $a \hookrightarrow a'$, $\mathcal{N}(a) \leq \mathcal{N}(a')$.

Proof. Suppose $a \hookrightarrow a'$, and $a \rightsquigarrow b$. Write $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$, and suppose $b = a_i \cdot_{d_{i-1}} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$. Likewise, write $a' = n'[\{d'_1\}a'_1, \dots, \{d'_{p'}\}a'_{p'}]$. Since $a \hookrightarrow a'$ we have $n' \geq n \geq 1$ and there is $j \in \{1, \dots, p'\}$ such that $d'_j \geq d_i \geq 1$ and $a_i \hookrightarrow a'_j$. Set $b' = a'_j \cdot (n'-1)[\{d'_1\}a'_1, \dots, \{d'_{p'}\}a'_{p'}]$, it is straightforward to check that

$b \hookrightarrow b'$. Therefore, \hookrightarrow is a simulation, which implies that if $a \hookrightarrow a'$ we have $\mathcal{N}(a) \leq \mathcal{N}(a')$. \square

From this it follows that permuting subtrees in a skeleton does not affect the possible reductions in any way. Perhaps more surprisingly, it shows that two identical subtrees can be merged without any effect on the possible reductions: the number of copies of identical subtrees does not matter. Following this idea, we are going to show that any skeleton embeds into a simple thread-like one, and this only increases the length of possible reductions.

Definition 7. Let $d, o, m \geq 1$ be natural numbers. The *thread-like skeleton* $\mathcal{T}(d, o, m)$ is:

$$\begin{aligned}\mathcal{T}(1, o, m) &= m \\ \mathcal{T}(d+1, o, m) &= m[\{o\}\mathcal{T}(d, o, m)]\end{aligned}$$

From the definition, we have that $\text{depth}(\mathcal{T}(d, o, m)) = d$, $\text{max}(\mathcal{T}(d, o, m)) = m$ and $\text{ord}(\mathcal{T}(d, o, m)) = o$. We also have:

Proposition 4. If a has depth d , order o and maximum m , then $\mathcal{N}(a) \leq \mathcal{N}(\mathcal{T}(d, o, m))$.

Proof. It is obvious that $a \hookrightarrow \mathcal{T}(d, o, m)$, therefore the result follows from Lemma 2. \square

4.1.3. *Constructions on skeletons.* If $(a_i)_{1 \leq i \leq n}$ is a finite family of skeletons, then writing $a_i = n_i[\{d_{i,1}\}b_{i,1}, \dots, \{d_{i,p_i}\}b_{i,p_i}]$, we define:

$$\begin{aligned}\bigsqcup_{i=1}^n a_i &= \left(\max_{1 \leq i \leq n} n_i\right) \cdot [\{d_{i,j}\}b_{i,j} \mid 1 \leq i \leq n \ \& \ 1 \leq j \leq p_i] \\ \sum_{i=1}^n a_i &= \left(\sum_{i=1}^n n_i\right) \cdot [\{d_{i,j}\}b_{i,j} \mid 1 \leq i \leq n \ \& \ 1 \leq j \leq p_i]\end{aligned}$$

so, they either take the maximum or the sum of the roots, and simply append all the subtrees of the a_i s. In the binary case, we write as usual $+$ for the sum.

Lemma 3. We have the following embeddings:

- If $(a_i)_{1 \leq i \leq n}$, $(b_i)_{1 \leq i \leq n}$ are finite families of skeletons such that for all $1 \leq i \leq n$, we have $a_i \hookrightarrow b_i$, then

$$\bigsqcup_{i=1}^n a_i \hookrightarrow \bigsqcup_{i=1}^n b_i \qquad \sum_{i=1}^n a_i \hookrightarrow \sum_{i=1}^n b_i$$

- If $(a_i)_{1 \leq i \leq n}$ is a finite family of skeletons and b is a skeleton, then

$$\bigsqcup_{i=1}^n (a_i \cdot b) \hookrightarrow \left(\bigsqcup_{i=1}^n a_i\right) \cdot b \qquad \sum_{i=1}^n (a_i \cdot b) \hookrightarrow \left(\sum_{i=1}^n b_i\right) \cdot b$$

- If a, b, c are skeletons and $d \in \mathbb{N}$, then:

$$a + b \cdot_d c \hookrightarrow (a + b) \cdot_d c$$

Proof. Direct from the definitions. \square

4.2. Upper bounds. We calculate upper bounds to the length of possible reductions on skeletons. This is done by adapting a technique used by Schwichtenberg [19] and Beckmann [4] to bound the length of possible β -reduction sequences on simply-typed λ -terms.

The idea of the proof is to define an inductive predicate $\frac{\alpha}{\rho}$ on terms/skeletons whose proofs/inhabitants combine aspects of a syntax tree and of a reduction tree — witnesses of this predicate are called *expanded reduction trees* by Beckmann [4]. Their mixed nature will allow us to define a transformation gradually eliminating their *syntactic* (or *static*) nodes, yielding an alternative expanded reduction tree for the term/skeleton under study, whose height is more easily controlled.

Definition 8. The predicate $\frac{\alpha}{\rho}$ (where ρ, α range over natural numbers) is defined on skeletons in the following inductive way.

- **BASE.** $\frac{\alpha}{\rho} 0[\{d_1\}a_1, \dots, \{d_p\}a_p]$
- **RED.** Suppose $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$ for $n \geq 1$. Then if for all a' such that $a \rightsquigarrow a'$ we have $\frac{\alpha}{\rho} a'$ and if we also have $\frac{\alpha}{\rho} \max(n-1, 0)[\{d_1\}a_1, \dots, \{d_p\}a_p]$, then $\frac{\alpha+1}{\rho} a$.
- **CUT.** If $\frac{\alpha}{\rho} a, \frac{\beta}{\rho} b$ and $d \leq \rho$, then $\frac{\alpha+\beta}{\rho} a \cdot_d b$.

Definition 9. A **context-skeleton** $a()$ is a finite tree whose edges are labeled by natural numbers, and whose nodes are labeled either by natural numbers, or by the variable x , with the constraint that all edges leading to x must be labeled by the same number d ; d is called the **type** of x in $a()$. We denote by $a(b)$ the result of substituting all occurrences of x in $a()$ by b . We denote by $a(\emptyset)$ the skeleton obtained by deleting in a all occurrences of x , along with the edges leading to them.

Lemma 4 (Monotonicity). If $\frac{\alpha}{\rho} a$, then $\frac{\alpha'}{\rho'} a$ for all $\alpha \leq \alpha'$ and $\rho \leq \rho'$, where the witness trees have the same number of occurrences of **CUT**.

Proof. Straightforward by induction on the derivation for $\frac{\alpha}{\rho} a$. □

Lemma 5 (Permutation lemma). If a is obtained from a' by permuting some subtrees in a , then for all ρ, α , $\frac{\alpha}{\rho} a$ iff $\frac{\alpha}{\rho} a'$.

Proof. Straightforward by induction on the derivation for $\frac{\alpha}{\rho} a$. □

Lemma 6 (Null substitution lemma). If $\frac{\alpha}{\rho} a(\emptyset)$ and the type of x in a is 0, then for all b we still have $\frac{\alpha}{\rho} a(b)$. Moreover, the witness includes as many **CUT** rules as for $\frac{\alpha}{\rho} a(\emptyset)$.

Proof. We prove by induction on derivations $\frac{\alpha}{\rho} a$ that the property above holds for all context-skeleton a' such that the type of x in a' is 0 and $a'(\emptyset) = a$.

- **BASE.** The root of a is 0, hence the result is trivial.
- **RED.** Suppose a' has the form $n[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{0\}x]$, where a'_1, \dots, a'_p possibly include occurrences of x (the case where x appears as a son of the root encompasses the other) and $a_i = a'_i(\emptyset)$. The premises of **RED** are then that for $1 \leq i \leq p$ such that $d_i \geq 1$, $\frac{\alpha-1}{\rho} a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$ and $\frac{\alpha-1}{\rho} \max(n-1, 0)[\{d_1\}a_1, \dots, \{d_p\}a_p]$. The **IH** on these premises give witnesses for the two following properties:

$$\frac{\alpha-1}{\rho} (a'_i \cdot_{d_i-1} (n-1)[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{0\}x])(b) \quad (4.1)$$

$$\left| \frac{\alpha-1}{\rho} \right| (\max(n-1, 0)[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{0\}x])(b) \quad (4.2)$$

This covers all the possible reductions of $a'(b)$, thus by RED we have $\left| \frac{(\alpha-1)+1}{\rho} \right| a'(b)$ as required.

- CUT. Let us suppose $\left| \frac{\alpha+\gamma}{\rho} \right| a$ is obtained by CUT, hence a has the form $a_1 \cdot_{d_1} a_2$. Let us assume that a' has the form $(a'_1 \cdot_{d'} a'_2) \cdot_0 x$ with $a'_1(\emptyset) = a_1$ and $a'_2(\emptyset) = a_2$, since again the case where x is a child of the root of a' encompasses the other. The premises of CUT are then $\left| \frac{\alpha}{\rho} \right| a_1$ and $\left| \frac{\gamma}{\rho} \right| a_2$, and $d' \leq \rho$. We also have $(a'_1 \cdot_0 x)(\emptyset) = a_1$, therefore the IH on $\left| \frac{\alpha}{\rho} \right| a_1$ along with $\left| \frac{\beta}{\rho} \right| b$ and $0 \leq \rho + 1$ implies that $\left| \frac{\alpha}{\rho} \right| a'_1(b) \cdot_0 b$. But by IH we also have $\left| \frac{\gamma}{\rho} \right| a'_2(b)$, hence by CUT:

$$\left| \frac{\alpha+\gamma}{\rho} \right| (a'_1(b) \cdot_0 b) \cdot_{d'} a'_2(b)$$

Which was what was required for $(a'_1(b) \cdot_{d'} a'_2(b)) \cdot_0 b$, thus it suffices since trees are considered up to permutation. \square

Lemma 7 (Main substitution lemma). If $\left| \frac{\alpha}{\rho} \right| a(\emptyset)$, $\left| \frac{\beta}{\rho} \right| b$ and $d \leq \rho + 1$ (where d is the type of x in a), then $\left| \frac{\alpha(\beta+1)}{\rho} \right| a(b)$

Proof. We prove by induction on derivations $\left| \frac{\alpha}{\rho} \right| a$ that the property above holds for all context-skeleton a' such that the type of x in a' is $d \leq \rho + 1$, and such that $a = a'(\emptyset)$.

- BASE. The root of a is 0, hence the result is trivial.
- RED. Suppose a' has the form $n[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{d\}x]$, where $a_1 = a'_1(\emptyset), \dots, a_p = a'_p(\emptyset)$ (the case where x appears as a son of the root encompasses the other). The premises of RED are that for $1 \leq i \leq p$ such that $d_i \geq 1$, $\left| \frac{\alpha-1}{\rho} \right| a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$ and $\left| \frac{\alpha-1}{\rho} \right| (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$. The IH on these premises give witnesses for the two following properties:

$$\left| \frac{(\alpha-1)(\beta+1)}{\rho} \right| (a'_i \cdot_{d_i-1} (n-1)[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{d\}x])(b) \quad (4.3)$$

$$\left| \frac{(\alpha-1)(\beta+1)}{\rho} \right| ((n-1)[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{d\}x])(b) \quad (4.4)$$

By hypothesis we have $\left| \frac{\beta}{\rho} \right| b$, hence by CUT (since $d-1 \leq \rho$), we have:

$$\left| \frac{(\alpha-1)(\beta+1)+\beta}{\rho} \right| b \cdot_{d-1} (n-1)[\{d_1\}a'_1(b), \dots, \{d_p\}a'_p(b), \{d\}b] \quad (4.5)$$

Using (4.3) for all $i \in \{1, \dots, p\}$, (4.4) (adjusted to $\left| \frac{(\alpha-1)(\beta+1)+\beta}{\rho} \right|$ by Lemma 4) and (4.5) we deduce by RED that

$$\left| \frac{(\alpha-1)(\beta+1)+\beta+1}{\rho} \right| n[\{d_1\}a'_1(b), \dots, \{d_p\}a'_p(b), \{d\}b]$$

Which is what was required.

- CUT. Let us suppose $\left| \frac{\alpha+\gamma}{\rho} \right| a$ is obtained by CUT, hence a has the form $a_1 \cdot_{d'} a_2$. Let us suppose that a' has the form $(a'_1 \cdot_{d'} a'_2) \cdot_d x$ with $a_1 = a'_1(\emptyset)$ and $a_2 = a'_2(\emptyset)$, since once again the case where x is a child of the root of a' encompasses the other. The premises of CUT are then $\left| \frac{\alpha}{\rho} \right| a_1$ and $\left| \frac{\gamma}{\rho} \right| a_2$, and $d' \leq \rho$. We also have $(a'_1 \cdot_d x)(\emptyset) = a_1$,

therefore the IH on $\left|\frac{\alpha}{\rho}\right| a_1$ along with $\left|\frac{\beta}{\rho}\right| b$ and $d \leq \rho + 1$ implies that $\left|\frac{\alpha(\beta+1)}{\rho}\right| a'_1(b) \cdot_d b$. But by IH we also have $\left|\frac{\gamma(\beta+1)}{\rho}\right| a'_2(b)$, hence by CUT:

$$\left|\frac{\alpha(\beta+1)+\gamma(\beta+1)}{\rho}\right| (a'_1(b) \cdot_d b) \cdot_{d'} a'_2(b)$$

Which is what was required, up to permutation. \square

The following lemma is the core of the proof, allowing to eliminate instances of the CUT rule in the expanded head reduction tree.

Lemma 8 (Cut elimination lemma). Suppose $\left|\frac{\alpha}{\rho+1}\right| a$. Then if $\alpha = 0$, $\left|\frac{0}{\rho}\right| a$. Otherwise, $\left|\frac{2^{\alpha-1}}{\rho}\right| a$.

Proof. By induction on the tree witness for $\left|\frac{\alpha}{\rho+1}\right| a$.

- BASE. Trivial.
- RED. Suppose $a = n[\{d_1\}a_1, \dots, \{d_p\}a_p]$. The premises of RED are $\left|\frac{\alpha-1}{\rho+1}\right| a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$ for all $i \in \{1, \dots, p\}$ and $\left|\frac{\alpha-1}{\rho+1}\right| (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$. If $\alpha \geq 2$, then it follows by IH that $\left|\frac{2^{\alpha-2}}{\rho}\right| a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$ and $\left|\frac{2^{\alpha-2}}{\rho}\right| (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$, which implies by RED and Lemma 4 that $\left|\frac{2^{\alpha-1}}{\rho}\right| a$. If $\alpha = 1$, then the premises of RED are $\left|\frac{0}{\rho+1}\right| a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$ for all $i \in \{1, \dots, p\}$ and $\left|\frac{0}{\rho+1}\right| (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p]$. By induction hypothesis this is still true with ρ instead of $\rho + 1$, thus by RED we have $\left|\frac{1}{\rho}\right| [\{d_1\}a_1, \dots, \{d_p\}a_p]$ which is what we needed to prove.
- CUT. Suppose $a = a_1 \cdot_d a_2$, the premises of CUT are $\left|\frac{\alpha}{\rho+1}\right| a_1$, $\left|\frac{\beta}{\rho+1}\right| a_2$ and $d \leq \rho + 1$. If $\alpha, \beta \geq 1$ then by IH it follows that $\left|\frac{2^{\alpha-1}}{\rho}\right| a_1$ and $\left|\frac{2^{\beta-1}}{\rho}\right| a_2$, in particular if we define a context-skeleton $a'_1 = a_1 \cdot_d x$ we have $\left|\frac{2^{\alpha-1}}{\rho}\right| a'_1(\emptyset)$, hence by the substitution lemma (since $d \leq \rho + 1$) we have $\left|\frac{2^{\alpha-1}(2^{\beta-1}+1)}{\rho}\right| a'_1(a_2) = a_1 \cdot_d a_2 = a$, thus $\left|\frac{2^{\alpha+\beta-1}}{\rho}\right| a$ thanks to Lemma 4 (since it is always true that $2^{\alpha+\beta-1} \geq 2^{\alpha-1}(2^{\beta-1}+1)$). If $\alpha = 0$ then by IH we have $\left|\frac{0}{\rho}\right| a_1$ and $\left|\frac{\beta'}{\rho}\right| a_2$. We use then the substitution lemma (since $d \leq \rho + 1$) to get $\left|\frac{0}{\rho}\right| (a_1 \cdot_d a_2)$, which is stronger than what was required whatever was the value of β . The last remaining case is when $\alpha = 1$ and $\beta = 0$, then by IH $\left|\frac{1}{\rho}\right| a_1$ and $\left|\frac{0}{\rho}\right| a_2$, thus by the substitution lemma we have as required $\left|\frac{1}{\rho}\right| (a_1 \cdot_d a_2)$. \square

The lemma above allows us to transform any expanded reduction tree into a purely dynamic one (using only rules BASE and RED²). Now, we show how an expanded reduction tree can be automatically inferred for any skeleton.

Lemma 9 (Recomposition lemma). Let a be a skeleton such that $\text{ord}(a) \geq 1$. Let $\text{nodes}(a)$ denote the multiset of labels of nodes in a . Then, $\left|\frac{\prod_{n \in \text{nodes}(a)} (n+1)}{\text{ord}(a)-1}\right| a$.

Proof. First, let us show that the following rule BASE' is admissible, for any α and ρ .

$$\left|\frac{\alpha+n}{\rho}\right| n$$

²In fact at this point the expanded reduction tree can still contain cuts of order 0, eliminated later.

If $n = 0$ this is exactly BASE. Otherwise we apply RED. There is no possible reduction, so the only thing we have to prove is $\left\lfloor \frac{\alpha+n-1}{\rho} (n-1) \right\rfloor$, which is by IH.

The lemma follows by applying Lemma 7 once for each node. \square

Now, we show how to deduce from a purely dynamic expanded reduction tree a bound to the length of possible reductions.

Lemma 10 (Bound lemma). Let a be a skeleton, then if $\left\lfloor \frac{\alpha}{0} a \right\rfloor$, $\mathcal{N}(a) \leq \alpha$.

Proof. First of all we prove that if there is a witness for $\left\lfloor \frac{\alpha}{0} a \right\rfloor$, then it can be supposed CUT-free. We reason by induction on $\left\lfloor \frac{\alpha}{0} a \right\rfloor$.

- BASE. The rule has no premise, so the witness tree for $\left\lfloor \frac{\alpha}{0} a \right\rfloor$ is already CUT-free.
- RED. By IH, CUT can be eliminated in the premises of $\left\lfloor \frac{\alpha+1}{0} a \right\rfloor$. Therefore by RED, there is a CUT-free witness for $\left\lfloor \frac{\alpha+1}{0} a \right\rfloor$.
- CUT. Suppose we have $\left\lfloor \frac{\alpha+\beta}{0} a \cdot_0 b \right\rfloor$ by CUT, whose premises are $\left\lfloor \frac{\alpha}{0} a \right\rfloor$ and $\left\lfloor \frac{\beta}{0} b \right\rfloor$. By IH, we can assume the witness trees for $\left\lfloor \frac{\alpha}{0} a \right\rfloor$ and $\left\lfloor \frac{\beta}{0} b \right\rfloor$ to be CUT-free. Let us form the context-skeleton $a() = a \cdot_0 x$. Then by Lemma 6 we have $\left\lfloor \frac{\alpha}{0} a(b) \right\rfloor = a \cdot_0 b$, and by Lemma 4 that $\left\lfloor \frac{\alpha+\beta}{0} a \cdot_0 b \right\rfloor$. Since the witness tree for $\left\lfloor \frac{\alpha}{0} a(\emptyset) \right\rfloor$ is CUT-free, so are the witness trees for $\left\lfloor \frac{\alpha}{0} a \cdot_0 b \right\rfloor$ and $\left\lfloor \frac{\alpha+\beta}{0} a \cdot_0 b \right\rfloor$.

Then, we prove the lemma by induction on the CUT-free witness tree for $\left\lfloor \frac{\alpha}{0} a \right\rfloor$:

- BASE. Necessarily, the root of a is 0, thus $\mathcal{N}(a) = 0$; there is nothing to prove.
- RED. The premises of $\left\lfloor \frac{\alpha}{0} a \right\rfloor$ include in particular that for all a' such that $a \rightsquigarrow a'$, we have $\left\lfloor \frac{\alpha-1}{0} a' \right\rfloor$. By IH, this means that for all such a' we have $\mathcal{N}(a') \leq \alpha - 1$, hence $\mathcal{N}(a) \leq \alpha$. \square

From all this, it is possible to give a first upper bound by using the recomposition lemma, then iterating the cut elimination lemma. However, we will first prove here a refined version of the cut elimination lemma when $\rho = 1$, which will allow to decrease by one the height of the tower of exponentials. First, we need the following adaptation of the substitution lemma:

Lemma 11 (Base substitution lemma). If $\left\lfloor \frac{\alpha}{0} a(\emptyset) \right\rfloor$, $\left\lfloor \frac{\beta}{0} b \right\rfloor$ and the type of x in a is 1, then $\left\lfloor \frac{\alpha+\beta}{0} a(b) \right\rfloor$.

Proof. We prove by induction on derivations $\left\lfloor \frac{\alpha}{0} a(\emptyset) \right\rfloor$ that the property above holds for all context-arena a' such that the type of x in a' is 1 and $a = a'(\emptyset)$.

- BASE. The root of a is 0, hence the result is trivial.
- RED. Suppose a' has the form $n[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{1\}x]$, with $a_i = a'_i(\emptyset)$ (the case where x appears as a son of the root encompasses the other). The premises of RED are then that for $1 \leq i \leq p$ such that $d_i \geq 1$, $\left\lfloor \frac{\alpha-1}{0} a_i \cdot_{d_i-1} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p] \right\rfloor$ and $\left\lfloor \frac{\alpha-1}{0} (n-1)[\{d_1\}a_1, \dots, \{d_p\}a_p] \right\rfloor$. The IH on these premises give witnesses for the two following properties:

$$\left\lfloor \frac{\alpha-1+\beta}{0} (a'_i \cdot_{d_i-1} (n-1)[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{1\}x])(b) \right\rfloor \quad (4.6)$$

$$\left\lfloor \frac{\alpha-1+\beta}{0} ((n-1)[\{d_1\}a'_1, \dots, \{d_p\}a'_p, \{1\}x])(b) \right\rfloor \quad (4.7)$$

By hypothesis we have $\left|\frac{\beta}{0}\right| b$, hence by Lemma 6 we have

$$\left|\frac{\beta}{0}\right| b \cdot_0 (n-1)[\{d_1\}a'_1(b), \dots, \{d_p\}a'_p(b), \{1\}b] \quad (4.8)$$

Hence, using (4.6) for all $i \in \{1, \dots, p\}$, (4.7) and (4.8) (adjusted to $\left|\frac{\alpha-1+\beta}{0}\right|$ by Lemma 4) we deduce by RED that

$$\left|\frac{\alpha+\beta}{0}\right| n[\{d_1\}a'_1(b), \dots, \{d_p\}a'_p(b), \{1\}b]$$

Which is what was required.

- **CUT.** Let us suppose $\left|\frac{\alpha+\gamma}{0}\right| a$ is obtained by CUT, hence a has the form $a_1 \cdot_0 a_2$. Let us suppose that a' has the form $(a'_1 \cdot_0 a'_2) \cdot_1 x$, with $a_i = a'_i(\emptyset)$, again the case where x is a child of the root encompasses the other. The premises of CUT are then $\left|\frac{\alpha}{0}\right| a_1$ and $\left|\frac{\gamma}{0}\right| a_2$. We also have $(a'_1 \cdot_1 x)(\emptyset) = a_1$, therefore the IH on $\left|\frac{\alpha}{0}\right| a_1$ along with $\left|\frac{\beta}{0}\right| b$ implies that $\left|\frac{\alpha+\beta}{0}\right| a'_1(b) \cdot_1 b$ and all that remains is to substitute $a'_2(b)$ in $(a'_1(b) \cdot_1 b) \cdot_0 x$. But since the type of x is 0, Lemma 6 implies that $\left|\frac{\alpha+\beta}{0}\right| (a'_1(b) \cdot_1 b) \cdot_0 a'_2(b)$, which concludes since trees are considered up to permutation. \square

Lemma 12 (Base cut elimination lemma). If $\left|\frac{\alpha}{1}\right| a$, then $\left|\frac{\alpha}{0}\right| a$.

Proof. By induction on the witness tree for $\left|\frac{\alpha}{1}\right| a$.

- **BASE.** Trivial.
- **RED.** Suppose a has the form $n[\{d_1\}a_1, \dots, \{d_p\}a_p]$. The premises of RED are that for all $i \in \{1, \dots, p\}$ we have $\left|\frac{\alpha-1}{1}\right| a_1 \cdot_{d_i-1} (n-1)\{d_1\}a_1, \dots, \{d_p\}a_p]$ and $\left|\frac{\alpha-1}{1}\right| (n-1)\{d_1\}a_1, \dots, \{d_p\}a_p]$. The result is then trivial by IH and RED.
- **CUT.** Suppose $a = a_1 \cdot_d a_2$ with $d \leq 1$, the premises of CUT are that $\left|\frac{\alpha}{1}\right| a_1$ and $\left|\frac{\beta}{1}\right| a_2$. If $d = 0$, then the result is trivial by the IH and CUT. If $d = 1$, we just apply Lemma 11 instead of CUT. \square

From all of this put together, we deduce the main theorem of this section.

Theorem 2 (Upper bound). If $\text{ord}(a), \text{depth}(a), \max(a) \geq 1$, $\mathcal{N}(a) \leq 2^{\frac{\text{depth}(a) \log(\max(a)+1)}{\text{ord}(a)-1}}$.

Proof. Let us set $a' = \mathcal{T}(\text{depth}(a), \text{ord}(a), \max(a))$. By definition, we have $\text{depth}(a') = \text{depth}(a)$, $\text{ord}(a') = \text{ord}(a)$ and $\max(a') = \max(a)$. Moreover by Proposition 4, we have:

$$\mathcal{N}(a) \leq \mathcal{N}(a')$$

Write $o = \text{ord}(a')$, $d = \text{depth}(a')$, $m = \max(a')$. By Lemma 9, we have $\left|\frac{\prod_{n \in \text{nodes}(a')}(n+1)}{o-1}\right| a'$. But all nodes in a' have the same label m and there are d nodes, therefore $\left|\frac{(m+1)^d}{o-1}\right| a'$. By $o-2$ applications of Lemma 8 it follows that $\left|\frac{2^{d \log(m+1)}}{o-1}\right| a'$, so $\left|\frac{2^{d \log(m+1)}}{o-1}\right| a'$ as well by Lemma 12. Finally, $\mathcal{N}(a) \leq \mathcal{N}(a') \leq 2^{\frac{d \log(m+1)}{o-1}}$ by Lemma 10. \square

This upper bound is optimal, since it yields bounds on linear head reduction that we will prove optimal in the next section.

4.3. On game situations. Before we conclude this section, let us mention a specialized form of our result of special importance to the previous section.

Theorem 3. If $n, p \geq 1$ and $d \geq 3$, then $\mathcal{N}(n[\{d\}p]) \leq 2^{\frac{2^{p^{n+1}-1}-1}{p-1}}$.

Proof. We show by induction on $n \geq 1$ that $\left\lfloor \frac{1+\sum_{k=1}^n 2p^k}{d-2} n[\{d\}p] \right\rfloor$. For $n = 1$, we need to show that $\left\lfloor \frac{1+2p}{d-2} 1[\{d\}p] \right\rfloor$. By RED, this amounts to $\left\lfloor \frac{2p}{d-2} 0[\{d\}p] \right\rfloor$ (true by BASE) and:

$$\left\lfloor \frac{2p}{d-2} p[\{d-1\}0[\{d\}p]] \right\rfloor$$

This follows from Lemma 7 from $\left\lfloor \frac{p}{d-2} p \right\rfloor$ and $\left\lfloor \frac{1}{d-2} 0[\{d\}p] \right\rfloor$, where the former follows from p instances of RED and one BASE and the latter follows from BASE.

Now, suppose $\left\lfloor \frac{1+\sum_{k=1}^{n-1} 2p^k}{d-2} (n-1)[\{d\}p] \right\rfloor$. The skeleton $n[\{d\}p]$ has one possible reduction:

$$n[\{d\}p] \rightsquigarrow p[\{d-1\}(n-1)[\{d\}p]]$$

We know by p instances of RED and one of BASE that $\left\lfloor \frac{p}{d-2} p \right\rfloor$. Therefore by Lemma 7:

$$\left\lfloor \frac{p(2+\sum_{k=1}^{n-1} 2p^k)}{d-2} p[\{d-1\}(n-1)[\{d\}p]] \right\rfloor$$

We also have $\left\lfloor \frac{p(2+\sum_{k=1}^{n-1} 2p^k)}{d-2} (n-1)[\{d\}p] \right\rfloor$ by IH and Lemma 4. Therefore by RED, $\left\lfloor \frac{1+\sum_{k=1}^n 2p^k}{d-2} n[\{d\}p] \right\rfloor$, which concludes the induction.

From this, we use as before $d-3$ times Lemma 8, Lemma 12 and Lemma 10, we deduce:

$$\mathcal{N}(n[\{d\}p]) \leq 2^{\frac{1+\sum_{k=1}^n 2p^k}{d-3}}$$

And finally $1 + \sum_{k=1}^n 2p^k = 2^{\frac{p^{n+1}-1}{p-1}} - 1$, yielding the announced result. \square

This result is particularly relevant in *game situations*: when studying the reduction length of one η -expanded Böhm tree applied to another. It also provides the answer to the question raised in the previous section about the possible length of bounded visible pointer structures — and hence of interactions between bounded strategies.

Remark 1. We finish this section by a few remarks on the above result:

- For $d = 2$, it is easy to see that $\mathcal{N}(n[\{2\}p]) = 2n$.
- For $d = 3$, experiments with an implementation of skeletons and their reductions suggest that, for $n \geq 0$ and $p \geq 2$, $\mathcal{N}(n[\{3\}p]) = 2^{\frac{p^{n+1}-1}{p-1}}$. This quantity is $\Theta(2^{(n-1)\log(p)})$ whereas our general bound predicts $\Theta(2^{n\log(p)})$. They differ but do match up to an exponential, being both of the form $2^{\Theta(n\log(p))}$.

In fact for any $d \geq 3$ we have $\mathcal{N}(n[\{d\}p]) = 2^{\frac{\Theta(n\log(p))}{d-2}}$. The upper bound is our theorem above, and the lower bound is provided by the reduction on skeletons corresponding to the visible pointer structures used in the proof of Theorem 1. So, in this sense our result is optimal on game situations, just as the upper bound of Theorem 2 will appear later to be optimal in the general situation.

5. SKELETONS AND LINEAR HEAD REDUCTION

Although it is generally understood that game-theoretic interaction (underlying interaction skeletons) has a strong operational content, the game-theoretic toolbox lacks results making this formal. One notable exception is the result of Danos, Herbelin and Regnier [9] already mentioned, which describes a step-by-step correspondence between the linear head reduction sequence of a game situation $M N_1 \dots N_n$ (where M, N_1, \dots, N_n are β -normal and η -long) and the interaction of the corresponding strategies. Along with Theorem 3, this connection suffices to immediately deduce an (optimal) upper bound for the length of reduction sequences on game situations. However, this reasoning has two drawbacks. Firstly it is rather indirect: the link it provides between linear head reduction and interaction skeletons, two relatively simple combinatorial objects, is obfuscated by the variety of mathematical notions involved. Indeed this connection requires elaborate semantic notions such as visible strategies and pointer structures, and third party results such as the (very technical) result of [9]. Secondly it only covers game situations, and it is not clear how to obtain from that general results on arbitrary terms.

In this final section we address these two points and proceed to analyse the direct connection between interaction skeletons and syntactic reduction. This study culminates in optimal upper bounds to the length of linear head reduction sequence on arbitrary simply-typed λ -terms. This requires us, on the one hand, to construct a generalization of game situations whose reduction follows the combinatorics of interaction skeletons and, on the other hand, to show that one can *compile* arbitrary terms into these generalized situations in a way allowing us to obtain our upper bounds.

In Subsection 5.1 we give the definition of linear head reduction and prove some basic properties. In Subsection 5.2 we define and study generalized game situations, and in Subsection 5.3 we prove the technical core of this section: the fact that lhr on generalized game situations can be simulated within interaction skeletons. Finally, Subsections 5.4 and 5.5 are devoted to dealing respectively with η -expansion and with λ -lifting in order to compile arbitrary terms to generalized game situations and deduce our results.

5.1. Linear head reduction. We start by recalling the definition of linear head reduction and proving some basic properties that are folklore, but to our knowledge unpublished under this formulation. Our notion of linear head reduction follows [9]. We use it rather than the more elegant approach of Accattoli [2] because we believe it yields a more direct relationship with games. Indeed the *multiplicative* reductions of Accattoli’s calculus have no counterpart in games/skeletons, which only take into account the variable substitutions.

5.1.1. Definition of linear head reduction. This work focuses strongly on *linear substitution*, for which only one variable occurrence is substituted at a time. In this situation, it is convenient to have a distinguished notation for particular *occurrences* of variables. We will use the notations x_0, x_1, \dots to denote particular occurrences of the same variable x in a term M . When in need of additional variable identifiers, we will use x^1, x^2, \dots . Sometimes, we will still denote occurrences of x by just x when their index is not relevant. If x_0 is a specific occurrence of x , we will use $M[N/x_0]$ for the substitution of x_0 by N , leaving all other occurrences of x unchanged.

Intuitively, lhr proceeds as follows. We first locate the head variable occurrence, *i.e.* the leftmost variable occurrence in the term M . Then we locate the abstraction, if any, that

binds this variable. Then we locate (again if it exists) the subterm N of M in argument position for that abstraction, and we substitute the head occurrence by N . We touch neither the other occurrences of x nor the redex. It is worth noting that locating the argument subterm can be delicate, as it is not necessarily part of a β -redex. For instance in $(\lambda y^A. (\lambda x^B. x_0 M)) N_1 N_2$, we want to replace x_0 by N_2 , even though N_2 is not directly applied to $\lambda x^B. x_0 M$. Therefore, the notion of redex will be generalized.

Note that a term is necessarily of the form $* M_1 \dots M_n, x_0 M_1 \dots M_n, \lambda x. M$ or $(\lambda x. M) M_1 \dots M_n$. This will be used quite extensively to define and reason on lhr. The **length** of a term M is the number of characters in M , *i.e.* $l(*) = 1, l(x_0) = 1, l(\lambda x. M) = l(M) + 1, l(M_1 M_2) = l(M_1) + l(M_2)$. Its **height** is $h(*) = 0, h(x_0) = 1, h(\lambda x. M) = h(M), h(M_1 M_2) = \max(h(M_1), h(M_2)) + 1$.

Definition 10. Given a term M , we define its set of **prime redexes**. They are written as pairs $(\lambda x, N)$ where N is a subterm of M , and λx is used to denote the (if it exists, necessarily unique by Barendregt's convention) subterm of M of the form $\lambda x. N'$. We define the prime redexes of M by induction on its length, distinguishing several cases depending on the form of M .

- If $* M_1 \dots M_n$ has no prime redex,
- If $x_0 M_1 \dots M_n$ has no prime redex,
- The prime redexes of $\lambda x. M'$ are those of M' ,
- The prime redexes of $(\lambda x. M') M_1 \dots M_n$, are $(\lambda x, M_1)$ plus those of $M' M_2 \dots M_n$.

The **head occurrence** of a term M is the leftmost occurrence of a variable or constant in M . If $(\lambda x, N)$ is a prime redex of M whose head occurrence is an occurrence x_0 of the variable x , then the **linear head reduct** of M is $M' = M[N/x_0]$. We write $M \rightarrow_{\text{lhr}} M'$.

Example 1. As an example, we give the lhr sequence of the term $(\lambda f. \lambda x. f (f x)) (\lambda y. y) *$.

$$\begin{aligned}
(\lambda f. \lambda x. f (f x)) (\lambda y. y) * &\rightarrow_{\text{lhr}} (\lambda f. \lambda x. (\lambda z. z) (f x)) (\lambda y. y) * \\
&\rightarrow_{\text{lhr}} (\lambda f. \lambda x. (\lambda z. f x) (f x)) (\lambda y. y) * \\
&\rightarrow_{\text{lhr}} (\lambda f. \lambda x. (\lambda z. (\lambda u. u) x) (f x)) (\lambda y. y) * \\
&\rightarrow_{\text{lhr}} (\lambda f. \lambda x. (\lambda z. (\lambda u. x) x) (f x)) (\lambda y. y) * \\
&\rightarrow_{\text{lhr}} (\lambda f. \lambda x. (\lambda z. (\lambda u. *) x) (f x)) (\lambda y. y) *
\end{aligned}$$

At this point the reduction stops since the head occurrence is a constant.

Given a term M , we overload the notation \mathcal{N} and write $\mathcal{N}(M)$ for the length of the lhr sequence of M . It is straightforward to see that lhr is compatible with β -reduction, in the sense that if $M \rightarrow_{\text{lhr}} M'$ we have $M \equiv_{\beta} M'$. Since redexes for lhr are not necessarily β -redexes, it will be necessary to consider the following generalization of redexes:

Definition 11 (Generalized redex). The **generalized redexes** of a term M are the prime redexes of all subterms of M . In particular, all prime redexes are generalized redexes.

Example 2. Consider the following λ -term:

$$M = (\lambda x. x) ((\lambda y. (\lambda z. u)) v w)$$

The only prime redex of M is $(\lambda x, (\lambda y. (\lambda z. u)) v w)$. The two other generalized redexes are $(\lambda y, v)$, which is also a β -redex, and $(\lambda z, w)$, which is not.

5.1.2. *Relating lhr with β -reduction.* Before we go on to relating lhr with β -reduction, let us include some preliminary investigations on its basic properties, and in particular how it relates to β -reduction. Consider λ -terms temporarily extended by a term \boxed{M} , for each term M . Those boxes can be opened when in head position. Formally, a head context is:

$$H[] = [] \mid H[] N \mid \lambda x. H[]$$

The additional reduction rule for opening boxes is then (where H is a head context):

$$H[\boxed{M}] \rightsquigarrow_{pop} H[M]$$

We extend lhr in the presence of \square by defining the prime redexes of M by induction on its length, as follows:

- $* M_1 \dots M_n, x_0 M_1 \dots M_n$ and $\boxed{M'} M_1 \dots M_n$ have no prime redex,
- The prime redexes of $\lambda x. M'$ are those of M' ,
- The prime redexes of $(\lambda x. M') M_1 \dots M_n$ are $(\lambda x, M_1)$ plus those of $M' M_2 \dots M_n$.

Writing $\square^{-1}(M)$ for the term obtained from M by removing all boxes, we see immediately that the boxes can only block prime redexes, so prime redexes of M are always in $\square^{-1}(M)$.

Writing $M \rightarrow_{\square^{-1}} \square^{-1}(M)$, we also notes that within this calculus, β -reduction on standard terms decomposes in two steps:

1. *Substitution* ($\beta\square$) $(\lambda x. M)N \rightsquigarrow_{\beta\square} M[\boxed{N}]/x$
2. *Unboxing.* (\square^{-1}) In one step, we remove all boxes in M .

Intuitively, boxes introduce a *delay* in the processing of β -reduction, and this delay corresponds exactly to the delay of lhr with respect to head β -reduction. Here, we exploit this fact to compare the two of them. In this Section 5.1.2 (but only in this section), terms are in this extended language unless otherwise specified. If M does not contain any boxes, we say it is a **standard** λ -term.

Lemma 13. If M, M' are standard terms such that $M \rightarrow_{lhr} M', N$ is a standard term, and the head occurrence x_0 of M is not an occurrence of y , then $M[\boxed{N}]/y \rightarrow_{lhr} M'[\boxed{N}]/y$.

Proof. Straightforward by induction on the length of M . □

Lemma 14. Suppose M is a standard term with a head occurrence x_0 of a variable x , and that $M \rightarrow_{\beta\square} M'$. Then for any standard term N such that no free variable of N is bound in M , we also have $M[N/x_0] \rightarrow_{\beta\square} M'[N/x_0]$.

Proof. Straightforward by induction on M . □

Write $\rightarrow_{lhr \vee pop}$ for exactly one reduction step of, either lhr, or *pop*.

Lemma 15. If M is a standard λ -term with $M \rightarrow_{\beta\square} N$ and $M \rightarrow_{lhr} M'$, then there is $N \rightarrow_{lhr \vee pop} N'$ such that $M' \rightarrow_{\beta\square}^* N'$. Moreover, if $M \rightarrow_{\beta\square} N$ is not a weak head reduction (so if it not a head reduction or operates under a lambda), none of the $M' \rightarrow_{\beta\square}^* N'$ is a weak head reduction.

Proof. By induction on the length of M , detailing only the case $M = (\lambda x. M') M_1 \dots M_n$.

- If $M = (\lambda x. M') M_1 \dots M_n$ where the head occurrence x_0 is an occurrence of x and the $\beta\square$ -reduction is:

$$(\lambda x. M') M_1 \dots M_n \rightarrow_{\beta\square} M'[\boxed{M_1}]/x] M_2 \dots M_n$$

In $M'[\overline{M_1}/x] M_2 \dots M_n$, one copy of $\overline{M_1}$ has replaced the occurrence x_0 , so there is $\overline{M_1}$ in head position. It follows that:

$$M'[\overline{M_1}/x] M_2 \dots M_n \rightarrow_{pop} M'[M_1/x_0][\overline{M_1}/x] M_2 \dots M_n$$

But we also have $(\lambda x. M'[M_1/x_0]) M_1 \dots M_n \rightarrow_{\beta\Box} M'[M_1/x_0][\overline{M_1}/x] M_2 \dots M_n$, so we set $N' = M'[M_1/x_0][\overline{M_1}/x] M_2 \dots M_n$.

- If $M = (\lambda x. M') M_1 \dots M_n$ whose head occurrence x_0 is an occurrence of x , and the $\beta\Box$ -reduction is inside M' . Then no free variable in M_1 can be bound in $\lambda x. M'$ by Barendregt's convention. Therefore, by Lemma 14, we have

$$M'[M_1/x_0] \rightarrow_{\beta\Box} M''[M_1/x_0]$$

therefore $(\lambda x. M'[M_1/x_0]) M_1 \dots M_n \rightarrow_{\beta\Box} (\lambda x. M''[M_1/x_0]) M_1 \dots M_n$.

- If $M = (\lambda x. M') M_1 \dots M_n$ where the head occurrence x_0 is an occurrence of x , and the $\beta\Box$ -reduction is $M_1 \rightarrow_{\beta\Box} M'_1$. Then we have:

$$(\lambda x. M'[M_1/x_0]) M_1 \dots M_n \rightarrow_{\beta\Box^2} (\lambda x. M'[M'_1/x_0]) M'_1 \dots M_n$$

providing the necessary commutation.

- If $M = (\lambda x. M') M_1 \dots M_n$ where the head occurrence x_0 is an occurrence of x , and the $\beta\Box$ -reduction is $M_i \rightarrow_{\beta\Box} M'_i$ with $i \geq 2$, then the commutation is trivial.
- If $M = (\lambda y. M') M_1 \dots M_n$ where the head occurrence x_0 is not an occurrence of y , and the $\beta\Box$ -reduction is:

$$(\lambda y. M') M_1 \dots M_n \rightarrow_{\beta\Box} M'[\overline{M_1}/y] M_2 \dots M_n$$

Necessarily, we have $M' M_2 \dots M_n \rightarrow_{\text{lhr}} M'' M_2 \dots M_n$. By Lemma 13, it follows that $(M' M_2 \dots M_n)[\overline{M_1}/y] \rightarrow_{\text{lhr}} (M'' M_2 \dots M_n)[\overline{M_1}/y]$. Since y can only appear in M' and M'' by Barendregt's convention, it follows that:

$$M'[\overline{M_1}/y] M_2 \dots M_n \rightarrow_{\text{lhr}} M''[\overline{M_1}/y] M_2 \dots M_n$$

which provides the required commutation.

- If $M = (\lambda y. M') M_1 \dots M_n$ where the head occurrence x_0 is not an occurrence of y , and the $\beta\Box$ -reduction is $M' \rightarrow_{\beta\Box} M''$, it follows directly from IH.
- If $M = (\lambda y. M') M_1 \dots M_n$ where the head occurrence x_0 is not an occurrence of y , and the $\beta\Box$ -reduction is $M_1 \rightarrow_{\beta\Box} M'_1$, trivial.
- If $M = (\lambda y. M') M_1 \dots M_n$ where the head occurrence x_0 is not an occurrence of y , and the $\beta\Box$ -reduction is $M_i \rightarrow_{\beta\Box} M'_i$ with $i \geq 2$. By definition of lhr:

$$M' M_2 \dots M_n \rightarrow_{\text{lhr}} M'' M_2 \dots M_n$$

but we also have $M' M_2 \dots M_n \rightarrow_{\beta\Box} M' M_2 \dots M'_i \dots M_n$. By IH, there is $M' M_2 \dots M'_i \dots M_n \rightarrow_{\text{lhr}} N$ such that $M'' M_2 \dots M_n \rightarrow_{\beta\Box}^* N$. But necessarily, N must have the form $S M_2 \dots M'_i \dots M_n$ since we have $M' M_2 \dots M'_i \dots M_n \rightarrow_{\text{lhr}} N$. So, $M'' M_2 \dots M_n \rightarrow_{\beta\Box}^* S M_2 \dots M'_i \dots M_n$. But the original $\beta\Box$ -reduction was not a weak head reduction, so by IH none of the reductions in $\rightarrow_{\beta\Box}^*$ is. So they must all either operate inside of M'' or M_i : in other words $M'' \rightarrow_{\beta\Box}^* S$ and $M_i \rightarrow_{\beta\Box}^* M'_i$. It follows that: $(\lambda y. M'') M_1 \dots M_n \rightarrow_{\beta\Box}^* (\lambda y. S) M_1 \dots M'_i \dots M_n$, providing the required commutation. \square

Lemma 16. Suppose $M \rightarrow_{\text{lhr}\vee\text{pop}} N$, $M \rightarrow_{\square^{-1}} M'$ and $N \rightarrow_{\square^{-1}} N'$. Then,

- If $M \rightarrow_{\text{pop}} N$, then $M' = N'$.
- If $M \rightarrow_{\text{lhr}} N$, then $M' \rightarrow_{\text{lhr}} N'$.

Proof. Straightforward. □

Proposition 5. Linear head reduction terminates on standard terms.

Proof. Note first that from Lemma 15 it follows that whenever $M \rightarrow_{\beta\square}^* N$ and $M \rightarrow_{\text{lhr}} M'$, then there exists N' such that $N \rightarrow_{\text{lhr}\vee\text{pop}} N'$ and $M' \rightarrow_{\beta\square}^* N'$.

Suppose standard M has an infinite reduction chain:

$$M = M_0 \rightarrow_{\text{lhr}} M_1 \rightarrow_{\text{lhr}} M_2 \rightarrow_{\text{lhr}} \dots$$

and $M \rightarrow_{\beta\square} M'$. Then, by iterating the argument above, we get an infinite chain:

$$M' = M'_0 \rightarrow_{\text{lhr}\vee\text{pop}} M'_1 \rightarrow_{\text{lhr}\vee\text{pop}} M'_2 \rightarrow_{\text{lhr}\vee\text{pop}} \dots$$

Note that in this chain, there cannot be an infinite succession of *pop*, because each *pop* strictly decreases the length of terms. Therefore, there are an infinite number of \rightarrow_{lhr} in this sequence. Now, define $M' \rightarrow_{\square^{-1}} M''$. Likewise for each $i \in \mathbb{N}$, define $M'_i \rightarrow_{\square^{-1}} M''_i$. By Lemma 16, if $M'_i \rightarrow_{\text{pop}} M'_{i+1}$ we have $M''_i = M''_{i+1}$, whereas if $M'_i \rightarrow_{\text{lhr}} M'_{i+1}$, we still have $M''_i \rightarrow_{\text{lhr}} M''_{i+1}$; it follows that there is a chain:

$$M'' = M''_0 \rightarrow_{\text{lhr}\vee=} M''_1 \rightarrow_{\text{lhr}\vee=} M''_2 \rightarrow_{\text{lhr}\vee=} \dots$$

Since there is an infinite number of *lhr* in the sequence of M' , there is an infinite number of *lhr* in this sequence. Moreover since $M \rightarrow_{\beta\square} M' \rightarrow_{\square^{-1}} M''$, we have $M \rightarrow_{\beta} M''$. Therefore, we have proved that if the *lhr* reduction chain of M is infinite and $M \rightarrow_{\beta} M'$, then the *lhr* reduction chain of M' is infinite as well. But by normalization of β -reduction, we know that $M \rightarrow_{\beta}^* K$, where K is β -normal. But if K has no β -redex it has no prime redexes either, so K has no *lhr* reduction; absurd. Therefore, the *lhr* reduction chain of M was finite. □

Lemma 17. Let M be a standard term with a head occurrence x_0 , such that there is no prime redex $(\lambda x, N)$ within M . Let S be another term, and y a free variable in M . Then, there is still no prime redex $(\lambda x, N)$ in $M[S/y]$.

Proof. Straightforward by induction on the length of M . □

Lemma 18. If M is standard, normal for *lhr* and $M \rightarrow_{\beta} M'$, then M' is normal for *lhr*.

Proof. By induction on the length of M . We only detail $M = (\lambda x. M') M_1 \dots M_n$, the other cases being trivial. Several subcases arise depending on the location of the β -reduction; the only non-trivial case is $M \rightarrow_{\beta} M'[M_1/x] M_2 \dots M_n$. If the head occurrence of M is a constant $*$, it is still the head occurrence of $M'[M_1/x] M_2 \dots M_n$, which is therefore *lhr*-normal. Otherwise it is a variable occurrence y_0 , such that there is no prime redex $(\lambda y, N)$ in M . It follows by definition of prime redexes that there is no prime redex either in $M' M_2 \dots M_n$. By Lemma 17, there is still no prime redex in $(M' M_2 \dots M_n)[M_1/x] = M'[M_1/x] M_2 \dots M_n$, but its head occurrence has not changed, so it is *lhr*-normal. □

Proposition 6. If a standard $M \rightarrow_{\beta} M'$, then $\mathcal{N}(M) \geq \mathcal{N}(M')$.

Proof. We decompose $M \rightarrow_{\beta\Box} M'' \rightarrow_{\Box^{-1}} M'$. Take a maximal lhr sequence:

$$M = M_0 \rightarrow_{\text{lhr}} M_1 \rightarrow_{\text{lhr}} \cdots \rightarrow_{\text{lhr}} M_n$$

by Lemma 15, there is a corresponding sequence of the same length:

$$M'' = M''_0 \rightarrow_{\text{lhr}\vee\text{pop}} M''_1 \rightarrow_{\text{lhr}\vee\text{pop}} \cdots \rightarrow_{\text{lhr}\vee\text{pop}} M''_n$$

with for all $i \in \{0, \dots, n\}$, $M_i \rightarrow_{\beta\Box}^* M''_i$. Since M_n is maximal, this means that its head occurrence does not appear in a prime redex, or is a constant. By unboxing, we get:

$$M' = M'_0 \rightarrow_{\text{lhr}\vee=} M'_1 \rightarrow_{\text{lhr}\vee=} \cdots \rightarrow_{\text{lhr}\vee=} M'_n$$

We have $M_n \rightarrow_{\beta}^* M'_n$ by construction, so by Lemma 18 we know that M'_n is lhr-normal as well, so this chain is maximal. So, the lhr sequence of M' is the subsequence of $M'_0 \rightarrow_{\text{lhr}\vee=} M'_1 \rightarrow_{\text{lhr}\vee=} \cdots \rightarrow_{\text{lhr}\vee=} M'_n$ keeping only the lhr steps, so its length is less or equal than n . \square

5.2. Generalized game situations. In this section, we detail the situation where the connection between lhr and skeletons is the most direct.

That involves two aspects. Firstly, since the game-theoretic interpretation of terms is invariant under η -expansion, it is clear from the start that terms whose dynamics match game-theoretic interaction should be η -expanded. Accordingly, we will give a new notion of non β -normal η -long terms adapted to lhr. Secondly, game-theoretic interaction is also *local*: strategies involved in the interpretation of a term will only communicate with other strategies with whom they share a redex, whereas substitution is non-local and can span over multiple redexes. Accordingly, we introduce a notion called *local scope* ensuring that the information flow of lhr is local. With the help of these two conditions we define *generalized game situations*, that we prove to elegantly connect to interaction skeletons.

We now investigate these two aspects, in turn.

5.2.1. Generalized η -long terms. Non β -normal η -long terms are often defined as those for which any further η -expansion creates new β -redexes. Here of course, since we work with generalized β -redexes, we will use instead the following definition:

Definition 12. A term M is η -**long** if whenever $M \rightarrow_{\eta} M'$, then M' has more generalized β -redexes than M .

We now prove that η -long terms are preserved by lhr. For that, we will make use of the following lemmas allowing us to compose and decompose η -long terms.

Lemma 19. If M is η -long, so are its subterms.

Proof. Straightforward by induction on M , since an η -expansion in a subterm N of M would create a new generalized redex in M that necessarily has to be in N . \square

Lemma 20. A term $(\lambda x. M) M_1 \dots M_n$ is η -long, iff so are $M M_2 \dots M_n$ and M_1 .

Proof. Straightforward by cases on the location of an η -expansion. \square

Lemma 21. If $\Gamma \vdash M : A \rightarrow B$ and $\Gamma \vdash N : A$ are η -long, so is $M N$.

Proof. Straightforward by induction on the length of M . \square

Using the lemmas above we prove the following substitution lemma, crucial in proving the stability of η -long terms by lhr.

Lemma 22. If $\Gamma \vdash M : A$ is η -long and with head occurrence x_0 of a variable $x : B$, and $\Gamma \vdash N : B$ is η -long, then $M[N/x_0]$ is η -long.

Proof. By induction on the length of M .

- If $M = x_0 M_1 \dots M_n$, then necessarily each M_i is η -long by Lemma 19. It follows by Lemma 21 that $N M_1 \dots M_n$ is η -long as well.
- If $M = \lambda y. M'$, it follows directly from the IH.
- If $M = (\lambda y. M') M_1 \dots M_n$, it follows directly from IH and Lemma 20. \square

Lemma 23. If M is η -long and $M \rightarrow_{\text{lhr}} M'$, then M' is η -long.

Proof. By induction on the length of M . We skip all the trivial cases, and only detail $M = (\lambda x. M') M_1 \dots M_n$. Then, two subcases:

- If the head occurrence of M is an occurrence x_0 of x , then we have: $M \rightarrow_{\text{lhr}} M[M_1/x_0]$. But since M is η -long, M_1 is η -long as well by Lemma 19. So, by Lemma 22 we have that $M[M_1/x_0]$ is η -long as well.
- If the head occurrence of M is an occurrence y_0 of a variable or constant other than x , then if $M \rightarrow_{\text{lhr}} M[N/y_0]$, for some subterm N of M , and we must have as well:

$$M' M_2 \dots M_n \rightarrow_{\text{lhr}} M'[N/y_0] M_2 \dots M_n$$

by IH, $M'[N/y_0] M_2 \dots M_n$ is η -long. Finally, it follows from Lemma 20 that M_1 is η -long and that recomposing $(\lambda x. M'[N/y_0]) M_1 \dots M_n$ yields an η -long term. \square

5.2.2. *Local scope.* Now that we have a notion of η -long term stable under composition, let us consider the syntactic counterpart of the second aspect of skeletons: that their reduction is local. It is not clear at first what *local* means in this context: just like a game situation consists in two η -long normal forms interacting, a generalized game situation will consist in a “tree” of η -long normal forms. Let us start with this slightly naive definition:

Definition 13. A term M is **strongly locally scoped** (abbreviated sls) iff for any generalized redex $(\lambda x, N)$ in M , N is closed.

Unfortunately, sls terms do not quite fit for a syntactic counterpart of skeletons: they are not preserved by lhr. It is easy to find a counter-example, for instance:

$$(\lambda y. (\lambda x. x y) (\lambda z. z)) * \rightarrow_{\text{lhr}} (\lambda y. (\lambda x. (\lambda z. z) y) (\lambda z. z)) *$$

Here, a new generalized redex $(\lambda z, y)$ is formed where y is obviously not closed. Therefore, we must make skeletons correspond instead with a generalization of sls terms preserved by lhr. This generalization comes from the observation that in the right hand side term above, the violation of strong local scope is mitigated by the fact that the violating variable y is part of a generalized redex — so its value is somehow already provided by an environment. Hence the following definition:

Definition 14. A variable x in M is **active** iff it is a free variable or if there is a generalized redex $(\lambda x, N)$ in M . It is **passive** otherwise. A term M is **locally scoped** (abbreviated ls) if for any generalized redex $(\lambda x, N)$ in M all the free variables in N are active in M .

Local scope will be sufficient to ensure that the interpretation to skeletons is a simulation, but the correspondence between terms and skeletons will be tighter for sls terms: the tree structure of the skeleton will match the tree structure of nested generalized redexes.

Of course, we now need to prove that locally scoped terms are preserved by lhr. However, this is still not true! Indeed, consider the following reduction:

$$\lambda y. (\lambda x. x y) (\lambda z. z) \rightarrow_{\text{lhr}} \lambda y. (\lambda x. (\lambda z. z) y) (\lambda z. z)$$

The left hand side term is (strongly) locally scoped, but the right hand side term is not because y is passive, but appears in the argument of a generalized redex. However, the problem disappears if we apply the two terms above to a constant $*$. In general, we will show that *closed* locally scoped terms of *ground type* are closed under lhr. This may seem like a big restriction but it is not: an arbitrary term can be made closed and of ground type without changing its possible reduction sequences significantly, by replacing its free variables with constants and applying it to as many constants as required.

To prove stability of local scope by lhr, we start by stability under substitution.

Lemma 24. If M is a locally scoped term of ground type with head occurrence x_0 of a variable x , and N is a locally scoped term, then $M[N/x_0]$ is locally scoped.

Proof. By induction on the length of M (of ground type, so not an abstraction).

- If $M = x_0 M_1 \dots M_n$, then the generalized redexes of $N M_1 \dots M_n$ are those of N, M_1, \dots, M_n (that are of the form $(\lambda y, S)$ where all free variables in S are active, by definition of local scope), with possibly the addition of generalized redexes of the form $(\lambda z, M_i)$. Free variables in M_i are free in $N M_1 \dots M_n$, so they are active.
- If $M = (\lambda y. M') M_1 \dots M_n$, it follows directly from IH. □

Lemma 25. If $\vdash M : o$ is locally scoped and $M \rightarrow_{\text{lhr}} M'$, then M' is locally scoped.

Proof. By induction on the length of M , writing $(\lambda x, S)$ for the prime redex fired in $M \rightarrow_{\text{lhr}} M'$. We only detail the non-trivial cases.

- If $M = (\lambda y. N) M_1 \dots M_n$ with $y \neq x$, then it direct that $N M_2 \dots M_n$ is still ls. By IH, $N[S/x_0] M_2 \dots M_n$ is ls as well, and from that follows that $M[S/x_0]$ is ls.
- If $M = (\lambda x. N) M_1 \dots M_n$, where x_0 is an occurrence of x . Then, necessarily $N M_2 \dots M_n$ is ls as well. Likewise, M_1 is ls, otherwise M could not be. Therefore by Lemma 24, $N[M_1/x_0] M_2 \dots M_n$ is ls, and from that follows that M' is ls. □

We say that a term M is a **generalized game situation** if it is closed, of ground type, and both η -long and locally scoped. By Lemmas 23 and 25, we know that generalized game situations are preserved by linear head reduction.

5.3. Simulation of generalized game situations. We start this subsection by showing how one can associate a skeleton with any generalized game situation – in fact with any term, although this connection will only yield a simulation for generalized game situations.

Definition 15. Let $\Gamma \vdash M : A$ be a term, with a **bs-environment** ρ , being defined as a partial function mapping each variable x of Γ on which it is defined to a skeleton $\rho(x)$.




Figure 3: Simulation of the reduction of Example 1 in skeletons

Then the skeleton $\llbracket M \rrbracket_\rho$ is defined by induction on the length of M , as follows:

$$\begin{aligned}
 \llbracket * M_1 \dots M_n \rrbracket_\rho &= \mathbf{0} \\
 \llbracket x_0 M_1 \dots M_n \rrbracket_\rho &= 1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho && \text{if } \rho(x) \text{ undefined} \\
 \llbracket x_0 M_1 \dots M_n \rrbracket_\rho &= (1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho) \cdot 1_{\text{v}(x)+1} \rho(x) && \text{if } \rho(x) \text{ defined} \\
 \llbracket \lambda x. M \rrbracket_\rho &= \llbracket M \rrbracket_\rho \\
 \llbracket (\lambda x. M) M_1 \dots M_n \rrbracket_\rho &= \llbracket M M_2 \dots M_n \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}}
 \end{aligned}$$

We write $\llbracket M \rrbracket$ for $\llbracket M \rrbracket_\emptyset$.

5.3.1. Simulation of generalized game situations. We now prove that $\llbracket - \rrbracket$ is a simulation. Because $\llbracket - \rrbracket$ over-approximates terms it will not directly relate \rightarrow_{lhr} and \rightsquigarrow . Rather, we will have that if $M \rightsquigarrow M'$, then there is a such that $\llbracket M \rrbracket \rightsquigarrow a \leftarrow \llbracket M' \rrbracket$. This relaxed simulation will suffice for our purposes since by Lemma 2 it implies that $\mathcal{N}(\llbracket M' \rrbracket) \leq \mathcal{N}(a)$. We show in Figure 3 the skeletons corresponding with all lhr-reducts of the term $(\lambda f^{o \rightarrow o}. \lambda x^o. f (f x)) (\lambda y^o. y) *_o$ from Example 1, with explicit typing.

We now aim to prove our simulation result for generalized game situations.

Definition 16. Let M be a term, and ρ, ρ' two bs-environments. We write $\rho \sim_M \rho'$ iff $\text{dom}(\rho) \cap \text{fv}(M) = \text{dom}(\rho') \cap \text{fv}(M)$, and that for all $x \in \text{dom}(\rho) \cap \text{fv}(M)$, $\rho(x) = \rho'(x)$.

Lemma 26. If M is a term, ρ and ρ' are such that $\rho \sim_M \rho'$, then $\llbracket M \rrbracket_{\rho'} = \llbracket M \rrbracket_{\rho}$.

Proof. Straightforward by induction on the length of M . \square

Lemma 27. Suppose $\Gamma \vdash M : A$ is such that $(x : B) \in \Gamma$ does not appear in any subterm N , for any generalized redex $(\lambda y, N)$ in M . Let ρ be a bs-environment, and a skeleton a . Then, we have $\llbracket M \rrbracket_{\rho \cup \{x \mapsto a\}} \hookrightarrow \llbracket M \rrbracket_{\rho} \cdot \text{lv}(x) + 1 \ a$.

Proof. By induction on the length of M , with $\rho' = \rho \cup \{x \mapsto a\}$. Most cases follow directly by unfolding the definition of $\llbracket - \rrbracket$, IH, Lemma 3 and basic manipulations on embeddings.

We detail the one non-trivial case where $M = (\lambda y. M') M_1 \dots M_n$. Then:

$$\begin{aligned} \llbracket (\lambda y. M') M_1 \dots M_n \rrbracket_{\rho \cup \{x \mapsto a\}} &= \llbracket M' M_2 \dots M_n \rrbracket_{\rho \cup \{x \mapsto a\} \cup \{y \mapsto \llbracket M_1 \rrbracket_{\rho \cup \{x \mapsto a\}}\}} \\ &= \llbracket M' M_2 \dots M_n \rrbracket_{\rho \cup \{x \mapsto a\} \cup \{y \mapsto \llbracket M_1 \rrbracket_{\rho}\}} \\ &\hookrightarrow \llbracket M' M_2 \dots M_n \rrbracket_{\rho \cup \{y \mapsto \llbracket M_1 \rrbracket_{\rho}\}} \cdot \text{lv}(x) + 1 \ a \\ &= \llbracket (\lambda x. M') M_1 \dots M_n \rrbracket_{\rho} \cdot \text{lv}(x) + 1 \ a \end{aligned}$$

where the second equation exploits Lemma 26 and the fact that x does not appear in M_1 (since M_1 is in a generalized redex $(\lambda y, M_1)$); the embedding is by IH, and the last equality is by definition. \square

Lemma 28. Let $\Gamma \vdash M : A_1 \rightarrow \dots \rightarrow A_n \rightarrow o$ be a locally scoped η -long term, and $\Gamma \vdash N_i : A_i$ be terms. Finally, let ρ be a bs-environment on $M N_1 \dots N_n$. Then:

$$\llbracket M N_1 \dots N_n \rrbracket_{\rho} \hookrightarrow \llbracket M \rrbracket_{\rho} \cdot \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho} \mid 1 \leq i \leq n$$

Proof. By induction on the length of M , detailing non-trivial cases.

- If $M = \lambda x. M'$, we calculate:

$$\begin{aligned} \llbracket (\lambda x. M') N_1 \dots N_p \rrbracket_{\rho} &= \llbracket M' N_2 \dots N_p \rrbracket_{\rho \cup \{x \mapsto \llbracket N_1 \rrbracket_{\rho}\}} \\ &\hookrightarrow \llbracket M' \rrbracket_{\rho \cup \{x \mapsto \llbracket N_1 \rrbracket_{\rho}\}} \cdot \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho \cup \{x \mapsto \llbracket N_2 \rrbracket_{\rho}\}} \mid 2 \leq i \leq p \\ &= \llbracket M' \rrbracket_{\rho \cup \{x \mapsto \llbracket N_1 \rrbracket_{\rho}\}} \cdot \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho} \mid 2 \leq i \leq p \\ &\hookrightarrow \llbracket M' \rrbracket_{\rho} \cdot \text{lv}(x) + 1 \ \llbracket N_1 \rrbracket_{\rho} \cdot \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho} \mid 2 \leq i \leq p \\ &= \llbracket M' \rrbracket_{\rho} \cdot \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho} \mid 1 \leq i \leq p \end{aligned}$$

The first equality is by definition, the first embedding by IH, the second equality by Lemma 26 since x cannot appear in N_i , the second embedding by Lemma 27 since $\lambda x. M'$ is ls, so x , being passive, cannot appear in N for any generalized redex $(\lambda z, N)$ of M' . Finally, the last equation is by definition.

- If $M = (\lambda x. M') M_1 \dots M_p$, we calculate:

$$\begin{aligned} &\llbracket (\lambda x. M') M_1 \dots M_p N_1 \dots N_n \rrbracket_{\rho} \\ &= \llbracket M' M_2 \dots M_p N_1 \dots N_n \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_{\rho}\}} \\ &\hookrightarrow \llbracket M' M_2 \dots M_p \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_{\rho}\}} \cdot \\ &\quad \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_{\rho}\}} \mid 1 \leq i \leq p \\ &= \llbracket (\lambda x. M') M_1 \dots M_p \rrbracket_{\rho} \cdot \{ \text{lv}(N_i) + 1 \} \llbracket N_i \rrbracket_{\rho} \mid 1 \leq i \leq p \end{aligned}$$

The first equality is by definition, the embedding by IH, the last equation by Definition 15 and Lemma 26, since x does not appear in N_i . \square

In order to prove our simulation result between lhr on terms and skeletons, we will have to generalize it to a connection between skeletons and terms-with-environments, or closures. Therefore, we will make use of the following notions.

Definition 17. The notions of *closure* and *environments* are defined by mutual induction:

- A **closure** is $(\Gamma \vdash M : A, \sigma)$ with $\Gamma \vdash M : A$ a term, and σ an **environment** on Γ .
- An **environment** σ on Γ is a partial function which, when defined, maps a variable $(x : A)$ to a closure $(\Gamma \vdash N : A, \tau)$. We write $\sigma_1(x) = N$ and $\sigma_2(x) = \tau$.

When $(\Gamma \vdash M : A, \sigma)$ is a closure, we will use the shorthand notation M^σ and leave Γ, A implicit. An environment σ for Γ is **flat** if for each $(x : A) \in \Gamma$, we have that $\sigma_2(x) \subseteq \sigma$.

Standard notions on terms directly translate to environments: σ on Γ is **η -long** iff for all $(x : A) \in \Gamma$, if $\sigma(x)$ is defined then $\sigma_1(x)$ is η -long and $\sigma_2(x)$ is η -long. It is **locally scoped** iff for all $(x : A) \in \Gamma$, if $\sigma(x)$ is defined then $\sigma_1(x)$ and $\sigma_2(x)$ are locally scoped.

Using environments, we can generalize lhr on terms to a linear reduction on closures.

Lemma 29. Define the following reduction:

$$\frac{}{x_i M_1 \dots M_n \rightarrow_\sigma \sigma_1(x) M_1 \dots M_n} \qquad \frac{M \rightarrow_\sigma M'}{\lambda y. M \rightarrow_\sigma \lambda y. M'}$$

$$\frac{M' M_2 \dots M_n \rightarrow_{\sigma \cup \{y \rightarrow M_1^\sigma\}} M'' M_2 \dots M_n}{(\lambda y. M') M_1 \dots M_n \rightarrow_\sigma (\lambda y. M'') M_1 \dots M_n}$$

Then, we have $M \rightarrow_\emptyset M'$ iff $M \rightarrow_{\text{lhr}} M'$.

Proof. We prove that for all σ :

- If the head occurrence of M is bound in M , then $M \rightarrow_{\text{lhr}} M'$ iff $M \rightarrow_\sigma M'$,
- If the head occurrence x_0 of M is an occurrence of a variable x free in M on which σ is defined, then $M \rightarrow_\sigma M[\sigma_1(x)/x_0]$.
- If the head occurrence of M is a $*$ or an occurrence x_0 of x on which σ is undefined, then both \rightarrow_{lhr} and \rightarrow_σ halt.

This is done by a straightforward induction on the length of M . □

Environments have a tree structure that corresponds closely to the tree structure of interaction skeletons. The following definition associates a skeleton with any environment.

Definition 18. If σ is an environment on Γ , then we define a bs-environment $\llbracket \sigma \rrbracket$ which for any $(x : A) \in \Gamma$, (1) is undefined if $\sigma(x)$ is undefined, (2) associates $\llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma_2(x) \rrbracket}$ otherwise.

Finally, we are in position to prove our simulation result.

Proposition 7. Let $\Gamma \vdash M, M' : o$ be η -long ls terms, and σ be an η -long ls flat environment on Γ such that $M \rightarrow_\sigma M'$. Then, $\llbracket M \rrbracket_{\llbracket \sigma \rrbracket} \rightarrow_{\text{bs}} \llbracket M' \rrbracket_{\llbracket \sigma \rrbracket}$. So in particular, if $M \rightarrow_{\text{lhr}} M'$ we have $\llbracket M \rrbracket \rightarrow_{\text{bs}} \llbracket M' \rrbracket$.

Proof. By induction on the definition of \rightarrow_σ .

- If $x_0 M_1 \dots M_n \rightarrow_\sigma \sigma_1(x) M_1 \dots M_n$. Since $\sigma_1(x)$ is locally scoped and η -long, by Lemma 28 we have:

$$\llbracket \sigma_1(x) M_1 \dots M_n \rrbracket_{\llbracket \sigma \rrbracket} \hookrightarrow \llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma \rrbracket} \cdot \{ \text{lv}(M_i) + 1 \} \llbracket M_i \rrbracket_{\llbracket \sigma \rrbracket} \mid 1 \leq i \leq n \}$$

But we have $\sigma_2(x) \subseteq \sigma$ since σ is flat, from which follows $\llbracket \sigma_2(x) \rrbracket \subseteq \llbracket \sigma \rrbracket$. Moreover, $\text{fv}(\sigma_1(x)) \cap \text{dom}(\sigma) = \text{fv}(\sigma_1(x)) \cap \text{dom}(\sigma_2)$ since both are defined on all free variables of $\sigma_1(x)$. So by Lemma 26, $\llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma \rrbracket} = \llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma_2(x) \rrbracket}$. Finally, we have:

$$\llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma_2(x) \rrbracket} \cdot \{ \text{lv}(M_i) + 1 \} \llbracket M_i \rrbracket_{\llbracket \sigma \rrbracket} \mid 1 \leq i \leq n \} \hookrightarrow \llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma_2(x) \rrbracket} \cdot \text{lv}(x) \left(\prod_{i=1}^n \llbracket M_i \rrbracket_{\llbracket \sigma \rrbracket} \right)$$

Where the right hand side is a \rightsquigarrow -reduct of

$$\llbracket x_0 M_1 \dots M_n \rrbracket_{\llbracket \sigma \rrbracket} = (1 + \prod_{i=1}^n \llbracket M_i \rrbracket_{\llbracket \sigma \rrbracket}) \cdot \text{lv}(x)+1 \llbracket \sigma_1(x) \rrbracket_{\llbracket \sigma_2(x) \rrbracket}$$

- If $\lambda y. M \rightarrow_{\sigma} \lambda y. M'$, then M and M' do not have ground type.
- If $(\lambda y. M) M_1 \dots M_n \rightarrow_{\sigma} (\lambda y. M') M_1 \dots M_n$, then note that $\sigma \cup \{y \mapsto M_1^{\sigma}\}$ is still flat (and η -long, locally scoped). Then by IH we have:

$$\llbracket M M_2 \dots M_n \rrbracket_{\llbracket \sigma \cup \{y \mapsto M_1^{\sigma}\} \rrbracket} \rightarrow_{\text{bs}} \llbracket M' M_2 \dots M_n \rrbracket_{\llbracket \sigma \cup \{y \mapsto M_1^{\sigma}\} \rrbracket}$$

But $\llbracket \sigma \cup \{y \mapsto M_1^{\sigma}\} \rrbracket = \llbracket \sigma \rrbracket \cup \{y \mapsto \llbracket M_1 \rrbracket_{\llbracket \sigma \rrbracket}\}$ by definition of interpretation of environments; the required reduction follows by def. of interpretation of terms. \square

5.3.2. Relating generalized game situations and their interpretation. To estimate lhr on generalized game situations, we need to define measures on terms that reflect the geometry of the corresponding skeletons. So instead of the quantities traditionally used to evaluate the complexity of λ -terms (like the height or length), we have two alternative quantities.

Definition 19. The **depth** $\text{depth}(M)$ of M is defined by induction on the length of M :

$$\begin{aligned} \text{depth}(* M_1 \dots M_n) &= 1 \\ \text{depth}(x_0 M_1 \dots M_n) &= \max_{1 \leq i \leq n} \text{depth}(M_i) \\ \text{depth}(\lambda x. M) &= \text{depth}(M) \\ \text{depth}((\lambda x. M) M_1 \dots M_n) &= \max(\text{depth}(M M_2 \dots M_n), \text{depth}(M_1) + 1) \end{aligned}$$

Likewise, the **local height** $\text{lh}(M)$ of a term M is defined by:

$$\begin{aligned} \text{lh}(* M_1 \dots M_n) &= 0 \\ \text{lh}(x_0 M_1 \dots M_n) &= 1 + \max_{1 \leq i \leq n} \text{lh}(M_i) \\ \text{lh}(\lambda x. M) &= \text{lh}(M) \\ \text{lh}((\lambda x. M) M_1 \dots M_n) &= \max(\text{lh}(M M_2 \dots M_n), \text{lh}(M_1)) \end{aligned}$$

We now aim to prove that these indeed reflect quantities on the corresponding skeletons. Lemma 30 deals with depth, Lemma 31 with local height and Lemma 32 with order.

Lemma 30. If ρ is a bs-environment, we set $\text{depth}(\rho) = \max_{x \in \text{dom}(\rho)} \text{depth}(\rho(x))$. Then, for each strongly locally scoped term M with a bs-environment ρ , we have:

$$\text{depth}(\llbracket M \rrbracket_{\rho}) \leq \max(\text{depth}(M), \text{depth}(\rho) + 1)$$

In particular, $\text{depth}(\llbracket M \rrbracket) \leq \text{depth}(M)$.

Proof. By induction on the length of M , detailing the non-trivial cases.

- If $M = x_0 M_1 \dots M_n$ and ρ is not defined on x , then we have $\text{depth}(M) = \max_{1 \leq i \leq n} \text{depth}(M_i)$. On the other hand, $\llbracket x_0 M_1 \dots M_n \rrbracket_\rho = (1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho)$. But then we have:

$$\begin{aligned} \text{depth}(1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho) &= \max_{1 \leq i \leq n} \text{depth}(\llbracket M_i \rrbracket_\rho) \\ &\leq \max_{1 \leq i \leq n} \max(\text{depth}(M_i), \text{depth}(\rho) + 1) \\ &= \max(\text{depth}(M), \text{depth}(\rho) + 1) \end{aligned}$$

where the first equality is by definition of $+$, depth , and \bigsqcup , the inequality is by IH, and the last equality is by definition of maximum and depth .

- If $M = x_0 M_1 \dots M_n$, $\rho(x)$ defined, we still have $\text{depth}(M) = \max_{1 \leq i \leq n} \text{depth}(M_i)$. On the other hand, $\llbracket x_0 M_1 \dots M_n \rrbracket_\rho = (1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho) \cdot_{\text{iv}(x)+1} \rho(x)$. But then:

$$\begin{aligned} \text{depth}((1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho) \cdot_{\text{iv}(x)+1} \rho(x)) &= \max(\text{depth}(1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho), \text{depth}(\rho(x)) + 1) \\ &\leq \max(\max_{1 \leq i \leq n} \text{depth}(\llbracket M_i \rrbracket_\rho), \text{depth}(\rho) + 1) \\ &\leq \max(\max_{1 \leq i \leq n} \text{depth}(M_i), \text{depth}(\rho) + 1) \end{aligned}$$

where the first equality is by definition of \cdot , the first inequality is by definition of $+$, \bigsqcup and $\text{depth}(\rho)$, the second inequality is by IH and definition of \max .

- If $M = \lambda x. M'$, then it directly follows from the IH.
- If $M = (\lambda x. M') M_1 \dots M_n$, then we have

$$\text{depth}(M) = \max(\text{depth}(M' M_2 \dots M_n), \text{depth}(M_1) + 1)$$

We calculate:

$$\begin{aligned} &\text{depth}(\llbracket (\lambda x. M') M_1 \dots M_n \rrbracket_\rho) \\ &= \text{depth}(\llbracket M' M_2 \dots M_n \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}}) \\ &\leq \max(\text{depth}(M' M_2 \dots M_n), \text{depth}(\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}) + 1) \\ &\leq \max(\text{depth}(M' M_2 \dots M_n), \max(\text{depth}(\rho), \text{depth}(\llbracket M_1 \rrbracket_\rho)) + 1) \\ &= \max(\text{depth}(M' M_2 \dots M_n), \max(\text{depth}(\rho), \text{depth}(\llbracket M_1 \rrbracket_\emptyset)) + 1) \\ &\leq \max(\text{depth}(M' M_2 \dots M_n), \max(\text{depth}(\rho), \max(\text{depth}(M_1), 1)) + 1) \\ &\leq \max(\text{depth}(M' M_2 \dots M_n), \text{depth}(M_1) + 1, \text{depth}(\rho) + 1) \\ &= \max(\text{depth}((\lambda x. M') M_1 \dots M_n), \text{depth}(\rho) + 1) \end{aligned}$$

Where the first equality is by definition of interpretation, the second line is by IH, the third line is by definition of depth on environments, the fourth line uses that since M is strongly locally scoped M_1 must be closed, therefore $\rho \simeq_{M_1} \emptyset$ hence by Lemma 26 we have $\llbracket M_1 \rrbracket_\rho = \llbracket M_1 \rrbracket_\emptyset$. The fifth line is by IH on M_1 , and the last two lines are by easy manipulations on maximums (using that depth is always greater than one) and definition of depth . \square

Lemma 31. If ρ is a bs-environment, we set $\max(\rho) = \max_{x \in \text{dom}(\rho)} \max(\rho(x))$. Then, for each term M with a bs-environment ρ , for any natural number $m \in \mathbb{N}$, we have:

$$\max(m + \llbracket M \rrbracket_\rho) \leq \max(\text{lh}(M) + m, \max(\rho))$$

In particular, $\max(\llbracket M \rrbracket) \leq \text{lh}(M)$.

Proof. By induction on the length of M , omitting basic manipulations of expressions.

- If $M = * M_1 \dots M_n$, it is direct.
- If $M = x_0 M_1 \dots M_n$ with $\rho(x)$ undefined, we calculate:

$$\begin{aligned} \max(m + \llbracket M \rrbracket_\rho) &= \max_{1 \leq i \leq n} ((m + 1) + \max(\llbracket M_i \rrbracket_\rho)) \\ &\leq \max_{1 \leq i \leq n} \max(\text{lh}(M_i) + m + 1, \max(\rho)) \\ &= \max(\text{lh}(M) + m, \max(\rho)) \end{aligned}$$

- If $M = x_0 M_1 \dots M_n$ and $\rho(x)$ is defined, we calculate:

$$\begin{aligned} \max(m + \llbracket M \rrbracket_\rho) &= \max\left(\left(\bigsqcup_{i=1}^n (m + 1) + \llbracket M_i \rrbracket_\rho\right) \cdot_{\text{lv}(x)+1} \rho(x)\right) \\ &\leq \max\left(\max_{1 \leq i \leq n} \max(m + 1 + \llbracket M_i \rrbracket_\rho), \max(\rho)\right) \\ &\leq \max\left(\max_{1 \leq i \leq n} \max(\text{lh}(M_i) + m + 1, \max(\rho)), \max(\rho)\right) \\ &= \max(\text{lh}(M) + m, \max(\rho)) \end{aligned}$$

- If $M = (\lambda x. M) M_1 \dots M_n$, then we calculate:

$$\begin{aligned} \max(m + \llbracket M \rrbracket_\rho) &= \max(m + \llbracket M M_2 \dots M_n \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}}}) \\ &\leq \max(\text{lh}(M M_2 \dots M_n) + m, \max(\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\})) \\ &= \max(\text{lh}(M M_2 \dots M_n) + m, \max(\rho), \max(\llbracket M_1 \rrbracket_\rho)) \\ &\leq \max(\text{lh}(M M_2 \dots M_n) + m, \max(\rho), \max(\text{lh}(M_1), \max(\rho))) \\ &\leq \max(\max(\text{lh}(M M_2 \dots M_n), \text{lh}(M_1)) + m, \max(\rho)) \\ &= \max(\text{lh}(M) + m, \max(\rho)) \end{aligned}$$

□

Lemma 32. If $\Gamma \vdash M : A$ is a term with a bs-environment ρ , we define $\text{ord}(\rho) = \max_{x \in \text{dom}(\rho)} \text{ord}(\rho(x))$. Then, we have:

$$\text{ord}(\llbracket M \rrbracket_\rho) \leq \max(\text{ord}(M), \text{ord}(\rho), \max_{x \in \text{dom}(\rho)} (\text{lv}(x) + 1))$$

In particular, $\text{ord}(\llbracket M \rrbracket) \leq \text{ord}(M)$.

Proof. By induction on the length of M , omitting some basic manipulations.

- If $M = x_0 M_1 \dots M_n$, $\rho(x)$ undefined, then $\llbracket M \rrbracket_\rho = 1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho$. We calculate:

$$\begin{aligned} \text{ord}(\llbracket M \rrbracket_\rho) &= \max_{1 \leq i \leq n} \text{ord}(\llbracket M_i \rrbracket_\rho) \\ &\leq \max_{1 \leq i \leq n} \max(\text{ord}(M_i), \text{ord}(\rho), \max_{x \in \text{dom}(\rho)} (\text{lv}(x) + 1)) \\ &= \max\left(\max_{1 \leq i \leq n} \text{ord}(M_i), \text{ord}(\rho), \max_{x \in \text{dom}(\rho)} (\text{lv}(x) + 1)\right) \\ &\leq \max(\text{ord}(M), \text{ord}(\rho), \max_{x \in \text{dom}(\rho)} (\text{lv}(x) + 1)) \end{aligned}$$

- If $M = x_0 M_1 \dots M_n$, $\rho(x)$ defined, then $\llbracket M \rrbracket_\rho = (1 + \bigsqcup_{i=1}^n \llbracket M_i \rrbracket_\rho) \cdot_{\text{lv}(x)+1} \rho(x)$.

$$\begin{aligned}
 \text{ord}(\llbracket M \rrbracket_\rho) &= \max(\max_{1 \leq i \leq n} \text{ord}(\llbracket M_i \rrbracket_\rho), \text{lv}(x) + 1, \text{ord}(\rho(x))) \\
 &\leq \max(\max_{1 \leq i \leq n} \max(\text{ord}(M_i), \text{ord}(\rho)), \max_{y \in \text{dom}(\rho)} (\text{lv}(y) + 1), \text{lv}(x) + 1, \text{ord}(\rho)) \\
 &= \max(\text{ord}(M), \text{ord}(\rho), \max_{y \in \text{dom}(\rho)} (\text{lv}(y) + 1))
 \end{aligned}$$

- If $M = \lambda x. M'$, then it is straightforward.
- If $M = (\lambda x^B. M') M_1 \dots M_n$, writing $\rho' = \rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}$:

$$\begin{aligned}
 \text{ord}(\llbracket M \rrbracket_\rho) &= \text{ord}(\llbracket M' M_2 \dots M_n \rrbracket_{\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}}) \\
 &\leq \max(\text{ord}(M' M_2 \dots M_n), \text{ord}(\rho \cup \{x \mapsto \llbracket M_1 \rrbracket_\rho\}), \max_{y \in \text{dom}(\rho')} (\text{lv}(y) + 1)) \\
 &= \max(\text{ord}(M' M_2 \dots M_n), \text{ord}(\llbracket M_1 \rrbracket_\rho), \text{ord}(\rho), \max_{y \in \text{dom}(\rho')} (\text{lv}(y) + 1)) \\
 &\leq \max(\text{ord}(M' M_2 \dots M_n), \text{ord}(M_1), \text{lv}(x) + 1, \text{ord}(\rho), \max_{y \in \text{dom}(\rho)} \text{lv}(y) + 1) \\
 &= \max(\text{ord}(M), \text{ord}(\rho), \max_{y \in \text{dom}(\rho)} \text{lv}(y) + 1)
 \end{aligned}$$

□

We can now summarize the results of this subsection with the following proposition.

Proposition 8. If M is a strongly locally scoped, η -long term of ground type, then:

$$\begin{array}{ll}
 \text{depth}(\llbracket M \rrbracket) \leq \text{depth}(M) & \max(\llbracket M \rrbracket) \leq \text{lh}(M) \\
 \text{ord}(\llbracket M \rrbracket) \leq \text{ord}(M) & \mathcal{N}(\llbracket M \rrbracket) \geq \mathcal{N}(M)
 \end{array}$$

Proof. For depth, local height and order, it is a consequence respectively of Lemmas 30, 31 and 32. For the norm, we prove it on locally scoped (not strongly) terms, by induction on $\mathcal{N}(\llbracket M \rrbracket)$. If $\mathcal{N}(\llbracket M \rrbracket) = 0$, then $\mathcal{N}(M) = 0$ as well by Proposition 7. If $\mathcal{N}(M) = 0$, this is obvious. Otherwise we have $M \rightarrow_{\text{lhr}} M'$. By Lemma 25, M' is still locally scoped. By Lemma 23, it is also still η -long. By Lemma 29, we have $M \rightarrow_{\emptyset} M'$. By Proposition 7, it follows that there is a skeleton a such that: $\llbracket M \rrbracket_{\emptyset} \rightarrow_{\text{bs}} a \leftrightarrow \llbracket M' \rrbracket_{\emptyset}$. By definition of norm and Lemma 2, we have $\mathcal{N}(\llbracket M \rrbracket_{\emptyset}) > \mathcal{N}(a) \geq \mathcal{N}(\llbracket M' \rrbracket_{\emptyset})$. But by IH, we know that $\mathcal{N}(M') \leq \mathcal{N}(\llbracket M' \rrbracket_{\emptyset})$. So, $\mathcal{N}(M) = \mathcal{N}(M') + 1 \leq \mathcal{N}(\llbracket M' \rrbracket_{\emptyset}) + 1 < \mathcal{N}(\llbracket M \rrbracket_{\emptyset}) + 1$. □

5.4. Bounds for strongly locally scoped terms. With Proposition 8 and Theorem 2 we can already deduce upper bounds for the length of lhr on sls η -long terms. However it will turn out that η -expansion does not change the asymptotic bounds, so we first deal with η -expansion and we will then formulate bounds for non necessarily η -long sls terms.

5.4.1. *How η -expansion affects the norm.*

Lemma 33. If M has a head occurrence x_0 of a variable x , and $M \rightarrow_{\eta} M'$, then for any term N , we have $M[N/x_0] \rightarrow_{\eta} M'[N/x_0]$.

Proof. Direct by induction on M . □

Lemma 34. If $(\lambda x, N)$ is a prime redex in $M N$, then for any N' , $(\lambda x, N')$ is a prime redex of $M N'$.

Proof. Direct by induction on M . □

Lemma 35. Suppose $M \rightarrow_\eta M'$, with $(\lambda x, N)$ prime redex in $M N$ and x_0 head occurrence of M , then there is a term M'' such that $M' N \rightarrow_{\text{lhr}}^+ M'' N$ and $M[N/x_0] N \rightarrow_\eta M'' N$.

Proof. By induction on the length of M . If the η -expansion is external, *i.e.* $M \rightarrow_\eta \lambda y. M y_0$, then by Lemma 34 we know that $(\lambda x, y_0)$ is a prime redex in $\lambda y. M y_0$. It follows that we have $(\lambda y. M y_0) N \rightarrow_{\text{lhr}} (\lambda y. M[y_1/x_0] y_0) N$. But then, y_1 becomes the head occurrence, and $(\lambda y, N)$ is a prime redex, therefore:

$$(\lambda y. M[y_1/x_0] y_0) N \rightarrow_{\text{lhr}} (\lambda y. M[N/x_0] y_0) N$$

and to conclude, we obviously have $M[N/x_0] N \rightarrow_\eta (\lambda y. M[N/x_0] y_0) N$.

If the η -expansion is internal, we reason by cases on the form of M .

- If $M = x_0 M_1 \dots M_n$, then M does not have any prime redex.
- If $M = \lambda x. M'$, then necessarily x_0 is an occurrence of x . Then we have $M' \rightarrow_\eta M''$, and the prime redex $(\lambda x, N)$ in $M N$ is still in $(\lambda x. M'') N$. Moreover, the head occurrence of M'' is still x_0 . Therefore, $(\lambda x. M'') N \rightarrow_{\text{lhr}} (\lambda x. M''[N/x_0]) N$. By Lemma 33, $M'[N/x_0] \rightarrow_\eta M''[N/x_0]$, so $(\lambda x. M'[N/x_0]) N \rightarrow_\eta (\lambda x. M''[N/x_0]) N$.
- If $M = (\lambda y. M') M_1 \dots M_n$, then by cases on the location of the η -expansion:
 - For $M \rightarrow_\eta (\lambda y. M'') M_1 \dots M_n$, it follows directly from IH.
 - For the remaining cases, the same lhr reduction as in M is possible and obviously commutes with the η -expansion. □

Lemma 36. If $M \rightarrow_\eta N$ and $M \rightarrow_{\text{lhr}} S$, then there is T such that $N \rightarrow_{\text{lhr}^+} T$ and $S \rightarrow_{\eta^+} T$.

Proof. By induction on M , detailing only non-trivial cases.

- If M has the form $\lambda x. M'$, then $M \rightarrow_{\text{lhr}} S$ with $S = \lambda x. S'$. Since we have $M \rightarrow_\eta N$, several cases arise:
 - If $N = \lambda y. M y_0$, it is immediate with $T = \lambda y. (\lambda x. S') y$.
 - If $M' \rightarrow_\eta N'$, IH provides T' with $N' \rightarrow_{\text{lhr}^+} T'$ and $S' \rightarrow_{\eta^+} T'$. Setting $T = \lambda x. T'$, the commutation follows.
- If M has the form $M_1 M_2$. Three cases arise:
 - If $M_1 M_2 \rightarrow_\eta \lambda x. M_1 M_2 x$, then the prime redexes of $\lambda x. M_1 M_2 x$ are included in those of $M_1 M_2$, and the head occurrence is the same. It follows that if $M_1 M_2 \rightarrow_{\text{lhr}} M'_1 M_2$, we have that $\lambda x. M_1 M_2 x \rightarrow_{\text{lhr}} \lambda x. M'_1 M_2 x$ as well. Setting $T = \lambda x. M'_1 M_2 x$ yields the required diagram.
 - If $M_1 M_2 \rightarrow_\eta M'_1 M_2$ (so $M_1 \rightarrow_\eta M'_1$). Then, two cases arise.
 - * If the reduction $M_1 M_2 \rightarrow_{\text{lhr}} S_1 M_2$ involves a prime redex $(\lambda y, M_2)$, and the head occurrence of M_1 is y_0 . Then by Lemma 35, there is a term T' such that $M'_1 M_2 \rightarrow_{\text{lhr}}^+ T' M_2$ and $M_1[M_2/y_0] M_2 \rightarrow_\eta T' M_2$. Setting $T = T' M_2$, we have the required commutation.
 - * If the reduction $M_1 M_2 \rightarrow_{\text{lhr}} S_1 M_2$ does not involve M_2 , then we also have $M_1 \rightarrow_\eta M'_1$, and $M_1 \rightarrow_{\text{lhr}} S_1$. By IH, there is T_1 such that $M'_1 \rightarrow_{\text{lhr}^+} T_1$ and $S_1 \rightarrow_{\eta^+} T_1$. Setting $T = T_1 M_2$ gives the required commutation.
 - If $M_1 M_2 \rightarrow_\eta M_1 M'_2$, then two sub-cases arise:

- * If $M_1 M_2 \rightarrow_{\text{lhr}} S_1 M_2$ is obtained by firing a prime redex $(\lambda x, M_2)$ with x_0 head occurrence of M_1 , then setting $T = M_1[M'_2/x_0]$ yields $M_1 M'_2 \rightarrow_{\text{lhr}} T$ and $S_1 M_2 = M_1[M_2/x_0] M_2 \rightarrow_{\eta^2} M_1[M'_2/x_0] M'_2$.
- * If $M_1 M_2 \rightarrow_{\text{lhr}} S_1 M_2$ is obtained by firing a prime redex not involving M_2 , then setting $T = S_1 M'_2$ immediately yields the required commutation. \square

Lemma 37. If $M \rightarrow_{\eta^n} N$ and $M \rightarrow_{\text{lhr}^p} S$, then there are T , $n' \geq n$ and $p' \geq p$ such that $N \rightarrow_{\text{lhr}^{p'}} T$ and $S \rightarrow_{\eta^{n'}} T$.

Proof. We reason by induction on the lexicographic ordering of pairs $(\mathcal{N}(N), n)$, using that $\mathcal{N}(N)$ is finite by Proposition 5. If $n = 0$ or $p = 0$, it is obvious. Otherwise, we have $M \rightarrow_{\eta} N_1 \rightarrow_{\eta^{n-1}} N$ and $M \rightarrow_{\text{lhr}} S_1 \rightarrow_{\text{lhr}^{p-1}} S$. By Lemma 36, there is N'_1 such that $M_1 \rightarrow_{\eta^+} N'_1$ and $N_1 \rightarrow_{\text{lhr}^+} N'_1$, let us write $M_1 \rightarrow_{\eta^q} N'_1$ and $N_1 \rightarrow_{\text{lhr}^r} N'_1$, with $r, q \geq 1$. We represent the situation on the following diagram:

$$\begin{array}{ccccc}
 M & \xrightarrow{\eta} & N_1 & \xrightarrow{\eta^{n-1}} & N \\
 \text{lhr} \downarrow & & \downarrow \text{lhr}^r & & \\
 S_1 & \xrightarrow{\eta^q} & N'_1 & & \\
 \text{lhr}^{p-1} \downarrow & & & & \\
 S & & & &
 \end{array}$$

We have $n - 1 < n$ and $\mathcal{N}(N)$ is unchanged, therefore by IH, there is N' as displayed here:

$$\begin{array}{ccccc}
 M & \xrightarrow{\eta} & N_1 & \xrightarrow{\eta^{n-1}} & N \\
 \text{lhr} \downarrow & & \downarrow \text{lhr}^r & & \downarrow \text{lhr}^{r'} \\
 S_1 & \xrightarrow{\eta^q} & N'_1 & \xrightarrow{\eta^{n'}} & N' \\
 \text{lhr}^{p-1} \downarrow & & & & \\
 S & & & &
 \end{array}$$

with $r' \geq r$ and $n' \geq n - 1$. Finally, $\mathcal{N}(N') < \mathcal{N}(N)$, therefore by IH there is T closing the diagram:

$$\begin{array}{ccccc}
 M & \xrightarrow{\eta} & N_1 & \xrightarrow{\eta^{n-1}} & N \\
 \text{lhr} \downarrow & & \downarrow \text{lhr}^r & & \downarrow \text{lhr}^{r'} \\
 S_1 & \xrightarrow{\eta^q} & N'_1 & \xrightarrow{\eta^{n'}} & N' \\
 \text{lhr}^{p-1} \downarrow & & & & \downarrow \text{lhr}^{p'} \\
 S & \xrightarrow{\eta^{n''}} & & & T
 \end{array}$$

with $p' \geq p - 1$ hence $p' + r' \geq p - 1 + r \geq p - 1 + 1 \geq p$, and $n'' \geq q + n' \geq 1 + n - 1 \geq n$. \square

Proposition 9. If $M \rightarrow_{\eta} N$, then $\mathcal{N}(N) \geq \mathcal{N}(M)$.

Proof. We show by induction on $\mathcal{N}(M)$ that if $M \rightarrow_{\eta}^* N$, then $\mathcal{N}(N) \geq \mathcal{N}(M)$. If $\mathcal{N}(M) = 0$, this is obvious. Otherwise, there is $M \rightarrow_{\text{lhr}} M'$. By Lemma 37 there is N' such that

$N \rightarrow_{\text{lh}^+} N'$ and $M' \rightarrow_{\eta}^* N'$. But $\mathcal{N}(M') < \mathcal{N}(M)$, so by IH we have $\mathcal{N}(N') \geq \mathcal{N}(M')$. Since $\mathcal{N}(M) = \mathcal{N}(M') + 1$ and $\mathcal{N}(N) \geq \mathcal{N}(N') + 1$, we have $\mathcal{N}(N) \geq \mathcal{N}(M)$ as well. \square

5.4.2. How η -expansion affects other quantities on terms.

Definition 20. An η -expansion step $M \rightarrow_{\eta} M'$ is **restricted** if M and M' have the same number of generalized redexes. We write $M \rightarrow_{\eta_r} M'$.

Lemma 38. On any term M , the expansion \rightarrow_{η_r} terminates.

Proof. Suppose A is a type, then its **size** written $\text{size}(A)$, is defined by induction on A by $\text{size}(o) = 1$ and $\text{size}(A \rightarrow B) = \text{size}(A) + \text{size}(B)$. If M is a term, its **deficiency** quantifies the lack of η -expansion within M . It is defined by induction on the length of M , as follows:

- If $M = x_0 M_1 \dots M_n$, where x_0 is an occurrence of a variable or constant, and if $x : A_1 \rightarrow \dots \rightarrow A_p \rightarrow o$, then set $\text{df}(M) = \sum_{i=1}^n \text{df}(M_i) + \sum_{i=n+1}^p \text{size}(A_i)$.
- If $M = \lambda x. M'$, we set $\text{df}(M) = \text{df}(M')$.
- If $M = (\lambda x. M') M_1 \dots M_n$, we set $\text{df}(M) = \text{df}(M' M_2 \dots M_n) + \text{df}(M_1)$.

Restricted η -expansion strictly decreases deficiency, by induction on the length of M :

- If $M = x_0 M_1 \dots M_n$ where x_0 is a variable or constant occurrence having type $A_1 \rightarrow \dots \rightarrow A_p \rightarrow o$, then by definition we have $\text{df}(M) = \sum_{i=1}^n \text{df}(M_i) + \sum_{i=n+1}^p \text{size}(A_i)$. We reason by cases on the location of the η -expansion. If it is:

$$x_0 M_1 \dots M_n \rightarrow_{\eta_r} \lambda y. x_0 M_1 \dots M_n y_0$$

then $\text{df}(\lambda y. x_0 M_1 \dots M_n y_0) = \sum_{i=1}^n \text{df}(M_i) + \text{df}(y_0) + \sum_{i=n+2}^p \text{size}(A_i)$. But it is obvious by definition that $\text{df}(y_0) = \text{size}(A_{n+1}) - 1$, so deficiency is reduced. If the η -expansion is $x_0 M_1 \dots M_n \rightarrow_{\eta_r} x_0 M_1 \dots M'_i \dots M_n$, then $M_i \rightarrow_{\eta_r} M'_i$, so by IH we have $\text{df}(M'_i) < \text{df}(M_i)$. So, $\text{df}(x_0 M_1 \dots M'_i \dots M_n) < \text{df}(x_0 M_1 \dots M_n)$.

- If $M = \lambda x. M'$, the η -expansion cannot be $\lambda x. M' \rightarrow \lambda y. (\lambda x. M') y_0$ as that would create a new generalized redex, so it is $M' \rightarrow_{\eta_r} M''$. The property follows from IH.
- If $M = (\lambda x. M') M_1 \dots M_n$, then the only possible restricted η -expansions are:

$$\begin{aligned} (\lambda x. M') M_1 \dots M_n &\rightarrow_{\eta_r} \lambda y. (\lambda x. M') M_1 \dots M_n y_0 \\ (\lambda x. M') M_1 \dots M_n &\rightarrow_{\eta_r} (\lambda x. M'') M_1 \dots M_n \\ (\lambda x. M') M_1 \dots M_n &\rightarrow_{\eta_r} (\lambda x. M') M_1 \dots M'_i \dots M_n \end{aligned}$$

All the others create a generalized β -redex. In all three cases, the result follows from IH and definition of deficiency.

We have a positive strictly decreasing measure for η_r , so it terminates. \square

Lemma 39. If M is a term and $M \rightarrow_{\eta_r}^* M'$, then $\text{lh}(M') \leq \text{lh}(M) + \text{ord}(M)$.

Proof. We define a quantity $\text{lh}'(M)$ such that $\text{lh}(M) \leq \text{lh}'(M) \leq \text{lh}(M) + \text{ord}(M)$ and we show that lh' is preserved by restricted η -expansion. It is defined as for lh , except for variables where we set, with x_0 occurrence of $x : A_1 \rightarrow \dots \rightarrow A_p \rightarrow o$:

$$\text{lh}'(x_0 M_1 \dots M_n) = 1 + \max\left(\max_{1 \leq i \leq n} \text{lh}'(M_i), \max_{1 \leq i \leq n} \text{lv}(A_i)\right)$$

Clearly, we have $\text{lh}(M) \leq \text{lh}'(M) \leq \text{lh}(M) + \text{ord}(M)$. We prove by induction on M that it is preserved by restricted η -expansion, skipping the trivial case.

- If $M = x_0 M_1 \dots M_n$ where x_0 is a variable occurrence of a variable $x : A_1 \rightarrow \dots \rightarrow A_p \rightarrow o$, then several cases depending on the location of the η -expansion. Firstly, if the η -expansion is: $x_0 M_1 \dots M_n \rightarrow_{\eta_r} \lambda y. x_0 M_1 \dots M_n y_0$ we have

$$\begin{aligned} \text{lh}'(x_0 M_1 \dots M_n) &= 1 + \max(\max_{1 \leq i \leq n} \text{lh}'(M_i), \max_{n+1 \leq i \leq p} \text{lv}(A_i)) \\ \text{lh}'(\lambda y. x_0 M_1 \dots M_n y) &= 1 + \max(\max_{1 \leq i \leq n} \text{lh}'(M_i), \text{lh}'(y), \max_{n+2 \leq i \leq p} \text{lv}(A_i)) \end{aligned}$$

But then, writing $y : B_1 \rightarrow \dots \rightarrow B_m \rightarrow o$, we have $\text{lh}'(y) = 1 + \max_{1 \leq i \leq m} \text{lv}(B_i) = \text{lv}(A_{n+1})$, so those two quantities are equal. Secondly, if the restricted η -expansion is within some M_i , then the result follows immediately by IH.

- If $M = (\lambda x. M') M_1 \dots M_n$, then several cases following the location of the η -expansion. For $(\lambda x. M') M_1 \dots M_n \rightarrow_{\eta_r} \lambda y. (\lambda x. M') M_1 \dots M_n y_0$, we calculate:

$$\begin{aligned} \text{lh}'((\lambda x. M') M_1 \dots M_n) &= \max(\text{lh}'(M' M_2 \dots M_n), \text{lh}'(M_1)) \\ &= \max(\text{lh}'(\lambda y. M' M_2 \dots M_n y_0), \text{lh}'(M_1)) \\ &= \max(\text{lh}'(M' M_2 \dots M_n y_0), \text{lh}'(M_1)) \\ &= \text{lh}'((\lambda x. M') M_1 M_2 \dots M_n y_0) \\ &= \text{lh}'(\lambda y. (\lambda x. M') M_1 M_2 \dots M_n y_0) \end{aligned}$$

If the η -expansion is within M' or M_i , it follows directly from IH. □

Lemma 40. If $M \rightarrow_{\eta_r} M'$, then $\text{depth}(M) = \text{depth}(M')$.

Proof. Immediate by induction on M . □

Lemma 41. If $M \rightarrow_{\eta} M'$, then $\text{ord}(M) = \text{ord}(M')$

Proof. There is a subterm N such that $N \rightarrow_{\eta} \lambda y. N y_0$. But $\text{lv}(\lambda y. N y_0) = \text{lv}(N)$ and $\text{lv}(N y_0) \leq \text{lv}(N)$, so the new subterms have lower level than the original ones. □

Finally, it remains to note that η -expansion preserves strong local scope.

Lemma 42. If M is strongly locally scoped and $M \rightarrow_{\eta_r} M'$, then M' is sls.

Proof. B straightforward induction on the length of M . □

Proposition 10. If M is a term, then there is an η -long term M' such that:

$$\begin{array}{ll} \text{lh}(M') \leq \text{lh}(M) + \text{ord}(M) & \text{depth}(M') = \text{depth}(M) \\ \text{ord}(M') = \text{ord}(M) & \mathcal{N}(M') \geq \mathcal{N}(M) \end{array}$$

Moreover if M was strongly locally scoped, so is M' .

Proof. By Lemma 38, there is M' such that $M \rightarrow_{\eta_r}^* M'$, and there is no further restricted η -expansion. By definition, M' is η -long. Moreover, the preservations of depth, order, local height and norm follow respectively from Lemmas 40, 41, 39 and Proposition 9. The construction preserves strong local scope by Lemma 42. □

5.4.3. *Bounds for strongly locally scoped terms.* Putting everything together, we estimate:

$$\text{Lls}_n(h, d) = \max\{\mathcal{N}(M) \mid \text{ord}(M) \leq n \ \& \ \text{lh}(M) \leq h \ \& \ \text{depth}(M) \leq d \ \& \ M \text{ sls}\}$$

Proposition 11. Suppose M is a sls term of order at least one. Then,

$$\mathcal{N}(M) \leq 2^{\text{depth}(M) \log(\text{lh}(M) + \text{ord}(M) + 1)}_{\text{ord}(M) - 1}$$

Proof. If $\Gamma \vdash M : A_1 \rightarrow \dots \rightarrow A_n \rightarrow o$ is a sls term, we first make it of ground type by forming $\Gamma \vdash M *_{A_1} \dots *_{A_n} : o$ – its norm can only increase, the other quantities stay unchanged and the term is still sls. By Proposition 10, there is M' η -long, of ground type, and sls such that $\text{lh}(M') \leq \text{lh}(M) + \text{ord}(M)$, $\text{depth}(M') = \text{depth}(M)$, $\text{ord}(M') = \text{ord}(M)$ and $\mathcal{N}(M') \geq \mathcal{N}(M)$. We conclude by Proposition 8 and Theorem 2. \square

We now prove the optimality of this upper bound by exhibiting a family of terms whose reduction length asymptotically reaches it. This family of terms is closely related to the example used in Section 3 for game situations. For $n, k, p \geq 0$ and $M : A_p$, we define:

$$[n]_p^0(M) = M \qquad [n]_p^{k+1}(M) = \underline{n}_{p+1} [n]_p^k(M)$$

One can immediately check that $[n]_p^k(M) : A_p$ and that for all $q \in \mathbb{N}$, $[n]_p^k(q_p) \rightarrow_{\beta}^* \underline{q}_{p-k}$. Exploiting this construction we set, for $n, k, p \geq 0$:

$$S_{n,k,p} = [n]_p^k(\underline{2}_p) \underline{2}_{p-1} \dots \underline{2}_0$$

For which it is immediate to check that for all $n, k, p \geq 0$ we have $S_{n,k,p} \rightarrow_{\beta}^* \underline{2}_{p-2n}^{2^{n+k}}$. Moreover, by construction of $S_{n,k,p}$, for $n \geq 2$ and $p, k \geq 1$ we have $\text{lh}(S_{n,k,p}) = n + 1$, $\text{depth}(S_{n,k,p}) = k + 1$ and $\text{ord}(S_{n,k,p}) = p + 3$, and $S_{n,k,p}$ is sls. To deduce a lower bound from this, we use:

Lemma 43. If $M \rightarrow_{\beta}^* \underline{n}_0$, then $\mathcal{N}(M \text{ id}_o) \geq n$, where $\text{id}_o = \lambda x^o. x$.

Proof. By induction on n , exploiting that lhr preserves β -equivalence. \square

Theorem 4. For fixed $n \geq 2$ we have $\text{Lls}_n(h, d) = 2^{\Theta(d \log(h))}_{n-1}$.

Proof. We start with $n \geq 3$, $n = 2$ requires a separate construction for the lower bound. Let us fix $h \geq 3$ and $d \geq 2$. By Proposition 11, we already know that $\text{Lls}_n(d, h) \leq 2^{\log(h+n+1)}_{n-1}$. Moreover, we have $\text{lh}(S_{h-1, d-1, n-3} \text{ id}_o) = h$ and $\text{depth}(T_{h-1, d-1, n-3} \text{ id}_o) = d$, and by Lemma 43 we have $\mathcal{N}(T_{h-1, d-1, n-3} \text{ id}_o) \geq 2^{(d-1) \log(h-1)}_{n-1}$. To summarize:

$$2^{(d-1) \log(h-1)}_{n-1} \leq \text{Lls}_n(d, h) \leq 2^{\log(h+n+1)}_{n-1}$$

Therefore, with $n \geq 3$ fixed and d, h parameters we have $\text{Lls}_n(h, d) = 2^{\Theta(d \log(h))}_{n-1}$.

For $n = 2$, the upper bound still holds. For $d, p \geq 2$, define:

$$U_{n,d} = \underline{n}_1 (\underline{n}_1 \dots (\underline{n}_1 \text{ id}_o) \dots)$$

with d copies of \underline{n}_1 in total. Then, $U_{n,d}$ is sls and $\text{lh}(U_{n,d}) = n + 1$, $\text{depth}(U_{n,d} \text{ id}_o) = d + 1$, $\text{ord}(U_{n,d} \text{ id}_o) = 2$ and $\mathcal{N}(U_{n,d}) \geq n^d = 2^{d \log(n)}$. It follows that $\text{Lls}_2(d, h) = 2^{\Theta(d \log(h))}$. \square

$$\begin{array}{c}
 \frac{y \in \text{fv}(M_1)}{(\lambda x. M) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda x. M[x \ y/x]) (\lambda y'. M_1[y'/y]) \dots M_n} \\
 \\
 \frac{M_i \rightarrow_{\lambda l} M'_i}{x_0 M_1 \dots M_n \rightarrow_{\lambda l} x_0 M_1 \dots M'_i \dots M_n} \qquad \frac{M \rightarrow_{\lambda l} M'}{\lambda x. M \rightarrow_{\lambda l} \lambda x. M'} \\
 \\
 \frac{M_1 \rightarrow_{\lambda l} M'_1}{(\lambda x. M) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda x. M) M'_1 \dots M_n} \\
 \\
 \frac{M M_2 \dots M_n \rightarrow_{\lambda l} M' M'_2 \dots M'_n \quad (M_1 \text{ closed})}{(\lambda x. M) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda x. M') M_1 M'_2 \dots M'_n}
 \end{array}$$

 Figure 4: Definition of the λ -lifting expansion $\rightarrow_{\lambda l}$

In particular, reduction length for sls second-order terms of fixed depth is bounded by a polynomial of degree less than the depth.

5.5. Generalization to arbitrary terms. In this final subsection, we deduce from the study of lhr of ls terms a bound on the length of lhr of arbitrary terms. The key observation is that any λ -term can be transformed into a locally scoped form through λ -lifting [16].

5.5.1. Lambda-lifting to sls terms. Take a term $M = \lambda x^A. (\lambda y^A. y) x$. Obviously, M is not sls: indeed there is a prime redex $(\lambda y, x)$ and the subterm x has x free. In order to make the variable x “local”, we modify the abstraction subterm $\lambda y. y$ to forward explicitly the variable x . We get the term $M' = \lambda x^A. (\lambda y^{A \rightarrow A}. y x)(\lambda x'^A. x')$. The type of y has changed, but not the type of the overall term. Note that the terms M and M' are still β -equivalent, although we are not going to use that explicitly. More importantly, the norm has increased, the order has increased by one, and the other quantities are essentially unchanged. We formalize this construction by the λ -lifting expansion $\rightarrow_{\lambda l}$, defined in Figure 4.

First, we prove that $\rightarrow_{\lambda l}$ indeed allows us to convert any term M into a sls M' . This is done by showing that $\rightarrow_{\lambda l}$ terminates, and that its normal forms are sls.

Lemma 44. Let M be any term. Then, $\rightarrow_{\lambda l}$ terminates on M .

Proof. For each occurrence x_0 of x in M , we define its **binding distance** $d_M(x_0)$:

$$\begin{array}{ll}
 d_{x_0 M_1 \dots M_n}(x_0) & = 0 \\
 d_{* M_1 \dots M_n}(x_0) & = d_{M_i}(x_0) \qquad (x_0 \in M_i) \\
 d_{y_0 M_1 \dots M_n}(x_0) & = d_{M_i}(x_0) \qquad (x_0 \in M_i) \\
 d_{\lambda x. M'}(x_0) & = d_{M'}(x_0) \\
 d_{(\lambda y. M') M_1 \dots M_n}(x_0) & = d_{M'}(x_0) \qquad (x_0 \in M') \\
 d_{(\lambda y. M') M_1 \dots M_n}(x_0) & = \max_{y_i \in \text{oc}_y(M')} d_{M'}(y_i) + d_{M_1}(x_0) + 1 \quad (x_0 \in M_1 \wedge x \in \text{fv}(M')) \\
 d_{(\lambda y. M') M_1 \dots M_n}(x_0) & = d_{M_1}(x_0) \qquad (x_0 \in M_1 \wedge x_0 \notin \text{fv}(M_1)) \\
 d_{(\lambda y. M') M_1 \dots M_n}(x_0) & = d_{M' M_2 \dots M_n}(x_0) \qquad (x_0 \in M_i \wedge i \geq 2)
 \end{array}$$

where $x_0 \in M$ means that x_0 is a variable occurrence x in M and $\text{oc}_y(M)$ is the set of occurrences of a variable y in M . Then, for any term M we consider the multiset $\mathcal{M}(M) =$

$\{d_M(x_0) \mid x_0 \in oc_x(M)\}$ where by $\{- \mid -\}$ we denote here multiset comprehension. Then, we show by induction on $M \rightarrow_{\lambda l} M'$ that $\mathcal{M}(M') < \mathcal{M}(M)$ for $<$ the multiset well-founded ordering on multisets of natural numbers.

- If $M = (\lambda x. M') M_1 \dots M_n \rightarrow_{\lambda l} (\lambda x. M'[x y/x]) (\lambda y'. M_1[y'/y]) \dots M_n = M''$:
 - Occurrences of variables other than y' in $M_1[y'/y]$ keep their binding distance,
 - Occurrences of y in M_1 of distance $d_M(y_i) = \max_{x_i \in oc_x(M')} d_{M'}(x_i) + d_{M_i}(y_i) + 1$ have been replaced by occurrences of y' of distance $d_{M''}(y'_i) = d_{M_i}(y_i)$.
 - New occurrences of y of distance less than $\max_{x_i \in oc_x(M')} d_{M'}(x_i)$ appear.
 So, each variable occurrence is either unchanged or replaced by occurrences of strictly smaller distance, so the reduction decreases the multiset ordering on $\mathcal{M}(M)$.
- The other cases follow from IH since $\mathcal{M}(-)$ is additive over all term constructors. \square

Lemma 45. If a term M is λl -normal, M is strongly locally scoped.

Proof. Straightforward induction on the length of M . \square

5.5.2. *How λ -lifting increases the norm.* Now, it remains to investigate how λ -lifting affects the relevant quantities on terms. First, we show that it can only increase the norm.

Lemma 46. If M has a head occurrence x_0 where x is free in M and N is a term, then if $M \rightarrow_{\lambda l} M'$, then the head occurrence of M' is still x_0 and $M[N/x_0] \rightarrow_{\lambda l} M'[N/x_0]$.

Proof. By induction on the length of M . The only non-trivial case is for the term $M = (\lambda y. M') M_1 \dots M_n$, where three subcases arise:

- If $M' \rightarrow_{\lambda l} M''$, then by IH we know that $M'[N/x_0] \rightarrow_{\lambda l} M''[N/x_0]$, therefore $(\lambda y. M'[N/x_0]) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda y. M''[N/x_0]) M_1 \dots M_n$.
- If $M_i \rightarrow_{\lambda l} M'_i$, then $(\lambda y. M'[N/x_0]) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda y. M'_i[N/x_0]) M_1 \dots M_n$.
- If $(\lambda y. M') M_1 \dots M_n \rightarrow_{\lambda l} (\lambda y. M'[y z/y]) (\lambda z. M_1) \dots M_n$, then since x is free in M' , necessarily $y \neq x$ (y is bounded). Hence $M'[y z/y][N/x_0] = M'[N/x_0][y z/y]$, so $(\lambda y. M'[N/x_0]) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda y. M'[y z/y][N/x_0]) (\lambda z. M_1) \dots M_n$. \square

Lemma 47. If M has a head occurrence x_0 where x is free in M and N is a term with $N \rightarrow_{\lambda l} N'$, then we have $M[N/x_0] \rightarrow_{\lambda l} M[N'/x_0]$.

Proof. Straightforward by induction on the length of M . \square

Lemma 48. If $M \rightarrow_{\lambda l} M'$ and $M \rightarrow_{\text{hr}} N$, then there exists S, T such that:

$$\begin{array}{ccc}
 M & \xrightarrow{\lambda l} & M' \\
 \downarrow \text{hr} & & \searrow \text{hr} \\
 N & \xrightarrow{\lambda l^+} & S & \xleftarrow{\beta_{si}^\bullet} & T
 \end{array}$$

Where in β_{si}^\bullet , by s we mean the β -reduction of a *spine redex*, i.e. a β -redex that is also a prime redex, by i we mean that it is *internal*, in the sense that it is not the leftmost redex, and by \bullet we mean that there are either zero or one reduction steps.

Proof. By induction on M , detailing only the non-trivial cases.

- If $M = \lambda x. M'$, it follows directly from IH.

- If $M = (\lambda x. M') M_1 \dots M_n$ and the head occurrence of M' is an occurrence x_0 of x , then $(\lambda x. M') M_1 \dots M_n \rightarrow_{\text{Ihr}} (\lambda x. M'[M_1/x_0]) M_1 \dots M_n$. There are several cases. If the $\rightarrow_{\lambda 1}$ step comes from $M' M_2 \dots M_n \rightarrow_{\lambda 1} M'' M'_2 \dots M'_n$, then by Lemma 46 we also have:

$$M'[M_1/x_0] M_2 \dots M_n \rightarrow_{\lambda 1} M''[M_1/x_0] M'_2 \dots M'_n$$

It immediately follows:

$$(\lambda x. M'[M_1/x_0]) M_1 \dots M_n \rightarrow_{\lambda 1} (\lambda x. M''[M_1/x_0]) M_1 M'_2 \dots M'_n$$

which provides the necessary commutation. If the $\rightarrow_{\lambda 1}$ comes from $M_1 \rightarrow_{\lambda 1} M'_1$, then by Lemma 47 we also have $M'[M_1/x_0] \rightarrow_{\lambda 1} M'[M'_1/x_0]$. It immediately follows:

$$(\lambda x. M'[M_1/x_0]) M_1 \dots M_n \rightarrow_{\lambda 1^2} (\lambda x. M'[M'_1/x_0]) M'_1 \dots M_n$$

which provides the necessary commutation. Finally, if the $\rightarrow_{\lambda 1}$ step is:

$$(\lambda x. M') M_1 \dots M_n \rightarrow_{\lambda 1} (\lambda x. M'[x z/x]) (\lambda z. M_1) \dots M_n$$

Then the head occurrence of $(\lambda x. M'[x z/x]) (\lambda z. M_1) \dots M_n$ is still x_0 . Moreover,

$$\begin{aligned} (\lambda x. M'[x z/x]) (\lambda z. M_1) \dots M_n &\rightarrow_{\text{Ihr}} (\lambda x. M'[x z/x][(\lambda z. M_1)/x_0]) (\lambda z. M_1) \dots M_n \\ &= (\lambda x. M'[((\lambda z. M_1) z)/x_0][x z/x]) (\lambda z. M_1) \dots M_n \\ &\rightarrow_{\beta_{si}} (\lambda x. M'[M_1/x_0][x z/x]) (\lambda z. M_1) \dots M_n \\ &\leftarrow_{\lambda 1} (\lambda x. M'[M_1/x_0]) M_1 \dots M_n \end{aligned}$$

- If $M = (\lambda y. M') M_1 \dots M_n$ whose head occurrence x_0 is not an occurrence of y , then three cases.
 - If the $\rightarrow_{\lambda 1}$ -reduction comes from $M' M_2 \dots M_n \rightarrow_{\lambda 1} M'' M'_2 \dots M'_n$, it follows directly from IH.
 - If the $\rightarrow_{\lambda 1}$ -reduction comes from $M_1 \rightarrow_{\lambda 1} M'_1$, then it is obvious.
 - If the $\rightarrow_{\lambda 1}$ -reduction is:

$$(\lambda y. M') M_1 \dots M_n \rightarrow_{\lambda 1} (\lambda y. M'[y z/y]) (\lambda z. M_1) \dots M_n$$

The head occurrence of M' is not an occurrence of y , so the lhr reduct of $(\lambda y. M') M_1 \dots M_n$ must have the form $(\lambda y. M'[N/x_0]) M_1 \dots M_n$ for some N . The head occurrence of $(\lambda y. M'[y z/y]) (\lambda z. M_1) \dots M_n$ is still x_0 , and:

- * If the associated prime redex $(\lambda x, N)$ has not changed, the same lhr is possible and commutes with the $\lambda 1$ -reduction.
- * If the associated prime redex has become $(\lambda x, N[y z/y])$, we have $M'[N/x_0][y z/y] = M'[N[y z/y]/x_0][y z/y]$, so the same reasoning as above applies. \square

Proposition 12. If $M \rightarrow_{\lambda 1} M'$, then $\mathcal{N}(M') \geq \mathcal{N}(M)$.

Proof. By induction on $\mathcal{N}(M')$. By Lemma 48, there are terms S, T such that:

$$\begin{array}{ccc} M & \xrightarrow{\lambda 1} & M' \\ \downarrow \text{Ihr} & & \searrow \text{Ihr} \\ N & \xrightarrow{\lambda 1^+} & S \end{array} \quad \begin{array}{c} \\ \\ \nearrow \beta^* \\ T \end{array}$$

By Proposition 6, we have $\mathcal{N}(S) \leq \mathcal{N}(T) = \mathcal{N}(M') - 1$. Moreover, we have a chain:

$$N = N_1 \rightarrow_{\lambda 1} N_2 \rightarrow_{\lambda 1} \cdots \rightarrow_{\lambda 1} N_n = S$$

But since $\mathcal{N}(S) < \mathcal{N}(M')$, it follows by immediate induction that $\mathcal{N}(N) \leq \mathcal{N}(S)$. Therefore, we have $\mathcal{N}(M) = \mathcal{N}(N) + 1 \leq \mathcal{N}(S) \leq \mathcal{N}(M') - 1 < \mathcal{N}(M')$. \square

5.5.3. *How λ -lifting preserves other quantities on terms.* Finally, We examine how λ -lifting affects the other quantities on terms. We first notice that it preserves the depth.

Lemma 49. If $M \rightarrow_{\lambda 1} M'$, then $\text{depth}(M) = \text{depth}(M')$.

Proof. First, we prove by induction on M that for any x free in M and variable y , we have $\text{depth}(M) = \text{depth}(M[x y/x])$. The lemma follows by induction on the definition of $\rightarrow_{\lambda 1}$. \square

Unfortunately it does affect the order and the local height; however we will show that multiple applications of $\rightarrow_{\lambda 1}$ can only change them by one. To prove that, we will construct *weighted* variants of order and local height that give different weight to variables according to their behaviour with respect to $\rightarrow_{\lambda 1}$. By design they will be preserved by $\rightarrow_{\lambda 1}$, but will remain within small bounds of the original level and local height. We start with the order.

Say that a variable x in a term M is **local** in M iff for any generalized redex $(\lambda y, N)$ of M , x does not appear free in N . If x is a variable of M define the **weighted level** $\text{lv}'(x)$ by $\text{lv}(x) + 1$ if x is local and $\text{lv}(x) + 2$ otherwise. Then, define the **weighted order** $\text{ord}'(M)$ as being the maximum over all $\text{lv}'(x)$ for variables x in M , and all $\text{lv}(A)$ for constants $*_A$ in M . First, we prove that $\text{ord}'(M)$ remains closely related to $\text{ord}(M)$.

Lemma 50. Let M be a closed term, then $\text{ord}(M) \leq \text{ord}'(M) \leq \text{ord}(M) + 1$.

Proof. We prove first that $\text{ord}(M) \leq \text{ord}'(M)$. If M admits as a subterm a constant $*_A$ with $\text{ord}(M) = \text{lv}(A)$, then $\text{ord}'(M) \geq \text{lv}(A) \geq \text{ord}(M)$. Otherwise, M has a subterm λx^A . M' such that $\text{ord}(M) = \text{lv}(A) + 1$ – indeed if a subterm of M of maximal level has the form $M_1 M_2$ then M_1 has higher level, and if it has the form λx^A . M' with $\text{lv}(M') > \text{lv}(A) + 1$ then M' still has maximal level so the property follows by induction. So in particular M has a variable x with $\text{ord}(M) = \text{lv}(x) + 1$, which is always less than $\text{ord}'(M)$.

We now prove that $\text{ord}'(M) \leq \text{ord}(M) + 1$. Firstly, if there is a constant $*_A$ in M such that $\text{ord}'(M) = \text{lv}(A)$, then $\text{ord}(M) + 1 \geq \text{ord}(M) \geq \text{lv}(A)$ as well. Secondly, if there is a local variable x^A such that $\text{ord}'(M) = \text{lv}(A) + 1$, then since M is closed it has a subterm of the form λx^A . M' , so $\text{ord}(M) + 1 \geq \text{lv}(\lambda x^A$. $M') + 1 \geq \text{lv}(A) + 2 > \text{lv}(A) + 1$. Finally if there is a non-local variable x^A such that $\text{ord}'(M) = \text{lv}(A) + 2$, same reasoning. \square

Now we prove that the weighted order is preserved by $\rightarrow_{\lambda 1}$.

Lemma 51. For any term M with $M \rightarrow_{\lambda 1} M'$, then $\text{ord}'(M) = \text{ord}'(M')$.

Therefore (by Lemma 50) if M is closed with $M \rightarrow_{\lambda 1}^* N$, then $\text{ord}(N) \leq \text{ord}(M) + 1$.

Proof. By induction on the definition of $\rightarrow_{\lambda 1}$.

- If $M_l = (\lambda x. M') M_1 \dots M_n \rightarrow_{\lambda 1} (\lambda x. M'[x y/x]) (\lambda y'. M_1[y'/y]) \dots M_n = M_r$, both terms contain the same constants. If z is a variable on the left hand side, reason by cases. For clarity we write the levels of variables as $\text{lv}_l(x)$ or $\text{lv}'_l(x)$ on the left hand side, and $\text{lv}_r(x)$ or $\text{lv}'_r(x)$ on the right hand side. We describe a mapping of variables of M_l to variables of M_r increasing the weighted level.

- We map x to x if $lv_l(x) \geq lv_l(y) + 1$. Then x is local in M_l iff it is local in M_r , so there is $i \in \{1, 2\}$ such that we have $lv'_l(x) = lv_l(x) + i$ and $lv'_r(x) = lv_r(x) + i = \max(lv_l(x), lv_l(y) + 1) + i = lv_l(x) + i$. If $lv_l(x) \leq lv_l(y)$, we associate y to x . Then, we note that x is local in M_l iff y is local in M_r ; it follows that $lv'_r(y) = lv_r(y) + i = lv_l(y) + i \geq lv_l(x) + i = lv'_l(x)$.
- We map y to x as well. Indeed, $lv'_l(y) = lv_l(y) + 2 \leq lv_r(x) + 1 \leq lv'_r(x)$.
- Any other variable is unchanged, its weighted level unaffected by the reduction.

Likewise, we give a mapping of variables in M_r to variables in M_l .

- We map x to x if $lv_l(x) \geq lv_l(y) + 1$ (then we have $lv'_r(x) = lv_r(x) + 1 = \max(lv_l(x), lv_l(y) + 1) + 1 = lv_l(x) + 1 = lv'_l(x)$), and y if $lv_l(x) \leq lv_l(y)$ (then $lv'_r(x) = lv_l(y) + 2 = lv'_l(y)$).
 - We map y and y' to y , the weighted level can only increase.
 - To any other z we leave it unchanged, its weighted level is unaffected.
- For all other cases, it directly follows from IH. □

For preservation of local height, we need a few preliminaries. If M is a term and x is a variable of M , we say that x is a **carrier variable** iff there is a generalized redex $(\lambda x, N)$ in M such that N has a free variable. We also define the **V -weighted local height** of M , parametrized by a set V of variables and written $lh_V(M)$, by induction on M as for lh except that the base case for variable occurrences is enriched with:

$$lh_V(x_0 M_1 \dots M_n) = 1 + \max(1, \max_{1 \leq i \leq n} lh_V(M_i))$$

when x_0 is an occurrence of $x \in V$. The **weighted local height** of M , written $lh'(M)$, is defined as $lh_V(M)$ where V is the set of carrier variables of M .

First, we show how the weighted local height compares with the local height.

Lemma 52. For any term M , $lh(M) \leq lh'(M) \leq lh(M) + 1$.

Proof. Immediate by induction on the length of M . □

Lemma 53. If M is a term such that $M \rightarrow_{\lambda l} M'$, then $lh'(M) = lh'(M')$.

Therefore (by Lemma 52) that for any term M , if $M \rightarrow_{\lambda l}^* M'$, then $lh(M') \leq lh(M) + 1$.

Proof. By induction on the definition of λl , detailing non-trivial cases.

- If we have, with $y \in fv(M_1)$:

$$M_l = (\lambda x. M) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda x. M[x y/x]) (\lambda y'. M_1[y'/y]) \dots M_n = M_r$$

Then, it is direct to show by induction on the length of a term M that if x, y are not bound in M and $x \in V$, $y \notin V$, and if V' is either V or $V \setminus \{x\}$, then $lh_V(M) = lh_{V'}(M[x y/x])$. Since y is free in M_l it is not a carrier variable, so applying the property above it is direct that $lh'(M_l) = lh'(M_r)$.

- Suppose we have $(\lambda x. M) M_1 \dots M_n \rightarrow_{\lambda l} (\lambda x. M') M_1 M'_2 \dots M'_n$ from $M_l = M M_2 \dots M_n \rightarrow_{\lambda l} M' M'_2 \dots M'_n$ and M_1 closed. Let V be the set of carrier variables of M_l . Since M_1 is closed, $x \notin V$. We calculate:

$$lh_V((\lambda x. M) M_1 \dots M_n) = \max(lh_V(M M_2 \dots M_n), lh_V(M_1))$$

But $lh_V(M M_2 \dots M_n) = lh'(M M_2 \dots M_n)$: by Barendregt's convention variables of V appearing in $M M_2 \dots M_n$ are bound in $M M_2 \dots M_n$ and therefore be carrier variables in $M M_2 \dots M_n$. Therefore, the property follows from IH. □

Putting all of the above together, we obtain the following lemma:

Lemma 54. For any closed term M , there is a strongly locally scoped M' such that:

$$\begin{array}{ll} \mathcal{N}(M') \geq \mathcal{N}(M) & \text{ord}(M') \leq \text{ord}(M) + 1 \\ \text{lh}(M') \leq \text{lh}(M) + 1 & \text{depth}(M') = \text{depth}(M) \end{array}$$

Proof. Starting from M , apply $\rightarrow_{\lambda 1}$ as much as possible. By Lemma 44, this reduction must terminate on a term M' . By Lemma 45, M' is strongly locally scoped. By Proposition 12, we have $\mathcal{N}(M') \geq \mathcal{N}(M)$. By Lemmas 51 and 53, we have $\text{lh}(M') \leq \text{lh}(M) + 1$ and $\text{ord}(M') \leq \text{ord}(M) + 1$. Finally by Lemma 49, we have $\text{depth}(M') = \text{depth}(M)$. \square

5.5.4. *Expanding variables.* Until now, terms have been measured through their local height and depth. However for general terms those are rather unnatural quantities, and bounds for λ -calculi are usually not expressed in terms of them.

The **height** of a term M , is the quantity defined by induction by $\text{h}(\ast) = 0, \text{h}(x_0) = 1, \text{h}(\lambda x. M) = \text{h}(M), \text{h}(M_1 M_2) = \max(\text{h}(M_1), \text{h}(M_2) + 1)$. We give a final norm-increasing term transformation, allowing us to convert depth and local height to height.

Lemma 55. For any term M , there exists a term M' such that $M \rightarrow_{\eta}^{\ast} M'$,

$$\begin{array}{ll} \text{lh}(M') \leq 2 & \text{depth}(M') \leq \text{h}(M) \\ \text{ord}(M') = \text{ord}(M) & \mathcal{N}(M') \geq \mathcal{N}(M) \end{array}$$

Proof. We define the *expansion* $\text{exp}(M)$ by induction on M as follows:

$$\begin{aligned} \text{exp}(x_0) &= \lambda y^{1A_1}. \dots \lambda y^{nA_n}. x_0 y_1 \dots y_n \\ \text{exp}(\ast) &= \ast \\ \text{exp}(\lambda x. M) &= \lambda x. \text{exp}(M) \\ \text{exp}(M N) &= \text{exp}(M) \text{exp}(N) \end{aligned}$$

where in the first case, x has type $A_1 \rightarrow \dots \rightarrow A_n \rightarrow o$. It is direct that $M \rightarrow_{\eta}^{\ast} \text{exp}(M)$, so by Proposition 9 and Lemma 41 we have $\mathcal{N}(\text{exp}(M)) \geq \mathcal{N}(M)$ and $\text{ord}(\text{exp}(M)) = \text{ord}(M)$. Finally, $\text{lh}(\text{exp}(M)) \leq 2$ and $\text{depth}(\text{exp}(M)) \leq \text{h}(M)$ are immediate by induction. \square

5.5.5. *Exact bounds for general terms.* We are now interested in estimating the quantity:

$$\text{Lgen}_n(h) = \max\{\mathcal{N}(M) \mid \text{ord}(M) \leq n \ \& \ \text{h}(M) \leq h\}$$

We do that by applying the tools developed earlier to get an upper bound on the length of reduction, and then prove a matching lower bound by providing terms whose length of reduction asymptotically reaches the upper bound.

Proposition 13 (Upper bound). Suppose M is a term. Then, $\mathcal{N}(M) \leq 2^{\frac{\text{h}(M) \log(\text{ord}(M)+5)}{\text{ord}(M)}}$.

Proof. Start with a term M , call it M_0 . Without loss of generality we can consider M closed, otherwise we replace occurrences of free variables by occurrences of constants without changing the norm, and only reducing the local height, depth and order. Then, we apply the following transformations.

First we expand variables: by Lemma 55, there is a term closed term M_1 with:

$$\begin{array}{ll} \mathcal{N}(M_1) \geq \mathcal{N}(M_0) & \text{ord}(M_1) = \text{ord}(M_0) \\ \text{lh}(M_1) \leq 2 & \text{depth}(M_1) \leq \text{h}(M_0) \end{array}$$

Then, we make it strongly locally scoped. By Lemma 54, there is M_2 sls such that:

$$\begin{array}{ll} \mathcal{N}(M_2) \geq \mathcal{N}(M_1) & \text{ord}(M_2) \leq \text{ord}(M_1) + 1 \\ \text{lh}(M_2) \leq \text{lh}(M_1) + 1 & \text{depth}(M_2) = \text{depth}(M_1) \end{array}$$

If $\text{ord}(M_2) = 0$, then M_2 does not have any prime redex (nor any application, in fact), so $\mathcal{N}(M_2) = 0$. Otherwise, by Proposition 11, $\mathcal{N}(M_2) \leq 2^{\frac{\text{depth}(M_2) \log(\text{lh}(M_2) + \text{ord}(M_2) + 1)}{\text{ord}(M_2) - 1}}$. By the inequalities above, we immediately deduce that $\mathcal{N}(M) \leq 2^{\frac{\text{h}(M) \log(\text{ord}(M) + 5)}{\text{ord}(M)}}$. \square

Theorem 5. For fixed $n \geq 3$ we have $\text{Lgen}_n(h) = 2_n^{\Theta(h)}$.

Proof. Upper bound. By Proposition 13.

Lower bound. The construction is essentially the same as the one used in [4] for the lower bound in terms of height. For $p \geq 1$ and $k \geq 0$, we define $b_0^p = \underline{2}_p$ and $b_{k+1}^p = \lambda x^{A_{p-1}}. b_k^p (b_k^p x)$. Then, we set:

$$B_k^p = b_k^p \underline{2}_{p-1} \cdots \underline{2}_0$$

Note that this term is *not* sls. By standard arithmetic of Church numerals, we have that for any $p \geq 1, k \geq 0$, $B_k^p \rightarrow_{\beta}^* \underline{2}_{p+2}^k$. By Lemma 43 it follows that $\mathcal{N}(B_k^p \text{id}_o) \geq 2_{p+2}^k$. It is direct to check that $\text{ord}(B_k^p) = p+2$ and $\text{h}(B_k^p) = k+3$ (for $k \geq 1$), concluding the proof. \square

For a term M of height h and order n , Beckmann's results [4] predict that any β -reduction chain of M terminates in less than $2_{n+1}^{\Theta(h)}$ steps. It might seem counter-intuitive that our bound (with lhr) is smaller than Beckmann's (with β -reduction) since we substitute only one occurrence at a time, which is obviously longer. However, Beckmann considers arbitrary β -reduction, not head β -reduction. The possibility of reducing in arbitrary locations of the term unlocks much longer reductions, since higher-order free variables or constants can isolate sections of the term that will never arrive in head position but can still be affected by arbitrary β -reduction. The fact that the length of lhr has the same order of magnitude as head β -reduction is not surprising in the light of Accattoli and Dal Lago's recent result [3] that a similar notion of lhr is quadratically related to head reduction.

Acknowledgment. The author is grateful to the anonymous referees for their very helpful comments and suggestions.

REFERENCES

- [1] Samson Abramsky and Guy McCusker. Linearity, sharing and state: a fully abstract game semantics for idealized algol with active expressions. *Electr. Notes Theor. Comput. Sci.*, 3:2–14, 1996.
- [2] Beniamino Accattoli. An abstract factorization theorem for explicit substitutions. In Tiwari [20], pages 6–21.

- [3] Beniamino Accattoli and Ugo Dal Lago. On the invariance of the unitary cost model for head reduction. In Tiwari [20], pages 22–37.
- [4] Arnold Beckmann. Exact bounds for lengths of reductions in typed lambda-calculus. *J. Symb. Log.*, 66(3):1277–1285, 2001.
- [5] Alexis Bernadet and Stéphane Lengrand. Complexity of strongly normalising λ -terms via non-idempotent intersection types. In *FOSSACS*, pages 88–107, 2011.
- [6] Pierre Clairambault. Estimation of the length of interactions in arena game semantics. In *FOSSACS*, pages 335–349, 2011.
- [7] Pierre Clairambault. Bounding skeletons, locally scoped terms and exact bounds for linear head reduction. In Masahito Hasegawa, editor, *TLCA*, volume 7941 of *Lecture Notes in Computer Science*, pages 109–124. Springer, 2013.
- [8] Pierre Clairambault and Russ Harmer. Totality in arena games. *Ann. Pure Appl. Logic*, 161(5):673–689, 2010.
- [9] V. Danos, H. Herbelin, and L. Regnier. Game semantics and abstract machines. In *Logic in Computer Science, 1996. LICS'96. Proceedings., Eleventh Annual IEEE Symposium on*, pages 394–405. IEEE, 1996.
- [10] V. Danos and L. Regnier. How abstract machines implement head linear reduction. 2003. unpublished.
- [11] Vincent Danos and Jean-Baptiste Joinet. Linear logic and elementary time. *Inf. Comput.*, 183(1):123–137, 2003.
- [12] Daniel de Carvalho, Michele Pagani, and Lorenzo Tortora de Falco. A semantic measure of the execution time in linear logic. *TCS*, 412(20):1884–1902, 2011.
- [13] Jean-Yves Girard. Light linear logic. In Daniel Leivant, editor, *LCC*, volume 960 of *Lecture Notes in Computer Science*, pages 145–176. Springer, 1994.
- [14] Russ Harmer. Innocent game semantics. *Lecture notes*, 2004–2007.
- [15] J. M. E. Hyland and C.-H. Luke Ong. On full abstraction for PCF: I, II, and III. *Inf. Comput.*, 163(2):285–408, 2000.
- [16] Thomas Johnsson. Lambda lifting: Transforming programs to recursive equations. In *FPCA*, pages 190–203, 1985.
- [17] Ugo Dal Lago and Olivier Laurent. Quantitative game semantics for linear logic. In Michael Kaminski and Simone Martini, editors, *CSL*, volume 5213 of *Lecture Notes in Computer Science*, pages 230–245. Springer, 2008.
- [18] R. Nakajima. Infinite normal forms for the λ -calculus. *λ -Calculus and Computer Science Theory*, pages 62–82, 1975.
- [19] H. Schwichtenberg. An upper bound for reduction sequences in the typed λ -calculus. *Archive for Mathematical Logic*, 30(5):405–408, 1991.
- [20] Ashish Tiwari, editor. *23rd International Conference on Rewriting Techniques and Applications (RTA'12)*, *RTA 2012, May 28 - June 2, 2012, Nagoya, Japan*, volume 15 of *LIPICs*. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2012.