

ACTIONS OF NONCOMPACT SEMISIMPLE GROUPS ON LORENTZ MANIFOLDS

MOHAMED DEFFAF, KARIN MELNICK AND ABDELGHANI ZEGHIB

Abstract. The above title is the same, but with “semisimple” instead of “simple,” as that of a notice by Nadine Kowalsky. There, she announced many theorems on the subject of actions of simple Lie groups preserving a Lorentz structure. Unfortunately, she published proofs for essentially only half of the announced results before her premature death. Here, using a different, geometric approach, we generalize her results to the semisimple case, and give proofs of all her announced results.

1 Introduction

Isometric actions on Lorentz manifolds were first investigated in the compact case (see [Zi2], [G], [AS1,2], [Z3,4]). The natural question was then: how can a compact Lorentz manifold have a noncompact isometry group? There is strong evidence that such a question is in fact “decidable” for a wide class of geometric structures (see, for instance [DG], [Zi1]).

1.1 Framework. One new aspect of Kowalsky’s work was to deal with actions of groups on *noncompact* Lorentz manifolds. Obviously, nothing can be said about such actions without compensating for noncompactness with a dynamical counterpart ensuring some kind of recurrence. A natural and rather weak condition used by Kowalsky is nonproperness of the action.

The noncompact case is important, at least from a physical point of view, according to which compact spacetimes have little interest. Having a nonproper isometry group is a manifestation of the non-Riemannian character of the geometry of spacetime. It is in such spaces that one can observe “dilation of length” and “contraction of time.” It is surely interesting to try to classify spacetimes with nonproper isometry groups. This job, however,

Keywords and phrases: Lorentz isometry, Lorentz manifold, de Sitter space, Minkowski space, anti-de Sitter space

AMS Mathematics Subject Classification: 53 (differential geometry), 37 (dynamical systems), 22 (Lie groups)

does not seem to be easy. Some extra hypotheses are therefore in order. Kowalsky restricted her study to actions of *simple Lie groups*.

1.2 Kowalsky's main theorem. The de Sitter and anti-de Sitter spaces, dS_n and AdS_{n+1} , respectively, are the homogeneous spaces $O(1, n)/O(1, n-1)$ and $O(2, n)/O(1, n)$. Geometrically, they are the universal Lorentz spaces of constant positive and negative curvature, respectively. A striking fact proved by Kowalsky is that, *at a group level*, they are the only Lorentz nonproper G -spaces, with G simple.

Theorem 1.1 (Kowalsky [K1, 5.1]). *Let G be a simple Lie group with finite center acting isometrically and nonproperly on a connected Lorentz manifold. Then G is locally isomorphic to either $O(1, n)$, $n \geq 2$, or $O(2, n)$, $n \geq 3$.*

REMARK 1.2. The groups $O(1, 1)$ and $O(2, 2)$ are not simple.

1.3 Geometry of semisimple isometric actions. Once the acting group is known, the problem arises of understanding the geometry of the Lorentz space, or at least that of orbits. Here, one hopes the space looks like de Sitter or anti-de Sitter space, depending on whether G is locally $O(1, n)$ or $O(2, n)$. Nadine Kowalsky announced results to this effect in [K2] and wrote proofs for the $O(1, n)$ case in her thesis [K3]. We will recall their statements below in section 1.7. Unfortunately, she died before publishing proofs.

1.4 The technique. When a Lie group G acts on M preserving a pseudo-Riemannian metric, one can consider a Gauss map from M to $S^2(\mathfrak{g})$, the space of quadratic forms on the Lie algebra \mathfrak{g} of G . When $S^2(\mathfrak{g})$ is endowed with the natural adjoint G -action, the Gauss map is equivariant, and the G -space $S^2(\mathfrak{g})$ reflects the dynamics on M . It is via this map that the non-properness condition is translated as a geometric condition on the induced metrics on orbits. This idea, due to Kowalsky, has become a basic tool in similar questions on the subject, e.g. Adams–Stuck [AS1,2], and Bader–Nevo [BN]. (Note here that variants of the Gauss map, with other natural spaces instead of $S^2(\mathfrak{g})$ were used by other authors, e.g. Gromov [G] and Zimmer [Zi1].) However, this is the starting point; further work in the proof is algebraic and Lie theoretic.

1.5 Other works. Another proof of Theorem 1.1 was proposed by S. Adams [A3]; his methods involve an analysis similar to Kowalsky's, except that zero jets of germs of Killing vector fields are replaced by higher order jets. In other directions, Adams investigated Lorentz-isometric actions,

for simply connected Lie groups, with the stronger dynamical condition that some orbit either is not closed or has noncompact stabilizers ([A1,2]).

Concerning Kowalsky's unpublished proofs, we note a contribution by D. Witte Morris [W], in which he considers the homogeneous case. More precisely, he takes G locally isomorphic to the isometry group of de Sitter or anti-de Sitter space, respectively, and considers a nonproper Lorentz homogeneous space G/H – that is, H is noncompact, and the G -action on G/H preserves some Lorentz metric. He proves that \mathfrak{h} is isomorphic to a standard copy of $\mathfrak{o}(1, n - 1)$ in $\mathfrak{o}(1, n)$ or of $\mathfrak{o}(1, n)$ in $\mathfrak{o}(2, n)$; it follows that G/H is locally homothetic to de Sitter or anti-de Sitter space. Witte Morris' proof is quite algebraic.

1.6 On the present contribution. Our investigation here relies heavily on the approach of [ArDZ], although the two articles can be read completely independently. From [ArDZ], we will use the result recalled below as Theorem 2.2. In light of this result, on the structure of nonproper orbits of Lorentz type, the present paper addresses the case in which the acting group has a nonproper degenerate orbit.

Before stating our results, let us give some motivations and emphasize new features:

- *Completing Kowalsky's work:* One major goal is to prove the announced results of Nadine Kowalsky.

- *Geometric approach:* The approach here is different from that of Kowalsky (as well as from others', for instance Adams'). Together with [ArDZ], we get proofs of the main results, in particular, of Theorem 1.1, using many geometric arguments, where one sees the global structure of proofs.

- *From simple to semisimple:* More importantly, we generalize results to the semisimple case, assuming there are no local $SL_2(\mathbf{R})$ -factors. A semisimple Lie group is essentially a product of simple Lie groups, but, in general, a nonproper action of a product does not derive from a nonproper action of one factor. However, in the Lorentz setting, we conclude that it necessarily does – that is, the semisimple case reduces to the simple one. This is really an important fact, since it leads one to hope to reduce the remaining work to the case in which the group is solvable. Of course, the reason for this is the Levi decomposition of Lie groups, which says that a Lie group is essentially a semidirect product of a semisimple and a solvable group.

1.7 Kowalsky's legacy. In [K2], the following theorems are stated. For Theorem 1.3 below, see also [K3, 6.2]. The manifold M and group G are assumed connected throughout.

Theorem 1.3. *Let G be locally isomorphic to $O(1, n)$, $n \geq 3$, and suppose that G acts on a manifold M preserving a Lorentz metric. Then all noncompact stabilizers have a Lie algebra isomorphic to either $\mathfrak{o}(1, n)$, $\mathfrak{o}(1, n - 1)$, or $\mathfrak{o}(n - 1) \times \mathbf{R}^{n-1}$.*

Theorem 1.4. *Let G be locally isomorphic to $O(2, n)$, $n \geq 6$, with G having finite center. Suppose that G acts nontrivially on a manifold M preserving a Lorentz metric. Then all noncompact stabilizers have a Lie algebra isomorphic to $\mathfrak{o}(1, n)$.*

Theorem 1.5. *Let G and M be as in Theorem 1.4 above, and assume there is a point of M with noncompact stabilizer. Then the universal cover \widetilde{M} is Lorentz isometric to a warped product $L \times_w \widetilde{AdS}_{n+1}$, where \widetilde{AdS}_{n+1} is the simply connected $(n + 1)$ -dimensional Lorentz space of constant curvature -1 , and L is a Riemannian manifold. Further, the induced action of the universal cover \widetilde{G} on \widetilde{M} is via the canonical \widetilde{G} -action on \widetilde{AdS}_{n+1} and the trivial action on L .*

See section 2 below for the definition of warped product.

1.8 Results. As said above in section 1.6, here we provide proofs of all of Kowalsky's previous statements, together with some improvements.

A submanifold N in a Lorentz manifold is *degenerate* if $T_x N^\perp \cap T_x N \neq \mathbf{0}$. In Minkowski space $\mathbf{R}^{1,n}$, the simple subgroup $O(1, n) \subset \text{Isom}(\mathbf{R}^{1,n})$ has one degenerate orbit, which, together with the origin, forms the *light cone*, the set of all isotropic vectors in $\mathbf{R}^{1,n}$. The stabilizer in $O(1, n)$ of a nonzero vector in the light cone is isomorphic to $O(n - 1) \times \mathbf{R}^{n-1}$, where the action of $O(n - 1)$ on \mathbf{R}^{n-1} is the usual representation.

In the degenerate case, we have the following theorem, which says that a degenerate orbit for a simple group acting nonproperly is locally homothetic to the Minkowski light cone. Together with Theorem 1.5 of [ArDZ], which classifies nonproper orbits of Lorentz type, it implies Kowalsky's Theorems 1.3 and 1.4 above.

Theorem 1.6. *Suppose G is a connected group with finite center, locally isomorphic to $O(1, n)$ or $O(2, n)$ for $n \geq 3$. If G acts isometrically on a Lorentz manifold and has a degenerate orbit O with noncompact stabilizer $G(x)$, then*

- (1) G is locally isomorphic to $O(1, n)$;

- (2) The Lie algebra $\mathfrak{g}(x)$ is isomorphic to $\mathfrak{o}(n - 1) \ltimes \mathbf{R}^{n-1}$;
- (3) The orbit O is locally homothetic to the light cone in Minkowski space.

The following result implies Theorem 1.5 above.

Theorem 1.7. *If G , a group with finite center locally isomorphic to $O(2, n)$, $n \geq 3$, acts isometrically and nontrivially on a Lorentz manifold M , with some noncompact stabilizer, then, up to finite covers, M is equivariantly isometric to a warped product $L \times_w AdS_{n+1}$ of a Riemannian manifold L with the anti-de Sitter space AdS_{n+1} .*

We extend the above results to semisimple groups. Note that the noncompact stabilizer assumption is weakened to nonproperness of the action. A local factor of a semisimple Lie group G is a Lie group G_1 such that the Lie algebra \mathfrak{g}_1 is a direct summand of \mathfrak{g} . If $|Z(G)| < \infty$ and G acts on a manifold M , then a finite cover of any local factor G_1 acts on M . Below we will abuse terminology slightly by referring to the “ G_1 -action” and “ G_1 -orbits” on M .

Theorem 1.8. *Let G be a semisimple group with finite center and no local $SL_2(\mathbf{R})$ -factor, acting isometrically, faithfully, and nonproperly on a Lorentz manifold M . Then*

- (1) G has a local factor G_1 isomorphic to $O(1, n)$ or $O(2, n)$;
- (2) There exists a Lorentz manifold S , isometric, up to finite cover, to dS_n or AdS_{n+1} , depending whether G_1 is isomorphic to $O(1, n)$ or $O(2, n)$, and an open subset of M , in which each G_1 -orbit is homothetic to S ;
- (3) Any such orbit as above has a G_1 -invariant neighborhood isometric to a warped product $L \times_w S$, for L a Riemannian manifold.

For $O(1, n)$ -actions, we can also describe orbits with compact isotropy (Proposition 8.2). Nonproper $O(1, n)$ -actions for which compact isotropy occurs strongly resemble the standard action on $\mathbf{R}^{1, n}$. As a fusion, we can give the following “full” theorem:

Theorem 1.9. *Let G be a semisimple Lie group with finite center and no local $SL_2(\mathbf{R})$ -factor, acting nonproperly and isometrically on a Lorentz manifold M . Then, G has a simple local factor G_1 that acts nonproperly. There are two possibilities for G_1 :*

- (1) $G_1 \cong O(2, n)$. In this case, there is a Lorentz manifold S isometric, up to finite cover, to AdS_{n+1} , such that all G_1 -orbits are homothetic to S . In fact, up to finite cover, M is a warped product $L \times_w AdS_{n+1}$.
- (2) $G_1 \cong O(1, n)$. There are open sets U and V such that $M = U \sqcup \partial U \sqcup V$, where

- For any $x \in U$, there exists S , isometric to dS_n up to finite cover, such that the G_1 -orbit of x is homothetic to S . The component of x in U is G_1 -equivariantly isometric to a warped product $L \times_w S$, for some Riemannian manifold L .
- Orbits of G_1 on the boundary of U are either fixed points or locally homothetic to the light cone in Minkowski space $\mathbf{R}^{1,n}$; further $\partial U = \partial V$.
- For any $x \in V$, the G_1 -orbit of x is homothetic to \mathbf{H}^n . The set V is globally G_1 -equivariantly isometric to a warped product $L \times_w \mathbf{H}^n$, for some Lorentz manifold L .

The following two examples of nonproper $SL_2(\mathbf{R})$ -actions illustrate the necessity of the hypothesis of no $SL_2(\mathbf{R})$ -factors.

EXAMPLE 1.10. In this example, the manifold has constant curvature, but has no common finite cover with any of the constant-curvature models. The group $PSL_2(\mathbf{R})$ with the Lorentz metric arising from the Killing form is isometric to AdS_3 . For Γ a cocompact lattice in $PSL_2(\mathbf{R})$, the manifold $M = AdS_3/\Gamma$ admits an isometric, nonproper left-action by $PSL_2(\mathbf{R})$.

EXAMPLE 1.11. This example is a transitive, nonproper, isometric $SL_2(\mathbf{R})$ -action on a manifold with nonconstant curvature. Let g be the Killing form on $\mathfrak{sl}_2(\mathbf{R})$. Let λ_1, λ_2 and λ_3 be the standard basis for $(\mathbf{R}^3)^*$. There is an isomorphism $\mathfrak{sl}_2(\mathbf{R}) \cong \mathbf{R}^3$ for which the Killing form is $2\lambda_1 \cdot \lambda_2 + \lambda_3^2$. Under this isomorphism, an \mathbf{R} -split element of $\mathfrak{sl}_2(\mathbf{R})$ maps to the basis element \mathbf{e}_3 . Let $c \neq 1$ be a positive number. Let g' be the pullback of the form $2\lambda_1 \cdot \lambda_2 + c\lambda_3^2$ to $\mathfrak{sl}_2(\mathbf{R})$. The adjoint action of the \mathbf{R} -split torus A of $SL_2(\mathbf{R})$ preserves g' . Let Γ be a lattice in A . The form g' gives rise to an $SL_2(\mathbf{R})$ -invariant Lorentz metric on $M = SL_2(\mathbf{R})/\Gamma$. The isometric $SL_2(\mathbf{R})$ -action is nonproper, but M does not have constant curvature.

For Lorentz-isometric actions of $SL_2(\mathbf{R})$ on finite-volume manifolds, Gromov has shown that all stabilizers are discrete, and the universal cover is isometric to a warped product $L \times_w \widetilde{AdS}_3$ ([G, 5.4.A]).

This final example illustrates the necessity for the results above of the hypothesis that G has finite center. We thank the referee for bringing this example to our attention.

EXAMPLE 1.12. The main result of [A4] implies that, if Z is a central closed subgroup of a Lie group G , and if Z acts isometrically on a Lorentz manifold M , then the manifold $G \times_Z M$ admits a G -invariant Lorentz

metric. There exists a Lorentz manifold M , diffeomorphic to a torus, with an Anosov action of the integers \mathbf{Z} by isometries (see [BaZ, §6.3.3]); in fact, this action has a fixed point. Then $\tilde{O}(2, n)$ acts isometrically and nonproperly on $\tilde{O}(2, n) \times_{\mathbf{Z}} M$. All stabilizers are discrete, and there is obviously no warped product as in the theorems above.

Notation and terminology. The Lie algebra of a Lie group G will be denoted \mathfrak{g} . The stabilizer subgroup in G and corresponding subalgebra of a point x will be denoted $G(x)$ and $\mathfrak{g}(x)$, respectively.

Lie groups G and H are *locally isomorphic* if $\mathfrak{g} \cong \mathfrak{h}$. As discussed above, a group G_1 is a *local factor* of G if \mathfrak{g}_1 is a direct summand of \mathfrak{g} .

The dimension of the Lorentz manifold M will be denoted d throughout.

2 Background: Warped Product near Lorentz Orbits

DEFINITION 2.1. For two pseudo-Riemannian manifolds (L, λ) and (S, σ) , a warped product $L \times_w S$ is given by a positive function w on L : the metric at (l, s) is $\lambda_l + w(l)\sigma_s$.

2.1 Results of [ArDZ]. We will make use of the following theorem:

Theorem 2.2 [ArDZ, 1.5]. Let G be a connected semisimple Lie group acting isometrically on a Lorentz manifold M of dimension ≥ 3 . Suppose that no local factor of G is isomorphic to $SL_2(\mathbf{R})$ and that there exists an orbit O of Lorentz type with noncompact isotropy.

Then, up to a finite cover, G factors $G \cong G_2 \times G_1$, where

- (1) G_1 possesses an orbit O_1 which is a Lorentz space of constant, non-vanishing curvature, and G_1 equals $\text{Isom}^0(O_1)$;
- (2) There is a G -invariant neighborhood U of O_1 which is a warped product $L \times_w O_1$;
- (3) The factor O_1 corresponds to G_1 -orbits, and G_2 acts along the L -factor.

3 Properties of the Isotropy Representation

Here we collect some algebraic facts about the structure of nonproper degenerate orbits. Suppose that x is a point of M with degenerate G -orbit. Denote this orbit by O . Recall that d is the dimension of M , and assume that G is semisimple.

Fix an isometric isomorphism of $T_x M$ with $\mathbf{R}^{1, d-1}$, determining an isomorphism $O(T_x M) \cong O(1, d-1)$. Let V be the image of $T_x O$ in $\mathbf{R}^{1, d-1}$.

Let $\Phi : G(x) \rightarrow O(1, d - 1)$ be the resulting isotropy representation. Because G acts properly and freely on the bundle of Lorentz frames of M , the isotropy representation is an injective, proper map. The invariant subspace V is degenerate, so $\Phi(G(x))$ leaves invariant the line $V^\perp \cap V$. The stabilizer of an isotropic line is conjugate in $O(1, d - 1)$ to the parabolic

$$P = (K \times A) \ltimes U$$

where $U \cong \mathbf{R}^{d-2}$ is unipotent, $A \cong \mathbf{R}^*$, and $K \cong O(d - 2)$ with the conjugation action of $K \times A$ on U equivalent to the standard representation of the conformal group of \mathbf{R}^{d-2} . Denote by \mathfrak{p} the Lie algebra of P , and by \mathfrak{k} , \mathfrak{a} , \mathfrak{u} , the subalgebras corresponding to K , A and U . Let $\varphi : \mathfrak{g}(x) \rightarrow \mathfrak{o}(1, d - 1)$ be the Lie algebra representation tangent to Φ .

Note that $\mathfrak{g}/\mathfrak{g}(x)$ can be identified with $T_x O$ by the map

$$Y \mapsto \overline{Y}(x) = \left. \frac{\partial}{\partial t} \right|_0 e^{tY} x.$$

For $g \in G(x)$, differentiating the relation $ge^{tY}x = (ge^{tY}g^{-1})x$ gives

$$D_x g(\overline{Y}(x)) = \overline{\text{Ad}g(Y)}(x).$$

In other words, Φ restricted to V is equivalent to the representation $\overline{\text{Ad}}$ of $G(x)$ on $\mathfrak{g}/\mathfrak{g}(x)$ arising from the adjoint representation. Let $\overline{\text{ad}}$ be the representation tangent to $\overline{\text{Ad}}$.

An element Y of \mathfrak{g} is called *nilpotent* if $\text{ad}(Y)$ is nilpotent. An element Y is *\mathbf{R} -split* if $\text{ad}(Y)$ is diagonalizable over \mathbf{R} .

LEMMA 3.1. *The stabilizer subalgebra $\mathfrak{g}(x) \subset \mathfrak{g}$ has the following properties:*

- (1) *For all $Y \in \mathfrak{g}(x)$, the endomorphism $\overline{\text{ad}}(Y)$ has no real nonzero eigenvalues. In fact, the same is true for φ , so $\varphi(\mathfrak{g}(x))$ is conjugate to a subalgebra of $\mathfrak{k} \ltimes \mathfrak{u}$;*
- (2) *The stabilizer subalgebra $\mathfrak{g}(x)$ contains no element \mathbf{R} -split in \mathfrak{g} ;*
- (3) *There exists a subalgebra $\mathfrak{s}(x)$ in which $\mathfrak{g}(x)$ has codimension one such that $[\mathfrak{s}(x), \mathfrak{s}(x)] \subset \mathfrak{g}(x)$, and the representation of $\mathfrak{g}(x)$ on $\mathfrak{g}/\mathfrak{s}(x)$ is skew-symmetric – that is, every endomorphism is diagonalizable with purely imaginary eigenvalues.*

Proof. (1) Suppose that $\overline{\text{ad}}(Y)$ has eigenvalue $\lambda > 0$. Then λ is also an eigenvalue of $\varphi(Y)$ on V . Since $\varphi(Y)$ is skew-symmetric on $V/(V \cap V^\perp)$, the generalized eigenspace for λ in V is one-dimensional and equals $V^\perp \cap V$. The trace of $\varphi(Y)$ on V is λ , so the trace of $\overline{\text{ad}}(Y)$ on $\mathfrak{g}/\mathfrak{g}(x)$ is λ .

Next we will show that the trace of $\text{ad}(\varphi(Y))$ on $\varphi(\mathfrak{g}(x))$ is nonnegative. Recall that \mathfrak{p} decomposes $(\mathfrak{a} \times \mathfrak{k}) \ltimes \mathfrak{u}$. The restriction of $\text{ad}(\mathfrak{p})$ to \mathfrak{u} factors through the projection to $\mathfrak{a} \times \mathfrak{k}$, which acts as the standard conformal

representation on \mathbf{R}^{d-2} . If $\varphi(Y)$ has eigenvalue λ on $V \cap V^\perp$, then it projects to $\lambda I + K$ in $\mathfrak{a} + \mathfrak{k} \cong \mathbf{conf}(\mathbf{R}^{d-2})$. Therefore, the characteristic roots of $\text{ad}(\varphi(Y))$ on \mathfrak{u} are in $\lambda + i\mathbf{R}$. On $\mathfrak{p}/\mathfrak{u} \cong \mathfrak{a} \oplus \mathfrak{k}$, the characteristic roots of any adjoint endomorphism are all purely imaginary. Therefore, the characteristic roots of $\text{ad}(\varphi(Y))$ on \mathfrak{p} are in $(\lambda + i\mathbf{R}) \cup i\mathbf{R}$, implying that the trace of $\text{ad}(\varphi(Y))$ is nonnegative on $\varphi(\mathfrak{g}(x))$.

Then the trace of $\text{ad}(Y)$ on $\mathfrak{g}(x)$ is also nonnegative. Finally, the trace of $\text{ad}(Y)$ on \mathfrak{g} is positive, contradicting the unimodularity of \mathfrak{g} . If $\lambda < 0$, the same argument shows that the trace of $\text{ad}(Y)$ on \mathfrak{g} is negative. Therefore, no $\overline{\text{ad}}(Y)$ has any nonzero real eigenvalues, and no $\varphi(Y)$ has any nonzero real eigenvalues on V .

If $\varphi(Y)$ has a nonzero real eigenvalue on $\mathbf{R}^{1,d-1}$, then an eigenvector must be isotropic. It either lies in V or is not orthogonal to $V^\perp \cap V$. In either case, $\varphi(Y)$ has a nonzero real eigenvalue on V , a contradiction.

(2) If an \mathbf{R} -split element $H \in \mathfrak{g}(x)$, then by (1), all root vectors on which $\text{ad}(H)$ is nontrivial must project to $\mathbf{0}$ in $\mathfrak{g}/\mathfrak{g}(x)$. In this case, $\mathfrak{g}(x)$ contains a subalgebra isomorphic to $\mathfrak{sl}_2(\mathbf{R})$. Applying the monomorphism φ would yield a subalgebra isomorphic to $\mathfrak{sl}_2(\mathbf{R})$ in \mathfrak{p} , which is impossible.

(3) Take any Z' spanning $V^\perp \cap V$. Item (1) implies $\varphi(\mathfrak{g}(x))$ annihilates Z' . Take the corresponding vector in $\mathfrak{g}/\mathfrak{g}(x)$, and let Z be any lift to \mathfrak{g} . Then $\text{ad}(\mathfrak{g}(x))(Z) \subseteq \mathfrak{g}(x)$, so $\mathfrak{s}(x) = \mathbf{R}Z + \mathfrak{g}(x)$ is the desired subalgebra.

From the equivalence of $\varphi|_V$ with $\overline{\text{ad}}$, the representations $V/(V \cap V^\perp)$ and $\mathfrak{g}/\mathfrak{s}(x)$ are equivalent. The former is skew-symmetric. □

4 Root Spaces in Isotropy Subalgebra

By a nonproper orbit we will mean one with noncompact isotropy. Theorem 1.8 of [ArDZ] asserts the existence of a nonproper orbit under the assumptions of our Theorem 1.8 above. The proof in [ArDZ] was easily deduced from the following result of [K1].

PROPOSITION 4.1. *If the G -action is nonproper, then there is $x \in M$, and an \mathbf{R} -split element H of \mathfrak{g} such that the negative root space*

$$\Sigma_{\alpha(H) < 0} \mathfrak{g}_\alpha$$

is isotropic at x . If G has noncompact stabilizer at some point y of M , then we may take $y = x$ above.

If x has a nonproper degenerate orbit, the subalgebra $\mathfrak{s}(x)$ of Lemma 3.1 is exactly the maximal subspace $\mathfrak{s} \subseteq \mathfrak{g}$ such that $\{\overline{Y}(x) \in T_x M : Y \in \mathfrak{s}\}$ is an isotropic subspace.

Fact 4.2 of [ArDZ] is that

$$(\Sigma_{\alpha(H) < 0} \mathfrak{g}_\alpha) \cap \mathfrak{g}(x) \neq \mathbf{0}.$$

If this intersection were $\mathbf{0}$, then the subalgebra $\oplus_{\alpha(H) \geq 0} \mathfrak{g}_\alpha$ would have codimension one in \mathfrak{g} . Such a subalgebra could only exist if $\mathfrak{sl}_2(\mathbf{R})$ were a factor of \mathfrak{g} , but our hypotheses exclude this.

REMARK 4.2. Point stabilizers are discrete for the $SL_2(\mathbf{R})$ -actions given in Examples 1.10 and 1.11.

Denote by \mathfrak{a} and Δ the Cartan subalgebra and root system, respectively, of Proposition 4.1. The remainder of this section is devoted to showing the following proposition.

PROPOSITION 4.3. *Suppose $x \in M$ has a nonproper degenerate orbit. Then there exist $J \in \mathfrak{a}$ and $S \subset \Delta$ such that*

- (1) $\mathfrak{s}(x) = \mathbf{R}J + \mathfrak{g}(x)$;
- (2) $\alpha(J) < 0$ for all $\alpha \in S$;
- (3) $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha \subseteq \mathfrak{g}(x)$;
- (4) $\dim(\Sigma_{\alpha \in S} \mathfrak{g}_\alpha) \geq 2$.

Proof. Fix H as in Proposition 4.1. Let

$$\mathbf{0} \neq X \in (\Sigma_{\alpha(H) < 0} \mathfrak{g}_\alpha) \cap \mathfrak{g}(x).$$

There exist $J \in \mathfrak{a}$ and a nilpotent Y in \mathfrak{g} such that $[J, X] = -2X$ and $[X, Y] = J$ (see [S, 2.4.B]). The operator $\overline{\text{ad}}(X)$ is nilpotent; on the other hand, by Lemma 3.1 (3), $\overline{\text{ad}}(X)$ is skew-symmetric on $\mathfrak{g}/\mathfrak{s}(x)$, so $\text{ad}(X)(\mathfrak{g}) \subseteq \mathfrak{s}(x)$. Therefore, J belongs to $\mathfrak{s}(x)$.

Note $J \notin \mathfrak{g}(x)$ by Lemma 3.1 (2). Therefore $\mathfrak{s}(x) = \mathbf{R}J + \mathfrak{g}(x)$, proving (1).

Let S be the set of $\alpha \in \Delta$ such that $\alpha(H) < 0$ and $\alpha(J) < 0$, so (2) is obviously satisfied. From the relation $[J, X] = -2X$, any α such that X has a nontrivial component in \mathfrak{g}_α satisfies $\alpha(J) = -2$, so any such α belongs to S ; in particular S is not empty.

For $\alpha \in S$, we have $\mathfrak{g}_\alpha \subset \mathfrak{s}(x)$ and

$$\mathfrak{g}_\alpha = [J, \mathfrak{g}_\alpha] \subset [\mathfrak{s}(x), \mathfrak{s}(x)] \subset \mathfrak{g}(x),$$

by Lemma 3.1 (3), showing statement (3) above.

Now, replacing X by a nonzero element of some \mathfrak{g}_α , $\alpha \in S$, we may assume that $-J$ is a *basic translation* – that is, there exists $c_\alpha < 0$ such that

$$\alpha(J) = -2 \quad \text{and} \quad \alpha(Z) = c_\alpha \kappa(J, Z),$$

for any $Z \in \mathfrak{a}$, where κ denotes the Killing form. In this case, we have that for any root β , the reflection

$$\sigma_\alpha(\beta) = \beta + \beta(J)\alpha$$

is again a root (see [S, II.5.A]).

Now, to show (4) it suffices to show that $\dim(\mathfrak{g}_\alpha) \geq 2$ or that there exists some $\gamma \neq \alpha$ also in S .

Suppose $\dim(\mathfrak{g}_\alpha) = 1$. The assumption that \mathfrak{g} has no $\mathfrak{sl}_2(\mathbf{R})$ -factor implies that there exists some nonzero root $\delta \neq \alpha$ such that $\delta(J) \neq 0$. We may assume $\delta(J) < 0$. If $\delta(H) < 0$, then $\delta \in S$, as desired. So suppose that $\delta(H) \geq 0$. Now let

$$\gamma = -\sigma_\alpha(\delta) = -\delta - \delta(J)\alpha.$$

Then

$$\gamma(J) = -\delta(J) - \delta(J)\alpha(J) = -\delta(J)(1 - 2) = \delta(J) < 0,$$

and

$$\gamma(H) = -\delta(H) - \delta(J)\alpha(H) < 0,$$

so $\gamma \in S$. □

5 Nonproper Semisimple Actions: Proof of Theorem 1.8

Reduction of the proof. As discussed in the previous section, it is proved in [ArDZ] that G has an orbit O with a noncompact stabilizer. It is also proved (Theorem 1.5) that if O is Lorentzian, then the situation is exactly as described in Theorem 1.8. If an orbit is not Lorentzian, then it is degenerate, fixed, or Riemannian. Riemannian isometries always have compact isotropy, so the existence of noncompact stabilizers implies that either O is degenerate, or some simple local factor fixes it pointwise. The proof would be finished using the following two propositions, which state that in either situation, there is a nonproper Lorentzian orbit.

PROPOSITION 5.1. *Let O be a G -orbit on which some simple noncompact normal subgroup G_1 acts trivially. Then G has (near O) Lorentzian orbits with noncompact isotropy.*

PROPOSITION 5.2. *Let O be a degenerate G -orbit with noncompact isotropy. Then, G has (near O) Lorentzian orbits with noncompact isotropy.*

Proof of Proposition 5.1. Let x be a point of O . The isotropy representation $\Phi : G_1 \rightarrow O(T_x M)$ is faithful and proper. No noncompact simple subgroup of $O(T_x M) \cong O(1, d - 1)$, can preserve an isotropic line.

Therefore, Theorem 1.1 of [BoZ] applies, so $\Phi(G_1)$ preserves an $(n + 1)$ -dimensional Lorentz subspace V_1 of $T_x M$ for some $2 \leq n \leq d - 1$; further, $\Phi(G_1)$ contains a subgroup conjugate to the standard copy of $O^0(1, n)$ in $O(1, d - 1)$. Because $\Phi(G_1)$ is simple and G has no local $SL_2(\mathbf{R})$ -factors, G_1 is locally isomorphic to $O(1, n)$, and $n \geq 3$.

Any spacelike vector $v \in V_1$ has Lorentzian $\Phi(G_1)$ -orbit isometric to dS_n . The stabilizer of v is locally isomorphic to $O(1, n - 1)$; in particular, it is noncompact. The G_1 -orbit of $\exp_x(v)$ is again of Lorentzian type, by the Gauss lemma (see [O, 5.1]). The G -orbit of $\exp_x(v)$ is Lorentzian because it contains a Lorentzian submanifold, and this orbit has noncompact isotropy, as desired. \square

The remainder of the section is devoted to the proof of Proposition 5.2; the orbit O will be assumed degenerate with noncompact isotropy below.

5.1 The asymptotic geodesic hypersurface \mathcal{F}_x .

Fact 5.3. *For $x \in O$, let $\mathbf{R}n_x$ be the null direction in $T_x O$. Then, the orthogonal n_x^\perp is tangent to a lightlike geodesic hypersurface \mathcal{F}_x (defined in a neighborhood of x).*

Proof. Let X be a nilpotent element of $\mathfrak{g}(x)$ given by Proposition 4.3, and consider the isometry $f = e^{tX}$, for some $t \neq 0$. The derivative $D_x f(n_x) = n_x$. Recall that d is the dimension of M ; let g denote the Lorentz metric. The graph $\text{Graph}(f) \subset M \times M$ is an isotropic totally geodesic d -dimensional submanifold of $M \times M$, equipped with the metric $g \oplus (-g)$. The graphs $\text{Graph}(f^m)$ converge to E , a d -dimensional, isotropic, totally geodesic submanifold, which is no longer a graph, since f^m is divergent (see [Z2] or [DG, 7.4]). The intersection $E \cap (\{x\} \times M)$ is nontrivial, but has dimension at most 1, because it is isotropic, and M is Lorentzian. Therefore, the projection \mathcal{F}_x of E is a totally geodesic hypersurface in $M \times \{x\}$. Because the derivative $D_x(f^m)$ fixes n_x for all m , the vector $(n_x, n_x) \in T_{(x,x)}E$. Because $T_{(x,x)}E$ is isotropic, its intersection with $\mathbf{0} \times T_x M$ is exactly $\mathbf{R}n_x$. Then $(n_x, \mathbf{0}) \in T_{(x,x)}E$, so the projection $T_x(\mathcal{F}_x) = n_x^\perp$, as desired. \square

Fact 5.4. *The hypersurface \mathcal{F}_x carries a 1-dimensional foliation \mathcal{C} , such that*

- (1) *Any isotropic curve in \mathcal{F}_x is tangent to a leaf of \mathcal{C} ;*
- (2) *Each leaf of \mathcal{C} is an isotropic geodesic;*
- (3) *The (local) quotient space $\mathcal{F}_x/\mathcal{C}$ inherits a Riemannian metric, infinitesimally preserved by the elements of \mathfrak{g} preserving \mathcal{F}_x .*

Proof. At any point y of a degenerate hypersurface \mathcal{F} , there exists a unique tangent isotropic direction C_y . These lines determine a characteristic 1-dimensional foliation \mathcal{C} of \mathcal{F} , proving (1). Since \mathcal{F}_x is totally geodesic, (2) follows. For (3), it is known (see, for instance, [Z1]) that \mathcal{C} is transversally Riemannian if and only if \mathcal{F}_x is totally geodesic. Here, transversally Riemannian means that the flow along any parameterization of \mathcal{C} preserves the induced degenerate metric, or equivalently, that the degenerate metric can be projected as a Riemannian metric on the (local) quotient space $\mathcal{F}_x/\mathcal{C}$. Finally, if an isometric flow locally preserves \mathcal{F}_x , then it induces a local diffeomorphism of $\mathcal{F}_x/\mathcal{C}$ that is obviously an isometry by construction of the Riemannian metric. \square

Fact 5.5. *The subalgebra $\mathfrak{s}(x)$ preserves the isotropic geodesic \mathcal{C}_x , so it preserves \mathcal{F}_x .*

Proof. Indeed, any $Y \in \mathfrak{s}(x)$ has $\bar{Y}(x)$ isotropic, and hence the whole Y -orbit of x is isotropic. But, as stated above, isotropic curves of \mathcal{F}_x are contained in leaves of \mathcal{C} – that is, all Y -orbits through x are contained in \mathcal{C}_x . The image of \mathcal{C}_x by any element of the one-parameter group generated by Y is an isotropic geodesic tangent to \mathcal{C}_x at some point, thus equals \mathcal{C}_x . \square

Fact 5.6. *The action of $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ on the tangent space of the (local) quotient space $\mathcal{F}_x/\mathcal{C}$ at the point corresponding to \mathcal{C}_x is trivial.*

Proof. The tangent space to $\mathcal{F}_x/\mathcal{C}$ at \mathcal{C}_x is identified with $n_x^\perp/\mathbf{R}n_x$. Note that the subspace $\mathbf{R}J + \Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ as in Proposition 4.3 is in fact a subalgebra of $\mathfrak{s}(x)$. We have a representation ρ of $\mathbf{R}J + \Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ into the orthogonal algebra of $n_x^\perp/\mathbf{R}n_x$, which is endowed with a positive definite inner product. But in such an orthogonal algebra, an equality $[\rho(J), \rho(Y)] = \lambda\rho(Y)$, becomes trivial – that is $\rho(Y) = 0$ (since $\lambda \neq 0$); \square

COROLLARY 5.7. *$\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ acts trivially on the (local) quotient space $\mathcal{F}_x/\mathcal{C}$. That is, $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ preserves individually each leaf of \mathcal{C} .*

Proof. The action of $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ on $\mathcal{F}_x/\mathcal{C}$ is trivial, since it is a Riemannian action with a fixed a point and a trivial derivative at it. \square

COROLLARY 5.8. *Any point of \mathcal{F}_x has a noncompact isotropy algebra.*

Proof. Indeed, $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ has dimension ≥ 2 and has orbits of dimension 1. Therefore, stabilizers are nontrivial. They are not compact since all elements of $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ are nilpotent. \square

Fact 5.9. *Let Γ be the set of fixed points of $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ in \mathcal{F}_x . Then, Γ has an empty interior (in \mathcal{F}_x). In particular, the orbit of any point of $\mathcal{F}_x - \Gamma$ under $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ locally coincides with its \mathcal{C} -leaf.*

Proof. No element of $\Sigma_{\alpha \in S} \mathfrak{g}_\alpha$ can fix points of an open subset of \mathcal{F}_x . Indeed, in general, a Lorentz transformation fixing one point and acting trivially on a tangent lightlike hyperplane at that point has a trivial derivative, and is therefore trivial. \square

COROLLARY 5.10. *No point of \mathcal{F}_x has a spacelike G -orbit.*

Proof. If a point $y \in \mathcal{F}_x$ has a spacelike orbit, then all orbits of points in a neighborhood of y are spacelike, as well. However, any neighborhood of y meets $\mathcal{F}_x - \Gamma$; orbits of points in here cannot be spacelike, because they contain at least one isotropic geodesic. \square

5.2 End of the proof of Proposition 5.2.

Fact 5.11. *The degenerate orbit O cannot be locally contained in \mathcal{F}_x – that is, $\mathcal{F}_x \cap O$ does not contain an open subset of O .*

Proof. Suppose O is locally contained in \mathcal{F}_x . Then the group G locally preserves \mathcal{F}_x . From Corollary 5.7, the infinitesimal action of \mathfrak{g} on the quotient space $Q = \mathcal{F}_x/\mathcal{C}$ is not faithful. More precisely, any factor \mathfrak{b} of \mathfrak{g} which contains an element like J (in Proposition 4.3) must act trivially on Q . However, orbits of \mathfrak{b} cannot be 1-dimensional, since \mathfrak{g} has no $\mathfrak{sl}_2(\mathbf{R})$ -factor. Therefore, \mathfrak{b} acts trivially on \mathcal{F}_x . As in the proof of Fact 5.9, this implies \mathfrak{b} acts trivially on M . \square

Now, from Corollary 5.8, the proof of Proposition 5.2 would be finished once one proves that there is a point of \mathcal{F}_x with a Lorentz orbit. It suffices to show existence of nondegenerate orbits, since from Corollary 5.10, points of \mathcal{F}_x cannot have spacelike orbits. Assume, for a contradiction, that all G -orbits of points of \mathcal{F}_x are degenerate. For any $y \in \mathcal{F}_x - \Gamma$, the orbit Gy locally contains the isotropic geodesic \mathcal{C}_y by Fact 5.9; therefore, $T_y(Gy) \subset n_y^\perp = T_y(\mathcal{F}_x)$. In other words, for any Killing field $X \in \mathfrak{g}$ and any $y \in \mathcal{F}_x - \Gamma$, the evaluation $X(y)$ is tangent to \mathcal{F}_x , so X defines a vector field in this open subset of \mathcal{F}_x . The flow of any $y \in \mathcal{F}_x - \Gamma$ along X for sufficiently short time is again contained in \mathcal{F}_x . In particular, since $x \in \mathcal{F}_x - \Gamma$, the orbit O is locally contained in \mathcal{F}_x , contradicting the previous fact.

COROLLARY 5.12 (from proof). *There is a simple local factor G_1 of G for which the G_1 -orbit of x is a point or degenerate with noncompact stabilizer. In other words, if G has a nonproper orbit that is either a point or*

degenerate, then a subgroup G_1 locally isomorphic to $O(1, n)$ or $O(2, n)$ has an orbit with the same properties.

Proof. First suppose that G has a nonproper fixed or Riemannian orbit O_1 . Then any local factor G_1 in the kernel of the restriction to O_1 has the desired properties.

Now suppose that G has a nonproper degenerate orbit. We have seen that some nilpotent elements stabilizing a point in the degenerate orbit O stabilize a point y with Lorentzian orbit. But, from Theorem 2.2, a Lorentz orbit can be nonproper only if there is a local factor G_1 acting nonproperly with $O_1 = G_1y$ Lorentzian of constant curvature; moreover, there is a warped product $L \times_w O_1$ preserved by G , and G splits up to finite cover as $G_2 \times G_1$. If $X \in \mathfrak{g}(y)$, the projection of X on \mathfrak{g}_2 generates a precompact 1-parameter group, so X cannot be nilpotent unless it belongs to \mathfrak{g}_1 . We infer from this that G_1 acts nonproperly on the degenerate orbit $O = Gx$. Because G_1 is simple and the stabilizer $G_1(x)$ is noncompact, the orbit G_1x must be degenerate or one point. \square

6 Degenerate Stabilizers: Proof of Theorem 1.6

Now as in Theorem 1.6, assume $\mathfrak{g} \cong \mathfrak{o}(1, n)$ or $\mathfrak{o}(2, n)$ for some $n \geq 3$ and that \mathfrak{g} has a degenerate orbit M with noncompact stabilizer $\mathfrak{g}(x)$. It is proved in the first section below that $\mathfrak{g} \cong \mathfrak{o}(1, n)$ and in the second that $\mathfrak{g}(x) \cong \mathfrak{o}(n - 1) \times \mathbf{R}^{n-1}$. The final point of the theorem follows from these two.

6.1 Excluding $\mathfrak{o}(2, n)$. Let J, \mathfrak{a} , and $S \subset \Delta$ be as in Proposition 4.3. Observe that there can be at most one negative root α with $\mathfrak{g}_\alpha \cap \mathfrak{g}(x) \neq \mathbf{0}$. For if $X \in \mathfrak{g}_\alpha \cap \mathfrak{g}(x)$, then let $Y \in \mathfrak{g}_{-\alpha}$ be as in [S, 2.4.B], so $[X, Y] = H_\alpha \in \mathfrak{a}$. Since $\overline{\text{ad}}(X)$ is nilpotent but skew-symmetric on $\mathfrak{g}/\mathfrak{s}(x)$ (Lemma 3.1 (3)), the \mathbf{R} -split element H_α must belong to $\mathfrak{s}(x)$. Any $X' \in \mathfrak{g}_\beta$ would give rise to $H_\beta \in \mathfrak{a} \cap \mathfrak{s}(x)$. If $\alpha \neq \beta$, then for some $c \in \mathbf{R}$, the difference $H_\alpha - cH_\beta$ would be a nonzero \mathbf{R} -split element of $\mathfrak{g}(x)$, contradicting Lemma 3.1 (2).

Suppose that $\mathfrak{g} \cong \mathfrak{o}(2, n)$. Let β and γ be distinct negative roots, each with $(n - 2)$ -dimensional root spaces. The other negative roots are $\beta - \gamma$ and $\beta + \gamma$, with one-dimensional root spaces.

First suppose $X \in \mathfrak{g}_\beta \cap \mathfrak{g}(x)$. Let L be a generator of $\mathfrak{g}_{-\beta-\gamma}$. The adjoint $(\text{ad}X)^2(L)$ is a nonzero element of $\mathfrak{g}_{\beta-\gamma}$. On the other hand, Lemma 3.2 (3) implies that any nilpotent element of $\mathfrak{g}(x)$ has nilpotence order 2 on $\mathfrak{g}/\mathfrak{g}(x)$. Then we would have $\mathfrak{g}_{\beta-\gamma} \cap \mathfrak{g}(x) \neq \mathbf{0}$, a contradiction.

Therefore, X cannot be in \mathfrak{g}_β . The same argument shows X cannot be in \mathfrak{g}_γ . Proposition 4.3 says that $\mathfrak{g}(x)$ contains a sum of negative root spaces with total dimension at least 2, but since $\mathfrak{g}_{\beta-\gamma}$ and $\mathfrak{g}_{\beta+\gamma}$ are 1-dimensional, and because only one can be contained in $\mathfrak{g}(x)$, we have a contradiction.

6.2 Full stabilizer. Now \mathfrak{g} must be $\mathfrak{o}(1, n)$. Let α and J be as above, so $\alpha(J) < 0$ and $\mathfrak{g}_\alpha \subseteq \mathfrak{g}(x)$. Let \mathfrak{k} be the maximal compact subalgebra of the centralizer of J in \mathfrak{g} ; it is isomorphic to $\mathfrak{o}(n-1)$.

Suppose $Y \in \mathfrak{s}(x) \cap \mathfrak{g}_{-\alpha}$. By Lemma 3.1 (3),

$$J \in [Y, \mathfrak{g}_\alpha] \subset [\mathfrak{s}(x), \mathfrak{s}(x)] \subset \mathfrak{g}(x).$$

But this contradicts Lemma 3.1 (2). Therefore, $\mathfrak{s}(x) \cap \mathfrak{g}_{-\alpha} = \mathbf{0}$.

On the other hand, since $\overline{\text{ad}}(X)$ is nilpotent for all $X \in \mathfrak{g}_\alpha$, Lemma 3.1 (3) forces

$$\mathfrak{k} \subset [\mathfrak{g}_{-\alpha}, \mathfrak{g}_\alpha] \subset \text{ad}(\mathfrak{g}_\alpha)(\mathfrak{g}) \subset \mathfrak{s}(x).$$

Since $\mathfrak{g} = \mathfrak{g}_{-\alpha} + \mathfrak{k} + \mathbf{R}J + \mathfrak{g}_\alpha$, the algebra $\mathfrak{s}(x)$ is exactly $\mathfrak{k} + \mathbf{R}J + \mathfrak{g}_\alpha$. Suppose there were $X = cJ + M \in (\mathbf{R}J + \mathfrak{k}) \cap \mathfrak{g}(x)$ for some nonzero $c \in \mathbf{R}$. The subspace $\mathfrak{g}_{-\alpha}$ is $\text{ad}(X)$ -invariant and maps onto $\mathfrak{g}/\mathfrak{s}(x)$. But $\text{ad}(X)$ is clearly not skew-symmetric here, contradicting Lemma 3.1 (3). Therefore, $\mathfrak{g}(x)$ is exactly $\mathfrak{k} + \mathfrak{g}_\alpha$, which is isomorphic as a Lie algebra to $\mathfrak{o}(n-1) \ltimes \mathbf{R}^{n-1}$.

7 Global AdS Warped Product: Proof of Theorem 1.7

Suppose G is locally isomorphic to $O(2, n)$, $n \geq 3$, with finite center, and G acts isometrically on M . By the argument of Kowalsky and the assumption of no local $SL_2(\mathbf{R})$ -factors, we know that there is a G -orbit with noncompact stabilizer (see section 4). By Theorem 1.6, any G -orbit with noncompact stabilizer is Lorentzian. Then by Theorem 2.2, a neighborhood of some G -orbit is a warped product of the form $L \times_w S$, where S is isometric to AdS_{n+1} up to finite cover. The set of orbits having a neighborhood isometric to $L \times_w S$, for some Riemannian manifold L and $w : L \rightarrow \mathbf{R}^+$, is open. Let us prove that this set is also closed, and thus equals the whole of M . A limit O of a sequence O_k of such orbits is a non-Riemannian orbit O of dimension $\leq n+1$. Suppose that such a limit O has compact isotropy $G(x)$ for $x \in O$. Then $G(x)$ is contained in a maximal compact subgroup K of G . The Lie algebra $\mathfrak{k} \cong \mathfrak{o}(2) \times \mathfrak{o}(n)$, which has codimension $2n$ in \mathfrak{g} . Since $n \geq 3$, this is impossible. Therefore, for any $x \in O$, the stabilizer $G(x)$ is noncompact. From Theorem 1.6, O cannot be degenerate; hence, it is Lorentzian. Then by the [ArDZ] result (Theorem 2.2 above), a neighborhood of this orbit is isometric to a warped product. Any orbit of M has a neighborhood isometric to $L \times_w S$, for some L and w .

From this, one sees in particular that the G -action determines a foliation \mathcal{O} . In addition, \mathcal{O} admits an orthogonal foliation \mathcal{O}^\perp . We will use the G -action to show that the pair of foliations \mathcal{O} and \mathcal{O}^\perp arise from a global warped product of the form $L \times_w AdS_{n+1}$ on a finite cover of M .

Choose a point $x_0 \in M$. Let O_0 and O_0^\perp be the leaves of x_0 in the foliations \mathcal{O} and \mathcal{O}^\perp , respectively. Let H_0 be the stabilizer of x_0 . Note that O_0^\perp is a component of the fixed set $\text{Fix}(H_0)$; this fixed set is a closed, totally geodesic submanifold, everywhere transverse to G -orbits. It is well known that $N(H_0)/H_0$ is finite, where $N(H_0)$ is the normalizer of H_0 in G .

Let i and i^\perp denote the respective inclusions of O_0 and $\text{Fix}(H_0)$ in M . Let G act on $\text{Fix}(H_0) \times O_0$ by $g(x, y) = (x, gy)$. Define a mapping $\phi : \text{Fix}(H_0) \times O_0 \rightarrow M$, by $\phi(x, gx_0) = g(i^\perp(x))$. One sees that ϕ is well defined, and it is in fact the G -equivariant extension of the inclusions.

Next, ϕ is a covering map. Clearly ϕ is a local diffeomorphism. It is also easy to see that ϕ is surjective: the orbit of any $y \in M$ is homothetic to S . Let H_y be its stabilizer. There is some $g \in G$ conjugating H_y to H_0 . Then $gy \in \text{Fix}(H_0)$, and $y = \phi(gy, g^{-1}x_0)$. Finally, ϕ is everywhere N -to-1, where $N = |N(H_0)/H_0|$, because every G -orbit in M is homothetic to S . An N -to-1 surjective local diffeomorphism is a covering map.

The submanifold $\text{Fix}(H_0)$ is Riemannian. Let $L = \text{Fix}(H_0)$ with the metric pulled back by i^\perp . By G -equivariance of ϕ , all leaves $\phi(L \times \{y\})$ are isometric. Also by G -equivariance, the metrics along G -orbits $\phi(\{y\} \times S)$ are all homothetic, with homothety factor depending only on $y \in L$. Since S and AdS_{n+1} have a common finite cover, M is isometric, up to finite cover, to a warped product $L \times_w AdS_{n+1}$. \square

8 Full Description: Proof of Theorem 1.9

Item (1) of Theorem 1.9 follows from Theorems 1.8 and 1.7.

In this section, we consider the case in which $G_1 \cong O(1, n)$. We will take U and V to be the points of M having G_1 -orbit homothetic, up to finite cover, to dS_n and \mathbf{H}^n , respectively. The first point of item (2) follows from Theorem 1.8 and an argument just as above in section 7, to obtain the warped product on each component (see also Lemma 8.1). In the first subsection below, we will deduce the second point of (2) from Theorem 1.6. Next, we address the last point of item (2), to obtain a warped product near Riemannian G_1 -orbits, as well as the global decomposition $M = U \sqcup \partial U \sqcup V$. These two subsections will complete the proof of Theorem 1.9.

8.1 Degenerate orbits. In this subsection, we assume G is locally isomorphic to $O(1, n)$ for some $n \geq 3$, and that G acts nonproperly on M . Let U be the set of points having G -orbit homothetic, up to finite cover, to dS_n . In order to deduce the second point of (2) from Theorem 1.6, it suffices to prove the following lemma:

LEMMA 8.1. *Let $x \in \partial U$. Then the G -orbit of x is either a fixed point or degenerate with noncompact isotropy.*

Proof. The G -orbit of x cannot be Riemannian and has dimension at most n . Suppose $G(x)$ is compact, so it is contained in a maximal compact subgroup K of G . The Lie algebra $\mathfrak{k} \cong \mathfrak{o}(n)$ and has codimension n , so $\mathfrak{g}(x) \cong \mathfrak{o}(n)$; further, Gx is either Lorentzian or degenerate and n -dimensional. In either case, the isotropy representation of $\mathfrak{g}(x)$ has a 1-dimensional invariant subspace tangent to Gx and an $(n - 1)$ -dimensional complementary representation in $T_x(Gx)$, which is necessarily trivial.

On the other hand, x is the limit of a sequence $x_i \in U$ for which the isotropy $\mathfrak{g}(x_i)$ is trivial on $T(Gx_i)^\perp$, because $\mathfrak{o}(1, n - 1) \cong \mathfrak{g}(x_i)$ has no nontrivial representation in $\mathfrak{o}(d - n)$. The limit $\lim \mathfrak{g}(x_i) \subseteq \mathfrak{g}(x)$, and, because these subalgebras have the same dimension, they must be equal. Then continuity of the action implies that $\mathfrak{g}(x)$ is trivial on $T(Gx)^\perp$. Now, whether Gx is Lorentzian or degenerate, the isotropy $\mathfrak{g}(x)$ is trivial on all of $T_x M$. But isometries fixing a point and having trivial derivative at that point are trivial, so we have a contradiction.

Now $G(x)$ is noncompact. If the orbit Gx were Lorentzian, then the result of [ArDZ] (Theorem 2.2 above) would give that $x \in U$. Therefore, the G -orbit of x is fixed or degenerate. \square

8.2 Riemannian orbits. As above, we assume in this subsection that G is locally isomorphic to $O(1, n)$ for some $n \geq 3$, and that G acts nonproperly on M . By Theorem 1.8 (3), the set U defined above is open and nonempty. The following proposition implies the last point of item (2) in Theorem 1.9, as well as the claimed decomposition of M into de Sitter orbits, fixed points and light cone orbits, and hyperbolic orbits.

See Chapter 6 of [K3] for local versions of many of the results below.

PROPOSITION 8.2. *Let U and V be the points of M having orbits homothetic up to finite cover to the de Sitter space dS_n and the hyperbolic space \mathbf{H}^n , respectively. Then*

- (1) *V is open, and is G -equivariantly isometric to a warped product $L \times_w \mathbf{H}^n$, for some Lorentz manifold L .*

- (2) $\partial U = \partial V$, and it consists of all fixed points and orbits locally homothetic to the Minkowski light cone.
- (3) $M = U \sqcup \partial U \sqcup V$.

NOTATION. There exists a neighborhood of the $\mathbf{0}$ -section of TM on which the exponential map is defined and injective on each fiber. We fix one such neighborhood and denote it by Ω below.

We first collect some lemmas for the proof.

8.2.1 Maximal subalgebras of $\mathfrak{o}(1, n)$. We begin with two facts about certain natural subalgebras of $\mathfrak{o}(1, n)$. A subalgebra $\mathfrak{h} \subset \mathfrak{g}$ is maximal if it is not contained nontrivially in another subalgebra: if $\mathfrak{h} \subseteq \mathfrak{h}' \subseteq \mathfrak{g}$, then $\mathfrak{h}' = \mathfrak{h}$ or $\mathfrak{h}' = \mathfrak{g}$.

Consider the infinitesimal action of $\mathfrak{g} = \mathfrak{o}(1, n)$ on the projectivization $\mathbf{P}(\mathbf{R}^{1,n})$ of the standard representation with the standard inner product of type $(1, n)$. For $\mathbf{v} \in \mathbf{R}^{1,n}$, denote by $\bar{\mathbf{v}}$ the image in $\mathbf{P}(\mathbf{R}^{1,n})$.

- If \mathbf{v} is spacelike, then $\mathfrak{g}(\mathbf{v}) = \mathfrak{g}(\bar{\mathbf{v}})$ is conjugate to $\mathfrak{g}(\mathbf{e}_2)$, which is isomorphic to $\mathfrak{o}(1, n - 1)$, with an obvious embedding in $\mathfrak{o}(1, n)$.
- For \mathbf{v} timelike, $\mathfrak{g}(\mathbf{v}) = \mathfrak{g}(\bar{\mathbf{v}})$ is conjugate to $\mathfrak{g}(\mathbf{e}_1)$, which is isomorphic to $\mathfrak{o}(n)$, with an obvious embedding in $\mathfrak{o}(1, n)$.
- Finally, for \mathbf{v} nonzero and isotropic, $\mathfrak{g}(\bar{\mathbf{v}})$ is conjugate to $\mathfrak{g}(\overline{\mathbf{e}_1 + \mathbf{e}_2})$, which is isomorphic to the parabolic subalgebra \mathfrak{p} as in section 3. This subalgebra is isomorphic to $\mathfrak{sim}(n - 1)$, the algebra of infinitesimal affine similarities of \mathbf{R}^{n-1} . The annihilator of a nonzero isotropic \mathbf{v} is a codimension-1 ideal of $\mathfrak{g}(\bar{\mathbf{v}})$, isomorphic to $\mathfrak{euc}(n - 1)$, the algebra of infinitesimal affine isometries of \mathbf{R}^{n-1} .

LEMMA 8.3. (1) $\mathfrak{sim}(n - 1)$ is a maximal subalgebra of $\mathfrak{o}(1, n)$.

(2) $\mathfrak{euc}(n - 1)$ is contained in exactly one maximal subalgebra, $\mathfrak{sim}(n - 1)$.

(3) Let \mathfrak{h} be a compact proper subalgebra of $\mathfrak{o}(1, n)$ containing two different conjugates \mathfrak{k} and \mathfrak{k}' , each isomorphic to $\mathfrak{o}(1, n - 1)$. Then \mathfrak{h} is conjugate to $\mathfrak{o}(n)$.

Proof. (1) The subalgebra $\mathfrak{sim}(n - 1)$ acts infinitesimally conformally on the projectivization of the light cone, which is conformally equivalent to S^{n-1} . It is the infinitesimal stabilizer of one point and acts transitively on the complement of this point. Then, for any $X \notin \mathfrak{sim}(n - 1)$, the algebra generated by X and $\mathfrak{sim}(n - 1)$ is transitive on S^{n-1} and contains the full stabilizer subalgebra of each point. It follows that any subalgebra properly containing $\mathfrak{sim}(n - 1)$ is $\mathfrak{o}(1, n) \cong \mathfrak{conf}(S^{n-1})$, so $\mathfrak{sim}(n - 1)$ is a maximal subalgebra.

(2) Let \mathfrak{h} be a maximal subalgebra of $\mathfrak{o}(1, n)$ containing $\mathfrak{euc}(n-1)$, and suppose that \mathfrak{h} does not preserve any isotropic line in $\mathbf{R}^{1, n}$. Because the corresponding connected group $H \subset O(1, n)$ is not compact, Theorem 1.1 of [BoZ] implies that \mathfrak{h} preserves some Lorentz subspace of $\mathbf{R}^{1, n}$ and contains the full infinitesimal linear isometry algebra of this subspace. Because $\mathfrak{euc}(n-1)$ preserves no proper Lorentz subspace, \mathfrak{h} must equal $\mathfrak{o}(1, n)$.

(3) Let H be the connected subgroup of $O(1, n)$ with Lie algebra \mathfrak{h} . If H is compact, it is conjugate to a subgroup of $O(n)$ containing two copies of $SO(n-1)$. By an argument similar to that for $\mathfrak{sim}(n-1)$, the subalgebra $\mathfrak{o}(n-1)$ is maximal in $\mathfrak{o}(n)$, so $\mathfrak{h} = \mathfrak{o}(n)$. \square

8.2.2 Fixed point sets. Recall that, for H a subgroup of G , the fixed set is denoted $\text{Fix}(H)$. Recall also that the exponential map is defined on Ω and injective on each fiber in it. Each component of $\text{Fix}(H)$ is a totally geodesic submanifold of M . Let D be the points of M having degenerate G -orbit with noncompact isotropy. By Theorem 1.6, any $x \in D$ has $\mathfrak{g}(x) \cong \mathfrak{o}(n-1) \times \mathbf{R}^{n-1} \cong \mathfrak{euc}(n-1)$.

LEMMA 8.4. *Let $x \in D$, and denote by O its G -orbit. Near x , the fixed set $\text{Fix}(G(x))$ coincides with $\exp_x(T_x O^\perp \cap \Omega)$. It has dimension $d-n$, and intersects O along the isotropic geodesic in O through x .*

Proof. By Proposition 5.2, we can approximate the degenerate orbit O by Lorentzian orbits O_i with noncompact isotropy. By Theorem 2.2, each O_i is homothetic, up to finite cover, to dS_n . Take $x_i \in O_i$ with $x_i \rightarrow x \in O$. The limit $\lim \mathfrak{g}(x_i)$ is contained in $\mathfrak{g}(x) \cong \mathfrak{euc}(n-1)$; since $\dim(\mathfrak{g}(x_i)) = \dim(\mathfrak{g}(x))$ for all i , the limit equals $\mathfrak{g}(x)$. As in the proof of Lemma 8.1, triviality of $\mathfrak{g}(x_i)$ on $(T_{x_i} O_i)^\perp$ for all i implies that, in the limit, the isotropy $\mathfrak{g}(x)$ is trivial on $(T_x O)^\perp$. It is easy to see from the form of the isotropy representation of $\mathfrak{g}(x) \cong \mathfrak{euc}(n-1)$ that the maximal trivial subrepresentation in $T_x M$ is exactly $(T_x O)^\perp$. Then the lemma follows with the exponential map and dimension counting. \square

LEMMA 8.5. *Let K be a maximal compact connected subgroup of $G(x)$. Then $\text{Fix}(K)$ has dimension $d-n+1$ and is of Lorentzian type. It meets O along the isotropic geodesic in O through x .*

Proof. The subgroup K is conjugate to $SO(n-1) \subset Euc(n-1)$. The isotropy representation of K at x fixes any nonzero normal vector $n_x \in (T_x O)^\perp \cap T_x O$ and acts irreducibly on an $(n-1)$ -dimensional complementary spacelike subspace $L_x \subset T_x O$. Then $\text{Fix}(K) \cap O = N_x$, where N_x is the

isotropic geodesic in O through x . Now, $L_x \oplus (T_x O)^\perp$ has codimension 1 in $T_x M$. There is a K -invariant 1-dimensional complementary representation, which is necessarily trivial. We have therefore a $(d - n + 1)$ -dimensional Lorentzian subspace L'_x containing $(T_x O)^\perp$, complementary to L_x , on which K acts trivially. This subspace is exactly the maximal trivial subrepresentation of K on $T_x M$. \square

LEMMA 8.6. *Let*

$$X_x = \cup \{T_x(\text{Fix}(K)) \setminus \{0\} : K \subset G(x) \text{ maximal compact connected}\}.$$

Then X_x is open in $T_x M$. More precisely,

$$X_x = ((T_x O)^\perp \times P^c) \cup ((T_x O)^\perp \setminus \{0\}),$$

where P is a hyperplane in a subspace F complementary to $(T_x O)^\perp$.

Proof. Fix one maximal connected compact $K_0 \subset G(x)$, locally isomorphic to $SO(n - 1)$. Let F_0 be a K_0 -invariant degenerate complement to $(T_x O)^\perp$, and let \mathbf{v}_0 be a nonzero isotropic vector in F_0 . Then $(T_x O)^\perp + \mathbf{R}\mathbf{v}_0 = T_x(\text{Fix}(K_0))$. Choose a generator \mathbf{n}_x of $(T_x O)^\perp \cap T_x O$; note that \mathbf{n}_x is fixed by $G^0(x)$. Now any other maximal compact, connected subgroup of $G(x)$ equals gK_0g^{-1} for some $g \in G^0(x)$. The fixed subspace $T_x(\text{Fix}(gK_0g^{-1})) = (T_x O)^\perp + \mathbf{R}g\mathbf{v}_0$. As g ranges over $G^0(x)$, the projection of $g\mathbf{v}_0$ to F_0 ranges over all vectors \mathbf{v} with $\langle \mathbf{v}, \mathbf{n}_x \rangle = \langle \mathbf{v}_0, \mathbf{n}_x \rangle$. Then the projection of $\mathbf{R}G^0(x)\mathbf{v}_0$ to F_0 ranges over all \mathbf{v} with $\langle \mathbf{v}, \mathbf{n}_x \rangle \neq 0$. Now the lemma is proved, with $F = F_0$ and $P = F_0 \cap \mathbf{n}_x^\perp$. \square

8.2.3 Hypersurface of degenerate orbits. Let O be as above, a degenerate orbit with isotropy $\mathfrak{euc}(n - 1)$. We will next show that any neighborhood of a point of O meets U or V . Then we will show that O lies in a hypersurface of degenerate orbits with the same isotropy, and, finally, that this hypersurface locally separates U from V .

Let X_x be as in Lemma 8.6 above. Consider $X(O) = \cup \{X_x \cap \Omega : x \in O\}$. This is an open set in the restriction $TM|_O$.

LEMMA 8.7. *The image $\exp(X(O)) \subset U \cup D \cup V$ – that is, for any $y \in \exp(X(O))$, the stabilizer $\mathfrak{g}(y) \cong \mathfrak{o}(n), \mathfrak{euc}(n - 1),$ or $\mathfrak{o}(1, n - 1)$.*

Proof. Denote by Φ the restriction of the exponential map $\Phi = \exp : X(O) \rightarrow M$. Reducing Ω if necessary, we may assume $\Phi(X(O))$ contains no fixed points. Each X_x is open in $T_x M$, so Φ has maximal rank, and is in particular an open map. Its image is an open set containing O in its closure. Any point y in this image is a regular value; the inverse image $S = \Phi^{-1}(y) \subset X(O)$ has dimension $\dim(X(O)) - \dim(M) = n$. Because \exp is

injective on each fiber of Ω , and hence of $X(O)$, the inverse image S projects injectively under $\pi : TM \rightarrow M$ onto some open set W in O . For any $z \in W$, there exists a maximal compact $K_z \subset G(z)$ such that $y \in \text{Fix}(K_z)$. Because $n \geq 3$, there exist $z, z' \in W$ which do not lie on a common isotropic geodesic of O , so that K_z and $K_{z'}$ are contained in $G(y)$ and are distinct. By Lemma 8.3 (3), either $\mathfrak{g}(y) \cong \mathfrak{o}(n)$, or it is noncompact. In the latter case, by [ArDZ] and Theorem 1.6, $\mathfrak{g}(y) \cong \mathfrak{o}(1, n - 1)$ or $\mathfrak{euc}(n - 1)$. \square

Let $x \in O$, and let R be a neighborhood of x in $\text{Fix}(G(x))$. Let \widehat{O} be the union of all the G -orbits of all points of R .

LEMMA 8.8. *For R sufficiently small, \widehat{O} is a degenerate hypersurface in which every orbit is degenerate with isotropy $\mathfrak{euc}(n - 1)$; further, $\widehat{O} \cup U \cup V$ is a neighborhood of \widehat{O} .*

Proof. From Lemma 8.4, R equals $\exp((T_x O)^\perp \cap \Omega)$ near x and has dimension $d - n$. Since $\mathfrak{g}(x) \cong \mathfrak{euc}(n - 1)$ lies in a unique maximal subalgebra of $\mathfrak{o}(1, n)$ by Lemma 8.3 (2), the orbit of any $z \in R$ is either a fixed point or degenerate. Shrink R so that the G -orbit of any $z \in R$ is degenerate; in this case, $\mathfrak{g}(z) \cong \mathfrak{euc}(n - 1)$. The tangent space $T_z \widehat{O} = T_z(Gz) + T_z(\text{Fix}(G(x))) = T_z(Gz) + (T_z(Gz))^\perp$, because z is also as in Lemma 8.4; hence, \widehat{O} is a degenerate hypersurface.

Now consider $X(\widehat{O}) = \cup\{X(Gz) : z \in R\} \subset T\widehat{O}$, and let Φ be the restriction of \exp to $X(\widehat{O})$. From Lemma 8.7, the image of the restriction of Φ_z to $X(Gz)$ is open and contained in $U \cup D \cup V$, and its closure is a neighborhood of Gz . If $y \in D$, then $\mathfrak{g}(y)$ is conjugate to $\mathfrak{euc}(n - 1)$. But y is close to a point of \widehat{O} , with stabilizer also conjugate to $\mathfrak{euc}(n - 1)$. Then y has the same stabilizer as a nearby point of \widehat{O} . The set Ω can be chosen sufficiently small that any such y is contained in \widehat{O} . Then the union

$$\bigcup_{z \in R} \Phi_z(X(Gz))$$

is contained in $U \cup \widehat{O} \cup V$, and its closure is a neighborhood of \widehat{O} . The subset of points of this closure with $\dim(\mathfrak{g}(y)) \leq \dim \mathfrak{euc}(n - 1)$ still forms a neighborhood of \widehat{O} , and is contained in $U \cup \widehat{O} \cup V$. \square

LEMMA 8.9. *Let Φ be the restriction of the exponential map to $X(\widehat{O})$, in a sufficiently small neighborhood of the $\mathbf{0}$ -section. One component of $\Phi(X(\widehat{O})) \setminus \widehat{O}$ lies in U , and the other is in V .*

Proof. From the last lemma, any point $x \in \widehat{O}$ has a neighborhood in which $\dim(\mathfrak{g}(x)) = \dim(\mathfrak{g}(y))$ for all y in the neighborhood; let N be this common dimension. In this neighborhood, the map $y \mapsto \mathfrak{g}(y) \in Gr^N(\mathfrak{g})$ is

continuous. There are smaller neighborhoods $Z \subset W$ of x such that W is a normal neighborhood of each of its points, and, whenever $z, z' \in Z$ are such that $\mathfrak{g}(z) = \text{Ad}(g)(\mathfrak{g}(z'))$, then $gz' \in W$. (*Note:* We take g to be the element of $\text{Ad}^{-1}(\text{Ad}(g))$ closest to the identity.)

Suppose for a contradiction that points on opposite sides of \widehat{O} have the same orbit type. Let Z_1, Z_2 be the components of $Z \setminus \widehat{O}$, and W_1, W_2 the components of $W \setminus \widehat{O}$. Take $z \in Z_1$ and $z' \in Z_2$. By assumption, $\mathfrak{g}(z)$ and $\mathfrak{g}(z')$ are conjugate in G . Let g be a conjugating element with minimal distance to the identity. Then $z, gz' \in W$ are connected by a unique geodesic, which necessarily passes through \widehat{O} . The common stabilizer $\mathfrak{g}(z) = \mathfrak{g}(gz')$ fixes the geodesic pointwise, so it fixes a point of \widehat{O} . Whether $\mathfrak{g}(z)$ is isomorphic to $\mathfrak{o}(n)$ or to $\mathfrak{o}(1, n - 1)$, neither embeds in the stabilizer of a point of \widehat{O} . □

8.2.4 Proof of Proposition 8.2. Suppose that $U \neq M$, so $\partial U \neq \emptyset$. By Lemma 8.9, not only is V nonempty, but the interior $\text{int}(V) \neq \emptyset$. The following fact will be key for the rest of the proof.

Fact 8.10. *For $x \in \text{int}(V)$ with orbit O , the isotropy representation of $\mathfrak{g}(x)$ is trivial on $(T_x O)^\perp$.*

Proof. The stabilizer $\mathfrak{g}(x) \cong \mathfrak{o}(n)$ acts on the orthogonal space $(T_x O)^\perp$ via a homomorphism $\mathfrak{o}(n) \rightarrow \mathfrak{o}(1, d - n - 1)$. There is a neighborhood W of $(x, \mathbf{0})$ in $(TO)^\perp$ on which the exponential map is a diffeomorphism onto its image. For any $y \in \exp_x(W \cap T_x M)$, the stabilizer $\mathfrak{g}(y) \subseteq \mathfrak{g}(x)$. If all orbits near O have the same dimension, then $\mathfrak{g}(x) = \mathfrak{g}(y)$ for all $y \in \exp(W \cap T_x M)$. Then the orthogonal representation of $\mathfrak{g}(x)$ is trivial. □

Now we can show that $\partial(\text{int}(V)) = \partial U$. Suppose $x_n \in \text{int}(V)$ and $x_n \rightarrow x$. Denote by O the orbit of x . As usual, we have $\lim \mathfrak{g}(x_n) \subseteq \mathfrak{g}(x)$, and $\dim(O) \leq n$. The orbit of x is either Riemannian, degenerate, or fixed.

First suppose O is Riemannian. Then $\mathfrak{g}(x)$ is compact, so it is conjugate to a subalgebra of $\mathfrak{o}(n)$. The dimension restriction forces $\mathfrak{g}(x)$ to be conjugate to $\mathfrak{o}(n)$, so $x \in V$. As in the proof of 8.1, the isotropy $\mathfrak{g}(x)$ is trivial on $(T_x O)^\perp$, so orbits near x have isotropy containing $\mathfrak{o}(n)$. The orbit O has a neighborhood consisting of Riemannian G -orbits. Because $\mathfrak{o}(n)$ is a maximal subalgebra, orbits near x all have isotropy $\mathfrak{g}(x)$. Then $x \in \text{int}(V)$, a contradiction.

If O is degenerate, the same argument as in Lemma 8.1 implies that $\mathfrak{g}(x)$ is noncompact. Then Proposition 5.2 implies that $x \in \partial U$. If x is a fixed point, then Proposition 5.1 implies that $x \in \partial U$. Now Lemma 8.1 gives item (2).

Now $U \cup \partial U \cup \text{int}(V)$ is closed. It is open by Lemma 8.8, so it equals M . Therefore, $\text{int}(V) = V$, and $M = U \sqcup \partial U \sqcup V$, proving item (3).

It remains to prove item (1), the warped product structure in V . As in the proof in [ArDZ, §2.3] of warped products around Lorentz orbits and as in section 7, the key properties are irreducibility of the isotropy along orbits and triviality of the isotropy orthogonal to orbits.

Fix $x \in V = \text{int}(V)$ with G -orbit O , and let $\mathcal{L}_x = \exp_x((T_x O)^\perp \cap \Omega)$. Since $G(x)$ is irreducible on $T_x O$ and trivial on $(T_x O)^\perp$, the leaf \mathcal{L}_x is exactly the set of common fixed points of $G(x)$ near x . Let $L = \text{Fix}(G(x)) \cap V$. Define $\phi : L \times O \rightarrow V$ by $\phi(y, gx) = gy$. This map is a local diffeomorphism because, for any $y \in \text{Fix}(G(x)) \cap V$, the fixed set L coincides near y with $\exp((T_y(Gy))^\perp \cap \Omega)$. If $\phi(y, gx) = \phi(y', g'x)$, then $g^{-1}g' \in N_G(G(x))$, and $g^{-1}g'y' = y$. The stabilizer $G(x)$ is a maximal compact subgroup of the semisimple connected group G , so $N_G(G(x)) = G(x)$ (see [E, 1.13.14 (4)]). It follows that ϕ is injective. Any $y \in V$ has $G(y)$ compact and $\dim G(y) = \dim G(x)$; because $G(x)$ is connected, there exists $g \in G$ such that $gG(y)g^{-1} = G(x)$. Then $\phi(gy, g^{-1}x) = y$, so ϕ is surjective. In conclusion, ϕ is a G -equivariant diffeomorphism $L \times O \rightarrow V$.

The orbit O is assumed to be homothetic, up to finite cover, to \mathbf{H}^n . Because \mathbf{H}^n is simply connected, it has no finite covers. Because any finite subgroup of $\text{Isom}(\mathbf{H}^n)$ has a common fixed point, there are no smooth finite quotients of \mathbf{H}^n . Therefore, the orbit O , and every other orbit of V , is globally homothetic to \mathbf{H}^n . The metric on V pulls back by ϕ to $L \times O$. The transverse fibers of the product are orthogonal by construction. All L -fibers are isometric because they are related by the action of G . Also by equivariance, O -fibers in $L \times O$ correspond to G -orbits in V , so each is homothetic to \mathbf{H}^n . In conclusion, there exists $w : L \rightarrow \mathbf{R}^+$ such that V is isometric to $L \times_w \mathbf{H}^n$. \square

The referee kindly communicated to us the following lemma. It synthesizes arguments that we used in many proofs, especially in the proof of Proposition 8.2 above. Actually, this proof above includes the proof of the lemma.

LEMMA 8.11 (The anonymous referee). *Let G act by isometries of a pseudo-Riemannian manifold M with no isotropic orbits. Suppose that, for all $x \in M$, the identity component $G^0(x)$ is conjugate to S , for S a fixed closed, connected subgroup of G . Assume that the adjoint representation of \mathfrak{s} on $\mathfrak{g}/\mathfrak{s}$ is absolutely irreducible. Let L be the set of S -fixed points in M . Then the G -action lifts to an N -to-1 cover M' of M , where $N = [N_G(S) : S]$.*

Further, M' is G -equivariantly isometric to a warped product $L \times_w G/S$, where G/S carries a nondegenerate G -invariant pseudo-Riemannian metric, and L carries the metric induced from M .

The hypothesis that all stabilizers are conjugate in G means that orbits are all isomorphic as homogeneous G -spaces. Together with the irreducibility hypotheses, one obtains that the set of points having a given stabilizer S' is transverse to the orbit foliation, and in fact equals $\text{Fix}(S')$. This situation is exactly that of the proof above. Not only in the proof of (1) in Proposition 8.2, but also in section 7, as part of the proof of Theorem 1.7, we prove a case of this lemma. Note that the hypothesis that all stabilizers are conjugate is strong, and, in each of our two cases, fulfilling the hypotheses of the lemma is intertwined with the rest of the proof.

Acknowledgements. We thank the referee for his many thoughtful comments and helpful suggestions. The first two authors thank the École Normale Supérieure de Lyon for their hospitality. During their stay, they received assistance from the ACI *Structures géométriques et trous noirs*.

References

- [A1] S. ADAMS, Orbit nonproper actions on Lorentz manifolds, *GAFA*, *Geom. Funct. Anal.* 11:2 (2001), 201–243.
- [A2] S. ADAMS, Orbit nonproper dynamics on Lorentz manifolds, *Illinois J. Math.* 45:4 (2001), 1191–1245.
- [A3] S. ADAMS, Dynamics of semisimple Lie groups on Lorentz manifolds, *Geom. Ded.* 105 (2004), 1–12.
- [A4] S. ADAMS, Induction of geometric actions, *Geom. Ded.* 88 (2001), 91–112.
- [AS1] S. ADAMS, G. STUCK, The isometry group of a compact Lorentz manifold I, *Inv. Math.* 129:2(1997), 239–261.
- [AS2] S. ADAMS, G. STUCK, The isometry group of a compact Lorentz manifold II, *Inv. Math.* 129:2 (1997), 263–287.
- [ArDZ] A. AROUCHE, M. DEFFAF, A. ZEGHIB, On Lorentz dynamics: From group actions to warped products via homogeneous spaces, *Trans. Amer. Math. Soc.* 359 (2007), 1253–1263.
- [BN] U. BADER, A. NEVO, Conformal actions of simple Lie groups on compact pseudo-Riemannian manifolds, *J. Differential Geom.* 60:3 (2002), 355–387.
- [BaZ] T. BARBOT, A. ZEGHIB, Group actions on Lorentz spaces, mathematical aspects: a survey, in “The Einstein Equations and the Large-Scale Behavior of Gravitational Fields” (P. Chrusciel, H. Friedrich, eds.) Birkhäuser, Basel (2004), 401–439.
- [BoZ] C. BOUBEL, A. ZEGHIB, Isometric actions of Lie subgroups of the Moebius group, *Nonlinearity* 17 (2004), 1677–1688.

- [DG] G. D'AMBRA, M. GROMOV, Lectures on transformation groups: geometry and dynamics, in "Surveys in Differential Geometry (Cambridge, MA, 1990)", Lehigh Univ., Bethlehem, PA (1991), 19–111.
- [E] P.B. EBERLEIN, Geometry of Nonpositively Curved Manifolds University of Chicago Press, Chicago, 1996.
- [G] M. GROMOV, Rigid transformations groups, in "Géométrie Différentielle" (D. Bernard, Y. Choquet-Bruhat, eds.) Hermann, Paris (1988), 65–139.
- [H] S. HELGASON, Differential Geometry, Lie Groups, and Symmetric Spaces, Academic Press, New York, 1978.
- [K1] N. KOWALSKY, Noncompact simple automorphism groups of Lorentz manifolds, *Ann. Math.* 144 (1997), 611–640.
- [K2] N. KOWALSKY, Actions of noncompact simple groups on Lorentz manifolds, *C.R. Acad. Sci. Paris Sér. I Math.* 321:5 (1995), 595–599.
- [K3] N. KOWALSKY, Actions of noncompact simple groups on Lorentz manifolds and other geometric manifolds, Thesis, University of Chicago, 1994.
- [O] B. O'NEILL, Semi-Riemannian Geometry, Academic Press, San Diego, 1983.
- [S] H. SAMELSON, Notes on Lie Algebras, van Nostrand, New York, 1969.
- [W] D. WITTE, Homogeneous Lorentz manifolds with simple isometry group, *Beiträge Alg. Geom.* 42:2 (2001) 451–461.
- [Z1] A. ZEGHIB, Geodesic foliations in Lorentz 3-manifolds, *Comm. Math. Helv.* 74 (1999), 1–21.
- [Z2] A. ZEGHIB, Remarks on Lorentz symmetric spaces, *Comp. Math.* 140 (2004), 1675–1678.
- [Z3] A. ZEGHIB, Sur les espaces-temps homogènes, The Epstein birthday schrift, *Geom. Topol. Publ. Coventry* (1998), 551–576.
- [Z4] A. ZEGHIB, The identity component of the isometry group of a compact Lorentz manifold, *Duke Math. J.* 92:2 (1998), 321–333.
- [Zi1] R.J. ZIMMER, Automorphism groups and fundamental groups of geometric manifolds, *Proc. Symp. Pure Math.* 54:3 (1993), 693–710.
- [Zi2] R.J. ZIMMER, On the automorphism group of a compact Lorentz manifold and other geometric manifolds, *Invent. Math.* 83:3 (1986), 411–424.

MOHAMED DEFFAF, Faculté des mathématiques, USTHB, BP 32 El'Alia, Bab Ezzouar, Alger, Algeria mhdeffaf@gmail.com

KARIN MELNICK, Department of Mathematics, Yale University, PO Box 208283, New Haven, CT, USA karin.melnick@yale.edu
www.math.yale.edu/~km494/

ABDELGHANI ZEGHIB, CNRS, UMPA, École Normale Supérieure de Lyon, 46, allée d'Italie, 69364 Lyon cedex 07, France Zeghib@umpa.ens-lyon.fr
www.umpa.ens-lyon.fr/~zeghib/

Received: May 2006

Revision: February 2007

Accepted: March 2007