



Coupling 1D modifications and 3D nuclear organization: data, models and function

Daniel Jost¹, Cédric Vaillant² and Peter Meister³

Over the past decade, advances in molecular methods have strikingly improved the resolution at which nuclear genome folding can be analyzed. This revealed a wealth of conserved features organizing the one dimensional DNA molecule into tridimensional nuclear domains. In this review, we briefly summarize the main findings and highlight how models based on polymer physics shed light on the principles underlying the formation of these domains. Finally, we discuss the mechanistic similarities allowing self-organization of these structures and the functional importance of these in the maintenance of transcriptional programs.

Addresses

¹ University Grenoble Alpes, CNRS, TIMC-IMAG lab, UMR 5525, Grenoble, F-38706 La Tronche, France

² Univ Lyon, ENS de Lyon, Univ Claude Bernard, CNRS, Laboratoire de Physique, F-69007 Lyon, France

³ Cell Fate and Nuclear Organization, Institute of Cell Biology, University of Bern, CH-3012 Bern, Switzerland

Corresponding author: Meister, Peter (peter.meister@izb.unibe.ch)

Current Opinion in Cell Biology 2017, **44**:20–27

This review comes from a themed issue on **Cell Architecture**

Edited by **Terry Lechler** and **Rafael Carazo Salas**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 29th December 2016

<http://dx.doi.org/10.1016/j.ceb.2016.12.001>

0955-0674/© 2016 Elsevier Ltd. All rights reserved.

Chromosome conformation capture and superresolution microscopy: 3D compartments in the nucleus


The past decade has seen major progresses in our understanding of chromosome organization in interphase nuclei with chromosome conformation capture techniques (3C) [1,2], supported by parallel superresolution microscopy studies [3,4[•],5[•],6]. 3C was designed to characterize long-range chromatin contacts based on chromatin crosslinking, restriction digestion of crosslinked complexes and ligation of digested fragments in their crosslinked chromatin context. Ligation events between restriction fragments not immediately adjacent along the linear genome imply that these fragments were present in the same crosslinked complex. While the original procedure [1] allowed the characterization of contacts between two target loci, later development (termed 4C, 5C, Hi-C)

coupled to high throughput sequencing allowed the identification of two-ways contacts of one, many or all fragments of the genome, respectively (for an in-depth technical review, see [2]).

In particular, genome-wide (Hi-C) maps provided a global view of possible contacts between any DNA segments of the genome [7]. Hi-C maps show a characteristic high contact frequency along the diagonal: sequences close to each other along the linear genome have a higher probability of contact. This mainly depends on the distance in base pairs between fragments, a property that can be used to build *de novo* chromosome backbones [8]. At the megabasepair (Mbp) scale, contact maps display a cell-type specific checker board pattern [4[•],7,9] (Figure 1a,b). This demonstrates the presence of two mutually exclusive contact compartments, named A and B [7,9] and the checker board pattern implies that sequences in the same compartment (A or B) contact each other more frequently than sequences in different ones (A-B). Strikingly, sequences in A compartments are gene rich, display higher expression than B sequences and A chromatin is generally more accessible and enriched for histone marks specific for active genes. In contrast, B compartments are more densely packed, harbor less genes and more repressed histone marks [7,9,10]. Together, these results demonstrate clustering of functionally similar chromatin, highly reminiscent of previous microscopy and biochemical studies in which spatial segregation between euchromatin (transcriptionally more active) and heterochromatin (transcriptionally more silent) was observed [11].

At higher resolution, Hi-C and 5C experiments highlighted compartmentalization into kbp- to Mbp sized contact domains [3,10], the so-called topologically-associating domains (TAD) (Figure 1b). TADs are defined as highly self-contacting portions of the genome: a sequence inside a TAD has a higher probability to contact sequences inside the same TAD than sequences in neighboring TADs [10]. TADs insulate sequences along the linear genome, thereby segmenting the 1D genome into 3D domains. The presence of TADs or similar 3D domains has been documented in most species in which genome-wide 3C has been carried out ranging from bacteria to mammalian cells and plants [12]. Most characterized promoter-enhancer interactions occur inside the same TAD [10,13[•]], suggesting that TADs allow insulation of promoters from enhancers located in neighboring TADs. Indeed, deletion of a TAD boundary is sufficient to induce ectopic new contacts with regulatory sequences

Figure 1



Contact maps in human cells. **(a)** Contact frequency map across one third of the right arm of human chromosome 7 in two different cell lines, based on Hi-C data [4**], at 50 kb bin size. The checkerboard contact frequency pattern is cell-type specific and allows the definition of two compartments A and B (middle bars). **(b)** A and B compartments in various cell lines (data from [4**]). **(c)** Conservation of Topologically Associating Domains (TADs) in two cell lines. Contact frequencies are drawn for a bin size of 5 kb. Arrows: 'corner' contacts, in which loci frequently contact each other, as the result of the creation of a loop. These loci are located at the base of the loops.

in the neighboring TAD and leads to aberrant gene expression even if not disrupting any regulatory segment [14]. In humans and mouse, this kind of deletion and subsequent gene misregulation leads to limb malformations such as polydactyly, finger malformations or sex reversal [15**,16]. The TAD structure of the genome is remarkably invariant over the course of development, between different cell types (Figure 1c) or even between species in conserved syntenic blocks [17**,18–20].


One of the key questions about both A/B compartments and TADs is how these domains are built. Several models based on polymer physics were suggested in the past

years, providing an interesting starting point to understand the minimal requirements for the creation of such higher-order chromatin structures.

Modeling chromatin as a polymer

Initial theoretical polymer models of chromosomes considered chromatin as a homogeneous polymer (Figure 2a). In those models, chromatin is described by a chain of connected identical monomers whose dynamics is controlled by thermal forces, excluded volume interactions (monomers cannot overlap) and optionally by the rigidity of the chromatin fiber or by non-specific dynamic interactions between monomers. Such homopolymer models

Figure 2



Polymer models of chromatin folding. **(a)** Homopolymer model: chromatin is constituted by a chain of connected monomers, each monomer representing a stretch of chromatin. Dynamics is controlled by thermal motion (i) and by steric interactions as two beads cannot overlap or cross each other (ii). **(b)** Copolymer model: a state is attributed to each monomer, depending on its underlying genomic or epigenomic information. Transient contacts may occur between loci of the same state (i, ii) or between a locus and the nuclear membrane (not shown). Long blocks of monomers having the same state will collapse into a TAD or may engage long-range interactions with distant regions. Alternative copolymer models were built with the assumption that dispersed monomers have increased affinities, which may lead to loop and TAD formation. **(c)** Loop extrusion model: extruders bind to chromatin (i), actively extrude DNA (ii), until they eventually detach from chromatin (iii). Sliding of extruders along chromatin is stopped or paused when encountering boundary elements in the proper orientation (iv), such as CTCF-bound sites in mammalian cells. This leads to the formation of TADs between convergent sites and boundaries between closely located divergent ones.

are sufficient to provide a quantitative and predictive description of chromatin organization in organisms with small genomes and relatively simple nuclear organization such as budding yeast [21–23]. Additionally, these models are able to predict generic aspects of chromosome folding in higher organisms, such as the average evolution of contact frequencies or distances between two loci as a function of their distance in base pairs along the linear genome, as well as chromosome intermingling [24–27]. In particular, large-scale organization of the genome is compatible with unknotted self-similar polymeric conformations, the so-called fractal or crumpled globule model [7,28,29], that are likely to result from the out-of-equilibrium folding of long polymers into a confined space [24]. However, these models fail to describe higher order structures such as A/B compartments or TADs.

Experimental data revealed that TAD epigenomic composition is often uniform for either active or inactive epigenetic marks [4^{••},9,20,30] while TAD boundaries are enriched for a number of chromatin binding proteins

or specific epigenomic states [4^{••},9,10,30–32]. This suggested that the 1D heterogeneous chromatin landscape along the genome plays a key role in the 3D folding. Motivated by these observations, heteropolymer models started recently to emerge, based on the homopolymer framework but explicitly considering the coupling between chromatin structure and function [33–38,39^{••},40,41,42^{••},43,44^{••},45].

The first major class of models (Figure 2b) considers that chromatin folding is driven by transient specific short-range or contact interactions between genomic loci or with nuclear landmarks such as the nuclear periphery. Mainly two types of polymers were considered: either (co-)polymers where large stretches of monomers have similar short-range interaction properties (Figure 2b, self-interacting) or where individual dispersed monomers have an affinity for each other, sometimes dependent on *trans* acting factors (Figure 2b, dispersed). TAD conformations vary from dense globule to more disordered structures, depending on the strength of interactions, the

size of the domain and the concentration of the ‘bridging’ molecules, as observed *in vivo* [33,39^{••},46–48]. In these frameworks, TAD formation results from an internal collapse due to attractive forces between the monomers composing a TAD. These approaches are experimentally supported by evidence that chromatin-binding proteins (transcription factors, transcription and replication machineries, lamins, cohesins, condensins, heterochromatin-associated complexes such as Polycomb or HP1, and so on) promote directly or effectively physical bridging [49–51]. In *Drosophila*, such heteropolymer models built directly from the 1D epigenomic landscape account well for the checker board patterns observed in Hi-C experiments (Figures 1a and 2b long range contacts) [39^{••},43,47], as well as the formation and properties of previously observed Polycomb bodies where silent H3K27me-enriched chromatin clusters [5[•],52[•]]. The prediction of long-range interactions is inherent to copolymer models arranged in blocks. Therefore, such models are well adapted to describe A/B compartment formation in mammals [53]. Interestingly, the temporal evolution which can be extracted from the models suggests that the internal collapse of epigenomic domains into TADs is a fast process while long-range contacts between TADs of the same epigenomic state are more slowly formed, more transient and dynamic [34,39^{••},47]. These predictions are consistent with Hi-C experiments in mammalian cells on synchronized or single cells [54,55]. Short-range interactions such as the ones used in heteropolymer models are however inconsistent with the observation that half of mammalian TADs contains strong loops between convergent CTCF (CCCTC-binding factor) sequence motifs, usually located at domain boundaries [4^{••},56,57^{••}]. While the formation of these loops can still be explained using dispersed models [34,42^{••}], this first class of models is however inconsistent with the observed preferential looping between convergent CTCF sites and with results obtained by experiments in which boundaries were deleted or inverted [42^{••},44^{••}].


To account for that, a second class of models has recently emerged (Figure 2c), based on early speculations on mitotic condensation [58,59]. These models consider the formation of TADs as the results of an active extrusion mechanism [42^{••},44^{••}]. Protein complexes, putatively structural maintenance of chromosome complexes (SMC; cohesins and condensins) bind to chromatin and extrude dynamically large chromatin loops. Eventually the complexes detach or stop extruding chromatin as they encounter boundary elements. In these models, the local 3D organization of the genome is controlled by the presence and orientation of proteins bound to 1D sequence elements that serve as barriers. Polymer models implementing this mechanism reproduce TAD formation as well as the observed loops between TAD boundaries [42^{••},44^{••}]. These models are experimentally supported in mammalian cells by the enrichment for cohesins

(a possible loop extrusion factor), CTCF and its cognate sequence motif at TAD boundaries, by the high contact frequencies of the sequences at the TAD boundaries (‘corners’ of the domains, Figure 1b, arrows), as well as by the preferential contacts between convergent CTCF binding sites [4^{••},10,42^{••},44^{••},57^{••}]. Moreover, these models can quantitatively predict the perturbed 3D organization after deletion, inversion or duplication of CTCF sites [14,42^{••}]. Similarly, TADs on the dosage compensated *C. elegans* X chromosome depend on sequence elements (*rex* sites) which are enriched at TAD boundaries, while the formation of these X-specific TADs is achieved by the loading of a condensin-like loop extrusion factor [60]. The loop extrusion models additionally suggest an elegant mechanism for the formation of mitotic chromosomes: the increase in the number of loop extruders coupled to a decrease in the number of boundary elements leads to higher compaction and an array of consecutive loops [61,62] in agreement with Hi-C data obtained for mitotic chromosomes [54]. Loop extrusion models reach their limits when trying to predict long-range (inter-TADs) contacts such as the formation of A/B compartments, or interactions with the nuclear envelope. As these processes are likely driven by epigenomic information, heteropolymer models accounting for both loop extrusion and epigenomically-driven short-range interactions remain to be developed in order to quantitatively describe within the same framework the local and higher-order chromosome organization.

Spatial chromatin compartments: ‘nano-reactors’ for gene regulation

The main question raised by the presence of nuclear compartments at different scales (A/B compartments, TADs, loops and so on) is whether this spatial organization of the genome is a by-product of genome activity or whether it is participating in the regulation of genome function. A popular hypothesis is that molecular crowding and spatial confinement increase binding affinities to the regulated sequences. This ‘high concentration’ paradigm has been formalized for the bacterial lac operon system and the necessity of DNA loops for efficient regulation has been demonstrated *in vivo* [63–67]. The lac operon is transcriptionally controlled by the binding of the lacI repressor to its lacO target site located 5′ of the controlled genes (Figure 3a). The association of the repressor to the functional lacO site (and the subsequent repression of transcription) is reinforced by the presence of additional lacO motifs in the vicinity. The ability of the tetrameric lacI repressor to bind simultaneously two operator sites thus favors DNA looping and the formation of a biochemical nano-reactor (Figure 3a) by enhancing the repression level, decreasing the sensitivity to number of repressors and reducing intrinsic noise. Nuclear compartments would similarly correspond to nano-reactors: a few number of reacting biomolecules co-localize in space, increasing their local concentration and thus promoting their

Figure 3



Nuclear compartments as self-assembled biochemical nano-reactors. **(a)** Schematic working principle of the *E. coli* lac operon nano-reactor. Dispersed lacO sites are located around the functional lacO site (5' of the transcribed region). lacI binding to these additional sites locally increases lacI concentration while lacI oligomerization leads to locus looping for efficient transcriptional repression. **(b)** Conserved features of eukaryotic nano-reactors leading to the formation of nuclear compartments. Dispersed binding sites for DNA-binding proteins are present on the same polymeric molecule, thereby providing a basal level of insulation as loci located on the same molecule are more likely to contact each other by looping. Multimerization of binding factors and/or interactions with common factors which themselves oligomerize lead to the formation of 'neighborhoods'. Finally, the formation of loops directed by sequence motifs on the 1D genome allows the insulation from each "neighborhood" and the formation of nuclear compartments.

biochemical activity on chromatin. Although proteins and sequences involved in the formation of specific compartments are different, the mechanisms driving spatial confinement may be similar to the lac operon case, based on three major features: (i) chain connectivity: the polymeric nature of chromatin induces a generic confinement as dispersed sequences on the same chain have a high probability to contact each other by looping. (ii) cooperativity: multimerization of DNA or chromatin binding proteins promote sequestration and physical bridging between target sequences. (iii) embedding and selectivity: the formation of TADs leads to 'insulated neighborhoods' [57^{••}], thus contributing to the structural but selective confinement of active or repressed modules.

At loop scales, enhancers recruit transcription factors, which in turn interact with the transcriptional mediator, thus establishing a transient physical contact with the regulated promoter leading to enhanced recruitment and stabilization of the transcriptional machinery at promoters [68] (Figure 3b). Interestingly, enhancer/promoter looping is observed even in the absence of transcriptional activity, suggesting that these loops constitute structural elements organizing the genome [13[•],19]. At larger scales, multimerization of Polycomb proteins in *Drosophila* [52[•],69] induces *cis* clustering of individual epigenomic

domains as well as *trans* clustering between distant domains (e.g., ANT-C and BX-C, located 10 Mb away from each other) [70]. This clustering of Polycomb-repressed genes into nuclear Polycomb bodies correlates with gene repression level [70]. Similar colocalisation of PcG repressed genes or HP1-regulated domains is observed in mammals, suggesting conservation of the regulatory logic [68]. At very large scale, the A/B compartments increase confinement of the TADs with similar transcriptional activity, providing a basal level of confinement and selectivity [17^{••},20].

Nuclear organization is known to increase over the course of development, with the appearance of numerous compartments (for review [71]). The creation of these is initiated by transcription factors binding specific genomic sites, which recruit chromatin-modifying enzymes [72–74]. This leads to spreading of specific chromatin marks *in cis* as well as *in trans* due to the spatial confinement by the insulated neighborhood [75–79]. Multimerizing proteins with specific affinities to these marks [50,80] then reinforce and enhance contacts between similar domains. Nucleation by a small number of factors coupled to self-assembly of the local chromatin therefore leads to the formation of nuclear compartments which further enhance, stabilize and/or perpetuate the active or repressed environment.

Ultimately, the assembly of nano-reactors is a conversion from a ‘hard-wired’ transcriptional program in which transcription factors determine the activity of genes to a ‘soft-wired’ self-reinforcing state in which spatially highly concentrated proteins maintain gene expression states, well in line with the fact that initiating transcription factors can disappear while cells maintain their fate.

Acknowledgements

We apologize to colleagues whose work could not be cited due to space constraints. We acknowledge our funding agencies: Agence Nationale de la Recherche (ANR-15-CE12-0006 EpiDevoMath; DJ/CV), Fondation pour la Recherche Médicale (DEI20151234396; DJ/CV), CNRS (DJ/CV), Institut Rhône-Alpin des Systèmes Complexes (DJ), program AGIR of University Grenoble Alpes (DJ), ENS de Lyon (CV), Swiss National Foundation (SNF) assistant professor PP00P3_133744/159320; PM), the Swiss Muscular Research Foundation (PM), the Novartis Biomedical Research Foundation (PM) and the University of Bern (PM).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Dekker J, Rippe K, Dekker M, Kleckner N: **Capturing chromosome conformation**. *Science* 2002, **295**:1306-1311.
2. de Wit E, de Laat W: **A decade of 3C technologies: insights into nuclear organization**. *Genes Dev* 2012, **26**:11-24.
3. Nora EP, Lajoie BR, Schulz EG, Giorgetti L, Okamoto I, Servant N, Piolot T, van Berkum NL, Meisig J, Sedat J *et al.*: **Spatial partitioning of the regulatory landscape of the X-inactivation centre**. *Nature* 2012, **485**:381-385.
4. Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, Robinson JT, Sanborn AL, Machol I, Omer AD, Lander ES *et al.*: **A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping**. *Cell* 2014, **159**:1665-1680.
- The authors perform high resolution Hi-C in different cell lines and different experimental conditions (fixed/unfixed). Beyond the TAD structure of the genome, they identify loops between convergent CTCF binding sites, which are conserved between the mouse and the human genome.
5. Boettiger AN, Bintu B, Moffitt JR, Wang S, Beliveau BJ, Fudenberg G, Imakaev M, Mirny LA, Wu C-t, Zhuang X: **Super-resolution imaging reveals distinct chromatin folding for different epigenetic states**. *Nature* 2016, **529**:418-422.
- Using fluorescent *in situ* hybridization and 3D Stochastic Optical Reconstruction Microscopy (STORM), a super resolution technique, the authors demonstrate that Polycomb, active and unmodified histone H3 epigenomic domains have different compaction levels. Moreover, they observe directly that epigenetically similar domains preferentially contact each other.
6. Fabre PJ, Benke A, Joye E, Nguyen Huynh TH, Manley S, Duboule D: **Nanoscale spatial organization of the HoxD gene cluster in distinct transcriptional states**. *Proc Natl Acad Sci U S A* 2015, **112**:13964-13969.
7. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I, Lajoie BR, Sabo PJ, Dorschner MO *et al.*: **Comprehensive mapping of long-range interactions reveals folding principles of the human genome**. *Science* 2009, **326**:289-293.
8. Marbouty M, Cournac A, Flot JF, Marie-Nelly H, Mozziconacci J, Koszul R: **Metagenomic chromosome conformation capture (meta3C) unveils the diversity of chromosome organization in microorganisms**. *Elife* 2014, **3**:e03318.
9. Sexton T, Yaffe E, Kenigsberg E, Bantignies F, Leblanc B, Hoichman M, Parrinello H, Tanay A, Cavalli G: **Three-dimensional folding and functional organization principles of the Drosophila genome**. *Cell* 2012, **148**:458-472.
10. Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B: **Topological domains in mammalian genomes identified by analysis of chromatin interactions**. *Nature* 2012, **485**:376-380.
11. Frenster JH, Allfrey VG, Mirsky AE: **Repressed and active chromatin isolated from interphase lymphocytes**. *Proc Natl Acad Sci U S A* 1963, **50**:1026-1032.
12. Dekker J, Heard E: **Structural and functional diversity of topologically associating domains**. *FEBS Lett* 2015, **589**:2877-2884.
13. Ghavi-Helm Y, Klein FA, Pakozdi T, Ciglar L, Noordermeer D, Huber W, Furlong EE: **Enhancer loops appear stable during development and are associated with paused polymerase**. *Nature* 2014, **512**:96-100.
- By focusing on enhancer/promoter contacts during two stages of *Drosophila* development and two different tissues, the authors observe that promoter/enhancer contacts are mainly unchanged between transcriptionally inactive and active states. This suggests that enhancer/promoter contacts are preformed and structurally organize the genome.
14. Guo Y, Xu Q, Canzio D, Shou J, Li J, Gorkin DU, Jung I, Wu H, Zhai Y, Tang Y *et al.*: **CRISPR inversion of CTCF sites alters genome topology and enhancer/promoter function**. *Cell* 2015, **162**:900-910.
15. Lupianez DG, Kraft K, Heinrich V, Krawitz P, Brancati F, Klopocki E, Horn D, Kayserili H, Opitz JM, Laxova R *et al.*: **Disruptions of topological chromatin domains cause pathogenic rewiring of gene-enhancer interactions**. *Cell* 2015, **161**:1012-1025.
- The authors of this study demonstrate that disruption of a TAD boundary in both mice and humans leads to ectopic contacts between a promoter and an enhancer located in a nearby TAD. This ultimately leads to misregulation of the EphA4 gene and finger malformations.
16. Franke M, Ibrahim DM, Andrey G, Schwarzer W, Heinrich V, Schopflin R, Kraft K, Kempfer R, Jerkovic I, Chan WL *et al.*: **Formation of new chromatin domains determines pathogenicity of genomic duplications**. *Nature* 2016, **538**:265-269.
17. Dixon JR, Jung I, Selvaraj S, Shen Y, Antosiewicz-Bourget JE, Lee AY, Ye Z, Kim A, Rajagopal N, Xie W *et al.*: **Chromatin architecture reorganization during stem cell differentiation**. *Nature* 2015, **518**:331-336.
- This paper describes TAD structures in human ES cells and 4 derived lineages. The TAD structure itself is remarkably stable between cell types, while the contact frequencies inside TADs and between TADs varies, affecting 36% of the genome.
18. Lonfat N, Montavon T, Darbellay F, Gitto S, Duboule D: **Convergent evolution of complex regulatory landscapes and pleiotropy at Hox loci**. *Science* 2014, **346**:1004-1006.
19. Jin F, Li Y, Dixon JR, Selvaraj S, Ye Z, Lee AY, Yen CA, Schmitt AD, Espinoza CA, Ren B: **A high-resolution map of the three-dimensional chromatin interactome in human cells**. *Nature* 2013, **503**:290-294.
20. Le Dily F, Bau D, Pohl A, Vicent GP, Serra F, Soronellas D, Castellano G, Wright RH, Ballare C, Filion G *et al.*: **Distinct structural transitions of chromatin topological domains correlate with coordinated hormone-induced gene regulation**. *Genes Dev* 2014, **28**:2151-2162.
21. Wong H, Marie-Nelly H, Herbert S, Carrivain P, Blanc H, Koszul R, Fabre E, Zimmer C: **A predictive computational model of the dynamic 3D interphase yeast nucleus**. *Curr Biol* 2012, **22**:1881-1890.
22. Kimura H, Shimooka Y, Nishikawa J-i, Miura O, Sugiyama S, Yamada S, Ohyama T: **The genome folding mechanism in yeast**. *J Biochem* 2013, **154**:137-147.
23. Avşaroğlu Bi, Bronk G, Gordon-Messer S, Ham J, Bressan DA, Haber JE, Kondev J: **Effect of chromosome tethering on nuclear organization in yeast**. *PLoS One* 2014, **9**:e102474.
24. Rosa A, Everaers R: **Structure and dynamics of interphase chromosomes**. *PLoS Comput Biol* 2008, **4**:e1000153.
25. Bohn M, Heermann DW: **Diffusion-driven looping provides a consistent framework for chromatin organization**. *PLoS One* 2010, **5**:e12218.

26. Halverson JD, Smrek J, Kremer K, Grosberg AY: **From a melt of rings to chromosome territories: the role of topological constraints in genome folding.** *Rep Prog Phys* 2014, **77**:022601.
 27. Gürsoy G, Xu Y, Kenter AL, Liang J: **Spatial confinement is a major determinant of the folding landscape of human chromosomes.** *Nucleic Acids Res* 2014, **42**:8223-8230.
 28. Grosberg A, Rabin Y, Halvin S, Neer A: **The crumpled globule model of the three-dimensional structure of DNA.** *Eur Phys Lett* 1993, **23**:373.
 29. Mirny LA: **The fractal globule as a model of chromatin architecture in the cell.** *Chromosome Res* 2011, **19**:37-51.
 30. Ho JW, Jung YL, Liu T, Alver BH, Lee S, Ikegami K, Sohn KA, Minoda A, Tolstorukov MY, Appert A *et al.*: **Comparative analysis of metazoan chromatin organization.** *Nature* 2014, **512**:449-452.
 31. Zhu Y, Chen Z, Zhang K, Wang M, Medovoy D, Whitaker JW, Ding B, Li N, Zheng L, Wang W: **Constructing 3D interaction maps from 1D epigenomes.** *Nat Commun* 2016, **7**:10812.
 32. Fraser J, Ferrai C, Chiariello AM, Schueler M, Rito T, Laudanno G, Barbieri M, Moore BL, Kraemer DC, Aitken S *et al.*: **Hierarchical folding and reorganization of chromosomes are linked to transcriptional changes in cellular differentiation.** *Mol Syst Biol* 2015, **11**:852.
 33. Barbieri M, Chotalia M, Fraser J, Lavitas L-M, Dostie J, Pombo A, Nicodemi M: **Complexity of chromatin folding is captured by the strings and binders switch model.** *Proc Natl Acad Sci U S A* 2012, **109**:16173-16178.
 34. Brackley CA, Johnson J, Kelly S, Cook PR, Marenduzzo D: **Simulated binding of transcription factors to active and inactive regions folds human chromosomes into loops, rosettes and topological domains.** *Nucleic Acids Res* 2016, **44**:3503-3512.
 35. Jerabek H, Heermann DW: **Expression-dependent folding of interphase chromatin.** *PLoS One* 2012, **7**:e37525.
 36. Benedetti F, Dorier J, Burnier Y, Stasiak A: **Models that include supercoiling of topological domains reproduce several known features of interphase chromosomes.** *Nucleic Acids Res* 2014, **42**:2848-2855.
 37. Doyle B, Fudenberg G, Imakaev M, Mirny LA: **Chromatin loops as allosteric modulators of enhancer-promoter interactions.** *PLoS Comput Biol* 2014, **10**:e1003867.
 38. Ganai N, Sengupta S, Menon GI: **Chromosome positioning from activity-based segregation.** *Nucleic Acids Res* 2014, **42**:4145-4159.
 39. Jost D, Carrivain P, Cavalli G, Vaillant C: **Modeling epigenome folding: formation and dynamics of topologically associated chromatin domains.** *Nucleic Acids Res* 2014, **42**:9553-9561.
- A heteropolymer model is introduced which suggests that epigenomic-driven interactions might be an important driver of chromosome folding at megabase scales. In agreement with experimental data, the model predicts that TAD formation is fast and results from the internal collapse of epigenomic domain and that inter-TAD interactions are dynamic and emerge at longer time-scales.
40. Tark-Dame M, Jerabek H, Manders EMM, Heermann DW, van Driel R: **Depletion of the chromatin looping proteins CTCF and cohesin causes chromatin compaction: insight into chromatin folding by polymer modelling.** *PLoS Comput Biol* 2014, **10**:e1003877.
 41. Nazarov LI, Tamm MV, Avetsov VA, Nechaev SK: **A statistical model of intra-chromosome contact maps.** *Soft Matter* 2015, **11**:1019-1025.
 42. Sanborn AL, Rao SS, Huang SC, Durand NC, Huntley MH, Jewett AI, Bochkov ID, Chinnappan D, Cutkosky A, Li J *et al.*: **Chromatin extrusion explains key features of loop and domain formation in wild-type and engineered genomes.** *Proc Natl Acad Sci U S A* 2015, **112**:E6456-E6465.
- A polymer model for chromatin extrusion is introduced and shown to agree quantitatively with experimental data. The role of oriented CTCF sites is explored experimentally, showing that the model is able to predict the effect of CTCF site inversions or deletions.
43. Ulianov SV, Khrameeva EE, Gavrilov AA, Flyamer IM, Kos P, Mikhaleva EA, Penin AA, Logacheva MD, Imakaev MV, Chertovich A *et al.*: **Active chromatin and transcription play a key role in chromosome partitioning into topologically associating domains.** *Genome Res* 2016, **26**:70-84.
 44. Fudenberg G, Imakaev M, Lu C, Goloborodko A, Abdennur N, ●● Mirny LA: **Formation of chromosomal domains by loop extrusion.** *Cell Rep* 2016, **15**:2038-2049.
- This paper introduces a rigorous analysis of the loop extrusion model and the role of boundary elements [note that preprint publication of this work was first released in bioRxiv in August 2015]. Compared to other possible models, the loop extrusion model is shown to fit the data best. Polymer simulations and genomic data are used to identify the functional role of CTCF and cohesin.
45. Tiana G, Amitai A, Pollex T, Piolot T, Holcman D, Heard E, Giorgetti L: **Structural fluctuations of the chromatin fiber within topologically associating domains.** *Biophys J* 2016, **110**:1234-1245.
 46. Giorgetti L, Galupa R, Nora EP, Piolot T, Lam F, Dekker J, Tiana G, Heard E: **Predictive polymer modeling reveals coupled fluctuations in chromosome conformation and transcription.** *Cell* 2014, **157**:950-963.
 47. Olarte-Plata JD, Haddad N, Vaillant C, Jost D: **The folding landscape of the epigenome.** *Phys Biol* 2016, **13**:026001.
 48. Care BR, Emeriau P-E, Cortini R, Victor J-M: **Chromatin epigenomic domain folding: size matters.** *AIMS Biophys* 2015, **2**:517-530.
 49. Canzio D, Liao M, Naber N, Pate E, Larson A, Wu S, Marina DB, Garcia JF, Madhani HD, Cooke R *et al.*: **A conformational switch in HP1 releases auto-inhibition to drive heterochromatin assembly.** *Nature* 2013, **496**:377-381.
 50. Isono K, Endo TA, Ku M, Yamada D, Suzuki R, Sharif J, Ishikura T, Toyoda T, Bernstein BE, Koseki H: **SAM domain polymerization links subnuclear clustering of PRC1 to gene silencing.** *Dev Cell* 2013, **26**:565-577.
 51. Hiragami-Hamada K, Soeroes S, Nikolov M, Wilkins B, Kreuz S, Chen C, De La Rosa-Velázquez IA, Zenn HM, Kost N, Pohl W *et al.*: **Dynamic and flexible H3K9me3 bridging via HP1 β dimerization establishes a plastic state of condensed chromatin.** *Nat Commun* 2016, **7**:11310.
 52. Wani AH, Boettiger AN, Schorderet P, Ergun A, Mürger C, ● Sadreyev RI, Zhuang X, Kingston RE, Francis NJ: **Chromatin topology is coupled to Polycomb group protein subnuclear organization.** *Nat Commun* 2016, **7**:10291.
- This study makes use of a polymer model using interacting binders, first introduced by Barbieri *et al.* [33], to mechanistically dissect the formation of Polycomb bodies in flies. Results are then compared to *in vivo* super-resolution microscopy data.
53. Brackley CA, Brown JM, Waithe D, Babbs C, Davies J, Hughes JR, Buckle VJ, Marenduzzo D: **Predicting the three-dimensional folding of cis-regulatory regions in mammalian genomes using bioinformatic data and polymer models.** *Genome Biol* 2016, **17**:59.
 54. Naumova N, Imakaev M, Fudenberg G, Zhan Y, Lajoie BR, Mirny LA, Dekker J: **Organization of the mitotic chromosome.** *Science* 2013, **342**:948-953.
 55. Nagano T, Lubling Y, Stevens TJ, Schoenfelder S, Yaffe E, Dean W, Laue ED, Tanay A, Fraser P: **Single-cell Hi-C reveals cell-to-cell variability in chromosome structure.** *Nature* 2013, **502**:59-64.
 56. Vietri Rudan M, Barrington C, Henderson S, Ernst C, Odom DT, Tanay A, Hadjuri S: **Comparative Hi-C reveals that CTCF underlies evolution of chromosomal domain architecture.** *Cell Rep* 2015, **10**:1297-1309.
 57. Downen JM, Fan ZP, Hnisz D, Ren G, Abraham BJ, Zhang LN, ●● Weintraub AS, Schuijers J, Lee TI, Zhao K *et al.*: **Control of cell identity genes occurs in insulated neighborhoods in mammalian chromosomes.** *Cell* 2014, **159**:374-387.
- From a genome-wide SMC1 ChIA-PET experiment performed in murine ESC, the authors identify for the first time cohesin-mediated, TAD-like interaction domains with CTCF-enriched boundaries. By specific deletion of these boundaries they show how these domains, by constituting

structural “insulated neighborhood” may contribute to the control of the superenhancer-mediated activity of pluripotent genes as well as the PcG-repression of lineage-specific genes.

58. Nasmyth K: **Disseminating the genome: joining, resolving, and separating sister chromatids during mitosis and meiosis.** *Annu Rev Genet* 2001, **35**:673-745.
59. Alipour E, Marko JF: **Self-organization of domain structures by DNA-loop-extruding enzymes.** *Nucleic Acids Res* 2012, **40**:11202-11212.
60. Crane E, Bian Q, McCord RP, Lajoie BR, Wheeler BS, Ralston EJ, Uzawa S, Dekker J, Meyer BJ: **Condensin-driven remodelling of X chromosome topology during dosage compensation.** *Nature* 2015, **523**:240-244.
61. Goloborodko A, Imakaev MV, Marko JF, Mirny L: **Compaction and segregation of sister chromatids via active loop extrusion.** *Elife* 2016:5.
62. Goloborodko A, Marko JF, Mirny LA: **Chromosome compaction by active loop extrusion.** *Biophys J* 2016, **110**:2162-2168.
63. Vilar JM, Leibler S: **DNA looping and physical constraints on transcription regulation.** *J Mol Biol* 2003, **331**:981-989.
64. Oehler S, Alberti S, Muller-Hill B: **Induction of the lac promoter in the absence of DNA loops and the stoichiometry of induction.** *Nucleic Acids Res* 2006, **34**:606-612.
65. Oehler S, Amouyal M, Kolkhof P, von Wilcken-Bergmann B, Muller-Hill B: **Quality and position of the three lac operators of *E. coli* define efficiency of repression.** *EMBO J* 1994, **13**:3348-3355.
66. Saiz L, Vilar JM: **DNA looping: the consequences and its control.** *Curr Opin Struct Biol* 2006, **16**:344-350.
67. Vilar JM, Saiz L: **Systems biophysics of gene expression.** *Biophys J* 2013, **104**:2574-2585.
68. Spitz F: **Gene regulation at a distance: from remote enhancers to 3D regulatory ensembles.** *Semin Cell Dev Biol* 2016, **57**:57-67.
69. Lanzuolo C, Orlando V: **The function of the epigenome in cell reprogramming.** *Cell Mol Life Sci* 2007, **64**:1043-1062.
70. Bantignies F, Roure V, Comet I, Leblanc B, Schuettengruber B, Bonnet J, Tixier V, Mas A, Cavalli G: **Polycomb-dependent regulatory contacts between distant Hox loci in *Drosophila*.** *Cell* 2011, **144**:214-226.
71. Püeschel R, Coraggio F, Meister P: **From single genes to entire genomes: the search for a function of nuclear organization.** *Development* 2016, **143**:910-923.
72. Beisel C, Paro R: **Silencing chromatin: comparing modes and mechanisms.** *Nat Rev Genet* 2011, **12**:123-135.
73. Zhang T, Cooper S, Brockdorff N: **The interplay of histone modifications - writers that read.** *EMBO Rep* 2015, **16**:1467-1481.
74. Chen T, Dent SY: **Chromatin modifiers and remodellers: regulators of cellular differentiation.** *Nat Rev Genet* 2014, **15**:93-106.
75. Obersiebenig MJ, Pallesen EM, Sneppen K, Trusina A, Thon G: **Nucleation and spreading of a heterochromatic domain in fission yeast.** *Nat Commun* 2016, **7**:11518.
76. Angel A, Song J, Dean C, Howard M: **A polycomb-based switch underlying quantitative epigenetic memory.** *Nature* 2011, **476**:105-108.
77. Dodd IB, Micheelsen MA, Sneppen K, Thon G: **Theoretical analysis of epigenetic cell memory by nucleosome modification.** *Cell* 2007, **129**:813-822.
78. Jost D: **Bifurcation in epigenetics: implications in development, proliferation, and diseases.** *Phys Rev E Stat Nonlin Soft Matter Phys* 2014, **89**:010701.
79. Soshnev AA, Josefowicz SZ, Allis CD: **Greater than the sum of parts: complexity of the dynamic epigenome.** *Mol Cell* 2016, **62**:681-694.
80. Simon JA, Kingston RE: **Occupying chromatin: polycomb mechanisms for getting to genomic targets, stopping transcriptional traffic, and staying put.** *Mol Cell* 2013, **49**:808-824.