

# Mid-Term Exam

In solving the following exercises you can use the statements of the main theorems proved during the course such as: Büchi's theorem about the decidability of the MSO theory of  $(\mathbb{N}, <)$  and Gödel's theorem about the undecidability of FO theory of  $(\mathbb{N}, +, \times)$ . On the other hand you have to be very precise and clear about how you invoke these theorems to solve the exercises.

**Exercise 1.** Let  $\mathbb{N}$  be the set of natural numbers. We have proved during the course, using the finite-automata method, that the First order (FO) theory of  $(\mathbb{N}, \leq, +)$ , known as *Presburger arithmetic*, is decidable.

Let  $Q \subseteq \mathbb{N}$  be the predicate (i.e., unary relation) on natural numbers which holds on the powers of 2:

$$Q = \{1, 2, 4, 8, 16, 32, \dots\} \quad \text{or, in other words,} \quad Q(n) \Leftrightarrow n = 2^m \text{ for some } m \in \mathbb{N}$$

**To do:** Is the FO-theory of  $(\mathbb{N}, \leq, +, Q)$  decidable?

**Exercise 2.** For a given  $k \in \mathbb{N}$ , let  $\times_k : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  be multiplication on natural numbers defined only on numbers less than  $k$ . Formally,

$$n \times_k m = \begin{cases} n \times m & \text{if both } n < k \text{ and } m < k \\ 0 & \text{otherwise} \end{cases}$$

where  $\times$  is the ordinary (full) multiplication on natural numbers.

**To do:** Is the FO-theory of  $(\mathbb{N}, \leq, +, \{\times_k\}_{k \in \mathbb{N}})$  decidable?

**Exercise 3.** Is it possible to write a formula  $\phi(x)$  in the WMSO-theory of  $(\mathbb{N}, <)$  such that

$$\phi(n) \text{ holds} \Leftrightarrow n \text{ is a prime number?}$$

*Hint:* to prove that it is possible, just write down the formula  $\phi$ . To prove that it is impossible typically we use the *pumping lemma*.

**Exercise 4.** Let  $L \subseteq \{a, b\}^\omega$  be defined as:

$$L = \{w \in \{a, b\}^\omega \mid w \text{ has exactly one position labeled by } a\}.$$

**To do:**

1. Write down a Büchi automaton  $A$  that accepts  $L$ .
2. Is the language  $L$  definable by a *deterministic* Büchi automaton?

**Exercise 5.** Let  $\Sigma$  be the alphabet  $\{0, 1, 2\}$ . Given an infinite word  $w \in \Sigma^\omega$ , we define  $\infty(w) \subseteq \Sigma$  as the set of elements of the alphabet that occur *infinitely many times* in  $w$ . For example:

1. if  $w = (12)^\omega = 1212121212\dots$  then  $\infty(w) = \{1, 2\}$ ,
2. if  $w = 0.(12)^\omega = 01212121212\dots$  then  $\infty(w) = \{1, 2\}$ ,
3. if  $w = 0^\omega = 0000000\dots$  then  $\infty(w) = \{0\}$ ,

Let  $L \subseteq \Sigma^\omega$  be the set

$$\{w \mid \text{the greatest number in } \infty(w) \text{ is even}\}$$

So for example:

- if  $w = (012)^\omega$  then  $w \in L$  because  $\infty(w) = \{0, 1, 2\}$  and  $\max\{0, 1, 2\} = 2$  is even.
- if  $w = 2.(01)^\omega$  then  $w \notin L$  because  $\infty(w) = \{0, 1\}$  and  $\max\{0, 1, 2\} = 1$  is not even.

**To do:**

- Is  $L$  definable by a Büchi automaton?
- Is  $L$  definable by a deterministic Büchi automaton?