

Approximation of a volume-preserving homeomorphism by a volume-preserving diffeomorphism

Jean-Claude Sikorav
September 24, 2007

The purpose of this note is to prove the following statement, which is "well known" but for which there does not seem to exist a published proof.

Proposition. *Let M be a smooth n -manifold equipped with a smooth volume element. Let φ be a volume-preserving homeomorphism. Assume that φ can be C^0 -approximated by smooth diffeomorphisms [this is always true if $n \leq 3$]. Then φ can be C^0 -approximated by smooth volume-preserving diffeomorphisms.*

Proof. To simplify and get at the heart of the matter, let us assume that M is compact and oriented, and that φ preserves the smooth oriented volume element $\omega \in \Omega^n(M)$.

Fix any Riemannian metric, and let $\varepsilon > 0$. Let K be a smooth triangulation such that $\text{diam}(\varphi(\sigma)) < \varepsilon$ for every n -simplex $\sigma \in K^{(n)}$. By hypothesis, for any $\eta > 0$ there exists $\psi_1 \in \text{Diff}(M)$ such that $d(\psi_1, \varphi) < \eta$. This implies

$$(*) \quad \max_{\sigma \in K^{(n)}} |\text{vol}(\psi_1(\sigma)) - \text{vol}(\sigma)| = O(\eta), \quad \eta \rightarrow 0.$$

Using an idea of E. Giroux, we choose some maximal tree T in the dual triangulation, and write T as a union of almost-disjoint simplicial paths P_1, \dots, P_k . Thus the n -simplices in all these paths are distinct and are all the n -simplices in $K^{(n)}$. We modify ψ_1 successively on each n -simplex on P_1, \dots, P_k to obtain ψ_2 such that $\text{vol}(\psi_2(\sigma)) = \text{vol}(\sigma)$ for every σ . If η is sufficiently small, (*) implies that this can be done with the property

$$(**) \quad (\forall \sigma \in K^{(n)}) \quad d(\psi_2, \varphi) + \text{diam}(\psi_2(\sigma)) < \varepsilon.$$

By construction, the n -form $\alpha := \psi_2^* \omega - \omega$ satisfies $\int_{\sigma} \alpha = 0$ for every n -simplex. Thus it admits a primitive β which vanishes on $K^{(n-1)}$. Indeed, by [ST], Lemma 2 p.148 [p.165 in the Springer edition] it has a primitive β_1 such that $\int_{\tau} \beta_1 = 0$ for every $\tau \in K^{(n-1)}$; thus β_1 is exact on $K^{(n-1)}$, thus there exists $\gamma \in \Omega^{n-2}(M)$ such that $d\gamma = \beta_1$ on $K^{(n-1)}$, and one can take $\beta = \beta_1 - d\gamma$.

Applying Moser's method to the path $\omega_t = \omega + td\beta$ provides a time-dependent vector field X_t such that $\omega_t(X_t, \cdot) = -\beta$, whose integration gives an isotopy φ_t with $\varphi_0 = \text{Id}$, $\varphi_t^* \omega_t = \omega$ so that $\psi = \psi_2 \circ \varphi_1$ preserves ω . Since $\beta|_{K^{(n-1)}} = 0$, we have $\varphi_t|_{K^{(n-1)}} = \text{Id}$, thus $\psi|_{K^{(n-1)}} = \psi_2$. Finally, (**) implies $d(\psi, \varphi) < \varepsilon$, which finishes the proof.

[ST] I.M. Singer, J.A. Thorpe, *Lecture notes on elementary topology and geometry*, Scott Foresman 1967. Republished in Springer Undergraduate Texts.

Remark (added on October 31, 2007). In fact, there exists a proof of this approximation result in a 2006 preprint of Yong-Geun Oh: " C^0 -coerciveness of Moser's problem and smoothing area preserving homeomorphisms", <http://xxx.lanl.gov/abs/math/0601183>. I thank Stefan Mueller for making me aware of this reference.

Oh's proof is more sophisticated, using in particular a key analytical estimate due to Dacorogna and Moser. Thus I believe that our proof is of independent interest.