

Bounded functions on the character variety

Konstantin Ardakov and Laurent Berger

With an appendix by Dragoş Crişan and Jingjie Yang

Abstract. This paper is motivated by an open question in p -adic Fourier theory, that seems to be more difficult than it appears at first glance. Let L be a finite extension of \mathbb{Q}_p with ring of integers \mathcal{O}_L and let \mathbb{C}_p denote the completion of an algebraic closure of \mathbb{Q}_p . In their work on p -adic Fourier theory, Schneider and Teitelbaum defined and studied the character variety \mathfrak{X} . This character variety is a rigid analytic curve over L that parameterizes the set of locally L -analytic characters $\lambda : (\mathcal{O}_L, +) \rightarrow (\mathbb{C}_p^\times, \times)$. One of the main results of Schneider and Teitelbaum is that over \mathbb{C}_p , the curve \mathfrak{X} becomes isomorphic to the open unit disk. Let $\Lambda_L(\mathfrak{X})$ denote the ring of bounded-by-one functions on \mathfrak{X} . If $\mu \in \mathcal{O}_L[[\mathcal{O}_L]]$ is a measure on \mathcal{O}_L , then $\lambda \mapsto \mu(\lambda)$ gives rise to an element of $\Lambda_L(\mathfrak{X})$. The resulting map $\mathcal{O}_L[[\mathcal{O}_L]] \rightarrow \Lambda_L(\mathfrak{X})$ is injective. The question is: do we have $\Lambda_L(\mathfrak{X}) = \mathcal{O}_L[[\mathcal{O}_L]]$?

In this paper, we prove various results that were obtained while studying this question. In particular, we give several criteria for a positive answer to the above question. We also recall and prove the “Katz isomorphism” that describes the dual of a certain space of continuous functions on \mathcal{O}_L . An important part of our paper is devoted to providing a proof of this theorem which was stated in 1977 by Katz. We then show how it applies to the question. Besides p -adic Fourier theory, the above question is related to the theory of formal groups, the theory of integer valued polynomials on \mathcal{O}_L , p -adic Hodge theory, and Iwasawa theory.

CONTENTS

1. Introduction	1000
1.1. Motivation	1000
1.2. The character variety	1001
1.3. Schneider and Teitelbaum’s uniformization	1001
1.4. The operators φ_q, ψ_q	1002
1.5. The polynomials P_n	1002
1.6. Galois-continuous functions and the Katz map	1003
1.7. Applications of the Katz isomorphism	1004
1.8. Other criteria	1005
1.9. Acknowledgements	1006

2.	The character variety	1006
2.1.	Notation	1006
2.2.	The p -adic Fourier transform	1006
2.3.	Lubin-Tate formal groups	1007
2.4.	A review of p -divisible groups	1007
2.5.	Cartier duality for p -divisible groups	1008
2.6.	The character $\tau : G_L \rightarrow o_L^\times$ and the period Ω	1009
2.7.	The Amice-Katz transform	1011
2.8.	The Schneider-Teitelbaum uniformisation	1012
2.9.	$\Lambda_L(\mathfrak{X})$ and the twisted G_L -action on $\mathbb{C}_p[[Z]]$	1013
2.10.	Some properties of $\Lambda_L(\mathfrak{X})$	1014
3.	The Katz isomorphism	1018
3.1.	The ψ_q -operator	1018
3.2.	The covariant bialgebra of \mathcal{G}	1021
3.3.	Gal-continuous functions	1026
3.4.	The largest ψ_q -stable o_L -submodule of $o_L[[Z]]$	1029
3.5.	The Newton polygon of $\Delta_1(Z) - 1$	1032
3.6.	Verifying Conjecture 3.3.11 in a special case	1034
4.	Integer-valued polynomials	1039
4.1.	The algebraic dual of $\mathcal{O}^\circ(\mathfrak{X}_K)$	1039
4.2.	The matrix coefficients $\rho_{i,j}(Y)$	1043
4.3.	Calculating the matrix coefficients $\sigma_{i,j}(Y)$	1050
5.	Consequences of the Katz isomorphism	1055
5.1.	Equivariant endomorphisms of L_∞	1055
5.2.	The dual of the ring of integers of a p -adic Lie extensions	1058
5.3.	The operator ψ and the span of the P_n	1062
6.	Other criteria	1064
6.1.	The Lubin-Tate derivative	1064
6.2.	Changing the base field	1065
	Appendix A. An algorithm for whether the $\sigma_{i,j}$'s span $\text{Int}(o_L, o_L)$	1066
	References	1079

1. INTRODUCTION

1.1. Motivation. Let L be a finite extension of \mathbb{Q}_p and let \mathbb{C}_p denote the completion of an algebraic closure of \mathbb{Q}_p . In their work on p -adic Fourier theory [28], Schneider and Teitelbaum defined and studied the character variety \mathfrak{X} . This character variety is a rigid analytic curve over L that parameterizes the set of locally L -analytic characters $\lambda : (o_L, +) \rightarrow (\mathbb{C}_p^\times, \times)$. One of the main results of Schneider and Teitelbaum is that over \mathbb{C}_p , the curve \mathfrak{X} becomes isomorphic to the open unit disk.

The ring $\mathcal{O}_L(\mathfrak{X})$ of holomorphic functions on \mathfrak{X} is a Prüfer domain, with an action of o_L coming from the natural action of o_L on the set of locally

L -analytic characters. One can then localize and complete $\mathcal{O}_L(\mathfrak{X})$ in order to obtain the Robba ring $\mathcal{R}_L(\mathfrak{X})$, and define (φ, o_L^\times) -modules over that ring and some of its subrings. These objects are defined and studied in [5], with the hope that they will be useful for a generalization of the p -adic local Langlands correspondence from $\mathrm{GL}_2(\mathbb{Q}_p)$ to $\mathrm{GL}_2(L)$.

In this paper, we instead consider a natural subring of $\mathcal{O}_L(\mathfrak{X})$, the ring $\Lambda_L(\mathfrak{X})$ of functions whose norms are bounded above by 1. If $\mu \in o_L[[o_L]]$ is a measure on o_L , then $\lambda \mapsto \mu(\lambda)$ gives rise to such a function $\mathfrak{X} \rightarrow \mathbb{C}_p$. The resulting map $o_L[[o_L]] \rightarrow \Lambda_L(\mathfrak{X})$ is injective. We do not know of any example of an element of $\Lambda_L(\mathfrak{X})$ that is not in the image of the above map.

Question 1.1.1. Do we have $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$?

This question seems to be more difficult than it appears at first glance, and so far we have not been able to answer it (except of course for $L = \mathbb{Q}_p$). The results of this paper were obtained while we were studying this problem. A related question is raised in remark 2.5 of [11]. We now give more details about the character variety \mathfrak{X} , and then explain our main results.

1.2. The character variety. Let \mathfrak{B} denote the open unit disk, seen as a rigid analytic variety. This space naturally parameterizes the set of locally \mathbb{Q}_p -analytic characters $\lambda : (\mathbb{Z}_p, +) \rightarrow (\mathbb{C}_p^\times, \times)$. Indeed, if K is a closed subfield of \mathbb{C}_p and $z \in \mathfrak{m}_K = \mathfrak{B}(K)$, then the map $\lambda_z : a \mapsto (1+z)^a$ is a K -valued locally \mathbb{Q}_p -analytic character on \mathbb{Z}_p , and every such character arises in this way. Note that $\lambda'_z(0) = \log(1+z)$. If $d = [L : \mathbb{Q}_p]$, then $o_L \simeq \mathbb{Z}_p^d$ and hence \mathfrak{B}^d parameterizes the set of locally \mathbb{Q}_p -analytic characters $\lambda : (o_L, +) \rightarrow (\mathbb{C}_p^\times, \times)$. Such a character is locally L -analytic if and only if $\lambda'(0)$ is L -linear. In coordinates $z = (z_1, \dots, z_d)$, there exists $\alpha_2, \dots, \alpha_d \in L$ such that the character corresponding to z is locally L -analytic if and only if $\log(1+z_i) = \alpha_i \cdot \log(1+z_1)$ for all $i = 2, \dots, d$. These $d-1$ Cauchy–Riemann equations cut out the character variety \mathfrak{X} inside \mathfrak{B}^d . Schneider and Teitelbaum showed [28] that \mathfrak{X} is a smooth rigid analytic group curve over L .

The ring of \mathbb{Q}_p -analytic distributions $D^{\mathbb{Q}_p\text{-an}}(o_L, L)$ on o_L is isomorphic to the ring of power series in d variables that converge on the open unit polydisk. Every distribution $\mu \in D^{\mathbb{Q}_p\text{-an}}(o_L, L)$ gives rise to an element of $\mathcal{O}_L(\mathfrak{X})$, defined by the map $\lambda \mapsto \mu(\lambda)$. This gives rise to a surjective (but not injective if $L \neq \mathbb{Q}_p$) map $D^{\mathbb{Q}_p\text{-an}}(o_L, L) \rightarrow \mathcal{O}_L(\mathfrak{X})$, whose restriction to $o_L[[o_L]]$ is injective and has image contained in $\Lambda_L(\mathfrak{X})$.

1.3. Schneider and Teitelbaum’s uniformization. We now explain why over \mathbb{C}_p , the curve \mathfrak{X} becomes isomorphic to the open unit disk. Let $G_L = \mathrm{Gal}(\overline{\mathbb{Q}_p}/L)$. Choose a uniformizer π of o_L and let \mathcal{G} denote the Lubin–Tate formal group attached to π . This gives us a Lubin–Tate character $\chi_\pi : G_L \rightarrow o_L^\times$ and, once we have chosen a coordinate Z on \mathcal{G} , a formal addition law $X \oplus Y \in o_L[[X, Y]]$, endomorphisms $[a](Z) \in o_L[[Z]]$ for all $a \in o_L$, and a logarithm $\log_{\mathrm{LT}}(Z) \in L[[Z]]$.

By the work of Tate on p -divisible groups, there is a non-trivial homomorphism $\mathcal{G} \rightarrow \mathbf{G}_m$ defined over $o_{\mathbb{C}_p}$. Concretely, there exists a power series $G(Z) \in o_{\mathbb{C}_p}[[Z]]$ (where $G(Z)$ is a generator of $\mathrm{Hom}_{o_{\mathbb{C}_p}}(\mathcal{G}, \mathbf{G}_m)$) such that $(1 + G(X \oplus Y)) = (1 + G(X)) \cdot (1 + G(Y))$. If $z \in \mathfrak{m}_{\mathbb{C}_p}$, then the map $\lambda_z : a \mapsto 1 + G([a](z))$ is a locally L -analytic character on o_L , and every such character arises in this way. This explains the main idea behind the proof of the statement that over \mathbb{C}_p , the curve \mathfrak{X} becomes isomorphic to the open unit disk.

In particular, $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$ is isomorphic to the ring of power series $\sum_{i \geq 0} a_i Z^i$ with $a_i \in \mathbb{C}_p$ that converge on the open unit disk. Let χ_{cyc} denote the cyclotomic character, and let $\tau : G_L \rightarrow o_L^\times$ denote the character $\tau = \chi_{\mathrm{cyc}} \cdot \chi_\pi^{-1}$. The Galois group G_L acts on $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$ by the formula $g(\sum_{i \geq 0} a_i Z^i) = \sum_{i \geq 0} g(a_i)[\tau(g)^{-1}](Z)^i$. This action is called the twisted Galois action, and we write $G_L, *$ to recall the twist. It follows from the Ax-Sen-Tate theorem that $\mathbb{C}_p^{G_L} = L$ and then, by unravelling the definitions, that $\mathcal{O}_L(\mathfrak{X}) = \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})^{G_L, *}$. At the level of bounded functions, this tells us that $\Lambda_L(\mathfrak{X}) = o_{\mathbb{C}_p}[[Z]]^{G_L, *}$. The natural map $o_L[[o_L]] \rightarrow \Lambda_L(\mathfrak{X})$ sends, for instance, the Dirac measure δ_a with $a \in o_L$ to $1 + G([a](Z)) \in o_{\mathbb{C}_p}[[Z]]^{G_L, *}$.

1.4. The operators φ_q, ψ_q . The monoid (o_L, \times) acts on o_L by multiplication, and hence on the set of locally L -analytic characters, on \mathfrak{X} , and on the ring $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$. If $a \in o_L$, this action is given by $f(Z) \mapsto f([a](Z))$. Let q denote the cardinality of the residue field k_L of o_L and let φ_q denote the action of π on $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$. The map φ_q is injective and the ring $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$ is a free $\varphi_q(\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}))$ -module of rank q . Let $\psi_q : \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}) \rightarrow \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$ be the map defined by $\varphi_q(\psi_q(f(Z))) = 1/q \cdot \mathrm{Tr}_{\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})/\varphi_q(\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}))}(f(Z))$. The action of o_L and the operator ψ_q commute with the twisted action of G_L , and therefore preserve $\mathcal{O}_L(\mathfrak{X})$. If we consider the image of the map $D^{\mathbb{Q}_p - \mathrm{an}}(o_L, L) \rightarrow \mathcal{O}_L(\mathfrak{X})$, we have $a \cdot \delta_b = \delta_{ab}$ and $\psi_q(\delta_b) = 0$ if $b \in o_L^\times$ and $\psi_q(\delta_b) = \delta_{b/\pi}$ if $b \in \pi o_L$. In particular, $o_L[[o_L]]^{\psi_q=0}$ coincides with $o_L[[o_L^\times]]$, those measures that are supported in o_L^\times . We use later on the fact (Lemma 5.1.9) that $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if $\Lambda_L(\mathfrak{X})^{\psi_q=0} = o_L[[o_L^\times]]$. Note that if $L \neq \mathbb{Q}_p$, then $\psi_q(\Lambda_{\mathbb{C}_p}(\mathfrak{X}))$ is not contained in $\Lambda_{\mathbb{C}_p}(\mathfrak{X})$ as $\mathrm{Tr}_{\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})/\varphi_q(\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}))}(f(Z))$ is divisible by π , but not always by q . Our first result is the following.

Theorem 1.4.1. *We have $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if $\psi_q(\Lambda_L(\mathfrak{X})) \subset \Lambda_L(\mathfrak{X})$.*

This is proved at the end of §3.1.

1.5. The polynomials P_n . Recall that $G(Z)$ is a generator of $\mathrm{Hom}_{o_{\mathbb{C}_p}}(\mathcal{G}, \mathbf{G}_m)$ and that $\tau = \chi_{\mathrm{cyc}} \cdot \chi_\pi^{-1}$. In fact, we have $G(Z) = \exp(\Omega \cdot \log_{\mathrm{LT}}(Z)) - 1 = \Omega \cdot Z + O(Z^2)$, where Ω is a certain special element of $\mathfrak{m}_{\mathbb{C}_p}$ such that $g(\Omega) = \tau(g) \cdot \Omega$. In particular, for all $n \geq 0$, there exists a polynomial $P_n(Y) \in L[Y]$ such that $1 + G(Z) = \sum_{n \geq 0} P_n(\Omega) \cdot Z^n$. For $n \geq 0$, the polynomial $P_n(Y)$ is of degree n , and its leading coefficient is $1/n!$. For example, assume that the

coordinate Z is chosen in a way that $\log_{\text{LT}}(Z) = \sum_{k \geq 0} Z^q / \pi^k$. Then we have (see Proposition 4.3.1 for more details)

$$P_n(Y) = \sum_{n_0 + qn_1 + \dots + q^d n_d = n} \frac{Y^{n_0 + \dots + n_d}}{n_0! \dots n_d! \cdot \pi^{1 \cdot n_1 + 2 \cdot n_2 + \dots + d \cdot n_d}}.$$

If $a \in o_L$, then $G([a](Z)) = \sum_{n \geq 1} P_n(\Omega) \cdot [a](Z)^n = \sum_{n \geq 1} P_n(a\Omega) \cdot Z^n$. This implies for instance that $P_n(a\Omega) \in o_{\mathbb{C}_p}$ for all $a \in o_L$. For $n \geq 0$ and $i \geq n$, let $\sigma_{n,i}(Y) \in L[Y]$ denote the polynomials such that $[a](Z)^n = \sum_{i \geq n} \sigma_{n,i}(a) Z^i$ for all $a \in o_L$. The $\sigma_{n,i}(Y)$ are all elements of Int , the o_L -submodule of $L[Y]$ of integer valued polynomials on o_L . The fact that $\sum_{n \geq 0} P_n(\Omega) \cdot [a](Z)^n = \sum_{n \geq 0} P_n(a\Omega) \cdot Z^n$ implies that $P_n(a\Omega) = \sum_{i=0}^n \sigma_{i,n}(a) P_i(\Omega)$.

If $\mu \in D^{\mathbb{Q}_p\text{-an}}(o_L, L)$, its image in $\mathcal{O}_L(\mathfrak{X})$ is therefore $f_\mu(Z) = \sum_{n \geq 0} Z^n \cdot \sum_{i=0}^n \mu(\sigma_{i,n}) P_i(\Omega)$. Let Pol denote the o_L -span of the $\sigma_{n,i}(Y)$ inside $L[Y]$, so that $\text{Pol} \subset \text{Int}$. The following gives a relation between our question and the theory of integer valued polynomials ([30], [31]):

Theorem 1.5.1. *If $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$, then $\text{Pol} = \text{Int}$.*

The proof can be found at the end of §4.2. The converse statement is not true, but “ $\text{Pol} = \text{Int}$ ” is equivalent to $U[[Z]]^{G_L,*} = o_L[[o_L]]$, where U is the o_L -submodule of $o_{\mathbb{C}_p}$ generated by $\{P_n(\Omega)\}_{n \geq 0}$. We have not been able to prove that $\text{Pol} = \text{Int}$, although we can show that Pol is p -adically dense in Int . Some numerical evidence indicates that $\text{Pol} = \text{Int}$ seems to hold: the details can be found in the Appendix by D. Crisan and J. Yang at the end of our paper.

We now explain how to compute the valuation of $P_n(\Omega)$ for certain n . The elements $z \in \mathfrak{m}_{\mathbb{C}_p}$ such that $G(z) = 0$ correspond to those locally L -analytic characters λ_z such that $\lambda_z(1) = 1$. Being locally L -analytic, they are necessarily trivial on an open subgroup of o_L , and correspond to certain torsion points of \mathcal{G} . We know the valuations of these torsion points, and this way we can determine the Newton polygon of $G(Z)$. Using this idea, we can prove the following. Let e be the ramification index of L/\mathbb{Q}_p . If $m \geq 0$, let $k_m = \lfloor (m-1)/e \rfloor$, so that $m = ek_m + r$ with $1 \leq r \leq e$. For $m \geq 0$, let $x_m = q^m/p^{k_m+1}$ (so that $x_0 = 1$ and $x_1 = q/p$). Write $m = en + r$ and let

$$y_0 = \frac{e}{p-1} - \frac{1}{q-1} \quad \text{and} \quad y_m = \frac{e}{p^n(p-1)} - \frac{r}{p^{n+1}} - \frac{1}{(q-1)p^{n+1}}.$$

Theorem 1.5.2. *For all $m \geq 0$, we have $\text{val}_\pi(P_{x_m}(\Omega)) = y_m$.*

For example, if $L = \mathbb{Q}_{p^2}$, then $\text{val}_p(P_{p^k}(\Omega)) = 1/p^{k-1}(q-1)$ for all $k \geq 0$.

1.6. Galois-continuous functions and the Katz map. Following Katz [19], we let $\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p})$ denote the o_L -module of Galois-continuous functions, namely those continuous functions $f : o_L \rightarrow o_{\mathbb{C}_p}$ such that $g(f(a)) = f(\tau(g) \cdot a)$ for all $a \in o_L$ and $g \in G_L$. If $P(T) \in L[T]$, then $a \mapsto P(a \cdot \Omega)$ is such a function. Let K be a closed subfield of \mathbb{C}_p containing L . The *dual Katz map* is the map $\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), o_K) \rightarrow o_K[[Z]]$ given by $\mu \mapsto \sum_{n \geq 0} \mu(P_n) \cdot Z^n$. Let

$o_K[[Z]]^{\psi_q\text{-int}}$ denote the set of $f(Z) \in o_K[[Z]]$ such that $\psi_q^n(f(Z)) \in o_K[[Z]]$ for all $n \geq 1$. Our main technical result is the following

Theorem 1.6.1. *Suppose that $L = \mathbb{Q}_{p^2}$.*

- (1) *The map $\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), o_K) \rightarrow o_K[[Z]]$ is injective.*
- (2) *Its image is equal to $o_K[[Z]]^{\psi_q\text{-int}}$.*

An important part of our paper is devoted to providing a proof of this theorem, which is completed at the end of §3.6. We note that Theorem 1.6.1 was stated by Katz at [19, p. 60], but he did not give a proof. The remarks contained in the last paragraph of [19, §IV] seem to indicate that his proof is different to ours.

The hardest part of the theorem is the claim concerning the image of \mathcal{K}^* . Note that when $L = \mathbb{Q}_{p^2}$, the dual of the p -divisible group attached to \mathcal{G} has dimension 1. Using this and Theorem 1.5.2 for $L = \mathbb{Q}_{p^2}$, we can prove (see Proposition 3.6.5) that every element of $o_\infty = o_{\mathbb{C}_p}^{\ker \tau}$ can be written as $\sum_{n \geq 0} \lambda_n \cdot P_n(\Omega)$ where $\lambda_n \in o_L$ and $\lambda_n \rightarrow 0$. This important ingredient of the proof of Theorem 1.6.1 is not known to be available if $L \neq \mathbb{Q}_{p^2}$.

1.7. Applications of the Katz isomorphism. Throughout this section, we assume that $L = \mathbb{Q}_{p^2}$ and $\pi = p$, so that $\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), o_K) \rightarrow o_K[[Z]]^{\psi_q\text{-int}}$ is an isomorphism. Let $L_\infty = \mathbb{C}_p^{\ker \tau}$ and $o_\infty = o_{\mathbb{C}_p}^{\ker \tau}$. Since $\pi = p$, L_∞ is also the completion of $L(\mathcal{G}[p^\infty])$.

Theorem 1.6.1 gives us an isomorphism $\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}), o_K) \rightarrow o_K[[Z]]^{\psi_q=0}$, and we have a natural isomorphism $\mathcal{C}_{\text{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}) \rightarrow o_\infty$. Applying this to $K = L$, we get the following result (Theorem 5.1.4), where $o_\infty^* = \text{Hom}_{o_L}(o_\infty, o_L)$:

Theorem 1.7.1. *The map \mathcal{K}^* gives rise to an isomorphism $o_\infty^* \simeq o_L[[Z]]^{\psi_q=0}$.*

Let $\Gamma_L^{\text{LT}} = \text{Gal}(L(\mathcal{G}[p^\infty])/L)$ and $\Gamma_{\mathbb{Q}_p}^{\text{cyc}} = \text{Gal}(\mathbb{Q}_p(\mu_{p^\infty})/\mathbb{Q}_p)$. In the cyclotomic setting, Perrin-Riou showed [25, Lemma 1.5] that $\mathbb{Z}_p[[Z]]^{\psi_p=0}$ is a free $\mathbb{Z}_p[[\Gamma_{\mathbb{Q}_p}^{\text{cyc}}]]$ -module of rank 1. She also raised the question of what happens in the present setting. Using Theorem 1.7.1, we show in Corollary 5.2.12 that $o_L[[Z]]^{\psi_q=0}$ is in fact *not* a free $o_L[[\Gamma_L^{\text{LT}}]]$ -module of rank 1.

We can also apply the isomorphism $\text{Hom}_{o_L}(o_\infty, o_K) \simeq o_K[[Z]]^{\psi_q=0}$ to $K = L_\infty$, and we get $\text{Hom}_{o_L}(o_\infty, o_\infty) \simeq o_\infty[[Z]]^{\psi_q=0}$. The natural action of G_L on the left is the twisted Galois action on the right. Since $\Lambda_L(\mathfrak{X}) = o_{\mathbb{C}_p}[[Z]]^{G_L,*} = o_\infty[[Z]]^{G_L,*}$, we get the following result (Theorem 5.1.6):

Theorem 1.7.2. *We have $\text{End}_{o_L}^{G_L}(o_\infty) \simeq \Lambda_L(\mathfrak{X})^{\psi_q=0}$.*

Recall that $o_L[o_L^\times] \subset \Lambda_L(\mathfrak{X})^{\psi_q=0}$. If $a \in o_L^\times$, then $\delta_a \in o_L[o_L^\times]$ acts on o_∞ by an element $g \in G_L$ such that $\tau(g) = a$. Since $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if $\Lambda_L(\mathfrak{X})^{\psi_q=0} = o_L[[o_L^\times]]$, we get the following criterion (Theorem 5.1.8):

Theorem 1.7.3. *We have $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if every continuous L -linear and G_L -equivariant map $f : L_\infty \rightarrow L_\infty$ comes from the Iwasawa algebra $L \otimes_{o_L} o_L[[\Gamma_L^{\text{LT}}]]$.*

In the cyclotomic case, Tate's normalized trace maps $T_n : \mathbb{Q}_p^{\text{cyc}} \rightarrow \mathbb{Q}_p(\mu_{p^n})$ are examples of continuous \mathbb{Q}_p -linear and $G_{\mathbb{Q}_p}$ -equivariant maps $f : \mathbb{Q}_p^{\text{cyc}} \rightarrow \mathbb{Q}_p^{\text{cyc}}$ that do not come from the Iwasawa algebra $L \otimes_{o_L} o_L[[\Gamma_{\mathbb{Q}_p}^{\text{cyc}}]]$. The lack of normalized trace maps in the Lubin–Tate setting is a source of many complications. In his PhD thesis, Fourquaux considered continuous L -linear and G_L -equivariant maps $f : L_\infty \rightarrow L_\infty$. We generalize some of Fourquaux's results: we prove in Proposition 5.1.13 that if $f \neq 0$ is such a map, then there exists $n \geq 0$ such that $f(L_\infty)$ contains a basis of the L_n -vector space $L_n[\log \Omega]$, where $L_n = L(\mathcal{G}[p^n])$. In particular, f necessarily has a very large image, so there can be no analogue of the equivariant trace maps T_n .

The Katz isomorphism also allows us to prove several results about the span of the polynomials P_n in $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$. Recall that by [28, Theorem 4.7], every Galois-continuous locally analytic function on o_L can be expanded as an overconvergent series in the P_n . One may then wonder about the existence of such an expansion for Galois-continuous functions. Let $\mathcal{C}^0(L)$ denote the set of sequences $\{\lambda_n\}_{n \geq 0}$ with $\lambda_n \in L$ and $\lambda_n \rightarrow 0$. The Katz isomorphism, and computations involving ψ_q , imply the following (Proposition 5.3.1, Corollary 5.3.4, and Corollary 5.3.9):

Theorem 1.7.4. *The map $\mathcal{C}^0(L) \rightarrow \mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$, given by*

$$\{\lambda_n\}_{n \geq 0} \mapsto \left[a \mapsto \sum_{n=0}^{\infty} \lambda_n \cdot P_n(a\Omega) \right]$$

is injective, has dense image, but is not surjective.

The same methods imply the following precise estimates for those elements of $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$ that are given by a polynomial function $a \mapsto Q(a\Omega)$ with $Q(T) \in L[T]$. See Proposition 5.3.6 and Corollary 5.3.12.

Theorem 1.7.5. *Assume that Z is a coordinate on \mathcal{G} such that $[p](Z) = Z^q + pZ$. Let $Q(T) \in L[T]$ be a polynomial such that $Q(a\Omega) \in o_{\mathbb{C}_p}$ for all $a \in o_L$, and write $Q(T) = \sum_{n=0}^{\deg Q} \lambda_n \cdot P_n(T)$.*

- (1) *We have $\lambda_n \in p^{-k} o_L$ if $n \leq q^k$.*
- (2) *For all k , there exists such a polynomial Q for which $\lambda_{q^k-1} = p^{-k}$.*

1.8. Other criteria. The following two criteria for our main question may be of interest.

Let $\partial : \mathbb{C}_p[[Z]] \rightarrow \mathbb{C}_p[[Z]]$ denote the invariant derivative $\partial = \log'_{\text{LT}}(Z)^{-1} \cdot d/dZ$. It does not commute with the twisted action of G_L , but $D = \Omega^{-1} \cdot \partial$ does. We get a map $D : \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}) \rightarrow \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$ that does not preserve $\Lambda_{\mathbb{C}_p}(\mathfrak{X})$ if $L \neq \mathbb{Q}_p$ since $\text{val}_p(\Omega^{-1}) < 0$. Note that $D(\delta_a) = a \cdot \delta_a$ if $a \in o_L$, so that D does preserve $o_L[[o_L]]$. We have the following result.

Theorem 1.8.1. *If $L = \mathbb{Q}_{p^2}$, then $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if we have $D^{q-1}(\Lambda_L(\mathfrak{X})) \subset \Lambda_L(\mathfrak{X})$.*

This Theorem follows from Theorem 1.4.1 and the following result, which is inspired by computations of Katz: assume that $L = \mathbb{Q}_{p^2}$ and that $\pi = p$. Let $\lambda = \Omega^{q-1}/p(q-1)! \in o_{\mathbb{C}_p}^\times$. If $f(Z) \in o_{\mathbb{C}_p}[[Z]]$, then

$$\varphi\psi_q(f) - \lambda \cdot D^{q-1}(f) \in o_{\mathbb{C}_p}[[Z]].$$

Here is another result concerning our main question. It says that if the answer is yes for a finite extension L/K , then the answer is also yes for K .

Theorem 1.8.2. *If L/K is finite and if $\Lambda_L(\mathfrak{X}_L) = o_L[[o_L]]$, then $\Lambda_K(\mathfrak{X}_K) = o_K[[o_K]]$.*

1.9. Acknowledgements. This paper grew out of a project started with Peter Schneider. The authors are very grateful to him for numerous discussions, interesting insights (in particular, considering the Katz isomorphism), and several invitations to Münster. Several results in this paper were obtained in collaboration with him. L.B. also thanks Pierre Colmez for some discussions about the main problem of this paper.

2. THE CHARACTER VARIETY

2.1. Notation. Let $\mathbb{Q}_p \subseteq L \subset \mathbb{C}_p$ be a field of finite degree d over \mathbb{Q}_p , o_L the ring of integers of L , $\pi \in o_L$ a fixed prime element, $k_L = o_L/\pi o_L$ the residue field, $q := |k_L|$ and e the absolute ramification index of L . We always use the absolute value $|\cdot|$ on \mathbb{C}_p which is normalized by $|p| = p^{-1}$. We let $G_L := \text{Gal}(\bar{L}/L)$ denote the absolute Galois group of L . Throughout our coefficient field K is a complete intermediate extension $L \subseteq K \subseteq \mathbb{C}_p$.

2.2. The p -adic Fourier transform. We are interested in the *character variety* \mathfrak{X} of the L -analytic commutative group $(o_L, +)$. We refer to [28, §2] for a precise definition, but recall that \mathfrak{X} is a rigid analytic variety defined over L , whose set of K -points (for K a field extension of L complete with respect to a non-archimedean absolute value extending the one on L) is the group $\mathfrak{X}(K)$ of K -valued characters $\chi : (o_L, +) \rightarrow (K^\times, \times)$ that are also L -analytic functions:

$$\mathfrak{X}(K) := \{f \in C^{L-\text{an}}(o_L, K) : f(a+b) = f(a)f(b) \text{ for all } a, b \in o_L\}.$$

Here $C^{L-\text{an}}(o_L, K)$ is the space of locally L -analytic K -valued functions on o_L . Let $D^{L-\text{an}}(o_L, K)$ be the K -algebra of locally L -analytic distributions on o_L , defined in [29, §2]. One of the main results of p -adic Fourier Theory — [28, Theorem 2.3] — tells us that there is a canonical isomorphism

$$\mathcal{F} : D^{L-\text{an}}(o_L, K) \rightarrow \mathcal{O}(\mathfrak{X} \times_L K)$$

called the *p -adic Fourier Transform*. This isomorphism is determined by

$$\mathcal{F}(\lambda)(\chi) = \lambda(\chi) \text{ for all } \lambda \in D^{L-\text{an}}(o_L, K), \chi \in \mathfrak{X}(K).$$

Since \mathfrak{X} is a rigid L -analytic variety, we have at our disposal the subalgebra $\mathcal{O}^\circ(\mathfrak{X})$ of $\mathcal{O}(\mathfrak{X})$ consisting of globally-defined, rigid analytic functions on \mathfrak{X} that are *power-bounded* — see [7, §1.2.5].

Definition 2.2.1. Write $\Lambda(\mathfrak{X}) := \mathcal{O}^\circ(\mathfrak{X})$.

The functorial definition of the character variety does not shed much light on its internal structure. It turns out that the base change $\mathfrak{X} \times_L K$ is isomorphic to the rigid analytic open unit disc over K , *provided* the field K is large enough. This isomorphism is obtained with the help of *Lubin-Tate formal groups* and their associated *p-divisible groups*.

2.3. Lubin-Tate formal groups. Let Z be an indeterminate and let

$$\mathcal{F}_\pi := (\pi Z + Z^2 o_L[[Z]]) \cap (Z^q + \pi o_L[[Z]])$$

be the set of possible *Frobenius power series*. Recall [21, Theorem 8.1.1]¹ that for every Frobenius power series $\varphi(Z) \in \mathcal{F}_\pi$, there is a unique formal group law $F_{\varphi(Z)} = Z_1 \oplus Z_2 \in o_L[[Z_1, Z_2]]$ such that $\varphi(Z)$ is an endomorphism of $F_{\varphi(Z)}$. Since we have fixed a coordinate Z on the power series ring $o_L[[Z]]$, this formal group law defines a *formal group*² (\mathcal{G}, \oplus) on the underlying formal affine scheme $\mathrm{Spf} o_L[[Z]]$, where we give $o_L[[Z]]$ the Z -adic topology. This formal group is called a *Lubin-Tate formal group*. Up to isomorphism of formal groups, it does not depend on the choice of the Frobenius power series $\varphi(Z)$, however it does depend on the choice of π . The base change of \mathcal{G} to the completion $\widehat{L}^{\mathrm{ur}}$ of the maximal unramified extension L^{ur} of L does not even depend on the choice of π .

The Lubin-Tate formal group \mathcal{G} is in fact a *formal o_L -module*. This means that there is a ring homomorphism $o_L \rightarrow \mathrm{End}(\mathcal{G})$, $a \mapsto [a](Z) \in o_L[[Z]]$, such that $[a](Z) \equiv aZ \pmod{Z^2 o_L[[Z]]}$ for all $a \in o_L$. In other words, the formal group \mathcal{G} admits an action of o_L by endomorphisms of formal groups, in such a way that the differential of this action at the identity element 1 of \mathcal{G} agrees with the natural o_L -action on the cotangent space of \mathcal{G} at 1. The action of $\pi \in o_L$ is given by the power series $[\pi](Z) = \varphi(Z)$.

2.4. A review of p -divisible groups. In his seminal paper [32], Tate introduced *p-divisible groups* and considered their relation to formal groups. Here we review some of his fundamental theorems.

Let R be a commutative base ring and let $\Gamma = (\mathrm{Spf} \mathcal{A}, *)$ be a commutative formal group over R where $\mathcal{A} = R[[X_1, \dots, X_d]]$ is a power series ring in d variables over R . Then we can associate with Γ the *p-divisible group* $\Gamma(p) = (\Gamma(p)_n, i_n)$ over R where $\Gamma(p)_n := \Gamma[p^n]$ is the subgroup of elements of Γ killed by p^n . More precisely, let $\psi : \mathcal{A} \rightarrow \mathcal{A}$ be the continuous R -algebra homomorphism which corresponds to multiplication by p on Γ and let J_n be the ideal $\mathcal{A}\psi^n(X_1) + \dots + \mathcal{A}\psi^n(X_d)$ of \mathcal{A} ; then \mathcal{A}/J_n is a Hopf algebra over R free of finite rank over R , and $\Gamma(p)_n = \mathrm{Spec}(\mathcal{A}/J_n)$ is the corresponding commutative finite flat group scheme over R . The closed immersions $i_n : \Gamma(p)_n \rightarrow \Gamma(p)_{n+1}$ are obtained from the R -algebra surjections $\mathcal{A}/J_{n+1} \twoheadrightarrow \mathcal{A}/J_n$.

¹Note that what Lang calls a formal group should really be called a *formal group law*.

²a group object in the category of formal schemes over $\mathrm{Spf} o_L$

Theorem 2.4.1 (§2.2, Proposition 1 [32]). *Let R be a complete Noetherian local ring whose residue field k is of characteristic $p > 0$. Then $\Gamma \mapsto \Gamma(p)$ is an equivalence between the category of divisible commutative formal groups over R and the category of connected p -divisible groups over R .*

Recall that the formal group Γ is said to be *divisible* if \mathcal{A}/J_1 is finitely generated as an R -module, and a p -divisible group (Γ_n, i_n) is said to be *connected* if every finite flat group scheme Γ_n is a connected scheme.

Remark 2.4.2. The fact that the functor $\Gamma \mapsto \Gamma(p)$ is fully faithful holds in greater generality: if R is *any* commutative ring and G, H are divisible formal groups defined over R such that $\mathcal{O}(G)$ and $\mathcal{O}(H)$ are power series rings in finitely many variables over R , then the natural map

$$\mathrm{Hom}_{R\text{-fgp}}(G, H) \rightarrow \mathrm{Hom}_{p\text{-div}}(G(p), H(p))$$

is a bijection.

Now we specialise to the case where R is our complete discrete valuation ring \mathcal{o}_L . The *Tate module* associated to a p -divisible group $\Gamma = (\Gamma_n, i_n)$ is by definition

$$T(\Gamma) := \varprojlim \Gamma_n(\bar{L})$$

where \bar{L} is the algebraic closure of L , $\Gamma_n(\bar{L}) = \mathrm{Hom}_{\mathcal{o}_L\text{-alg}}(\mathcal{O}(\Gamma_n), \bar{L})$ is the set of \bar{L} -points of Γ_n , and the connecting maps in the inverse limit are induced by the multiplication-by- p -maps $j_n : \Gamma_{n+1} \rightarrow \Gamma_n$. By functoriality, the Tate module $T(\Gamma)$ carries a natural action of the absolute Galois group $G_L = \mathrm{Gal}(\bar{L}/L)$, making $T(\Gamma)$ into a continuous \mathbb{Z}_p -linear representation of G_L of rank equal to the *height* h of Γ . Remarkably, it turns out that this Galois representation completely determines the p -divisible group Γ . More precisely, we have the following

Theorem 2.4.3 (§4.2, Corollary 1 [32]). *The functor $\Gamma \mapsto T(\Gamma)$ is a fully faithful embedding of the category of p -divisible groups over \mathcal{o}_L into the category of finite rank \mathbb{Z}_p -linear continuous representations of G_L .*

2.5. Cartier duality for p -divisible groups. The category of commutative finite flat group R -schemes admits a duality called *Cartier duality*: if G is a commutative finite flat group scheme over R , then its Cartier dual is defined by $G^\vee = \mathrm{Spec}(\mathcal{O}(G)^*)$ where $\mathcal{O}(G)^* := \mathrm{Hom}_R(\mathcal{O}(G), R)$ is the R -linear dual of the coordinate ring $\mathcal{O}(G)$. The group structure on G^\vee is obtained by dualising the multiplication map on $\mathcal{O}(G)$ and the scheme structure on G^\vee is obtained by dualising the comultiplication map on $\mathcal{O}(G)$ encoding the group structure on G .

Tate shows in [32, §2.3] that Cartier duality extends naturally to a duality $\Gamma \mapsto \Gamma^\vee$ on the category of p -divisible groups. He also shows in [32, §4] that when $R = \mathcal{o}_L$, the Tate-module functor to Galois representations converts Cartier duality into what is now called *Tate duality* on Galois representations,

namely $V \mapsto \operatorname{Hom}(V, \mathbb{Z}_p(1))$. In other words, there is a natural isomorphism of continuous G_L -representations on finite rank \mathbb{Z}_p -modules

$$T(\Gamma^\vee) \cong \operatorname{Hom}_{\mathbb{Z}_p}(T(\Gamma), \mathbb{Z}_p(1))$$

where $\mathbb{Z}_p(1) := T(\widehat{\mathbb{G}}_m(p))$ is the Tate module associated to the formal multiplicative group $\widehat{\mathbb{G}}_m$, the formal completion at the identity of the group scheme $\mathbb{G}_m := \operatorname{Spec} o_L[T, T^{-1}]$.

2.6. The character $\tau : G_L \rightarrow o_L^\times$ and the period Ω . We return to the Lubin-Tate formal group \mathcal{G} as in §2.3, which is easily seen to be divisible. Because \mathcal{G} is a formal o_L -module, the functoriality of $T(-)$ implies that the Tate module $T(\mathcal{G}(p))$ of the p -divisible group $\mathcal{G}(p)$ associated with \mathcal{G} is actually an o_L -module. It is a fundamental fact due to Lubin and Tate — see [23, Theorem 2] — that $T(\mathcal{G}(p))$ is a free o_L -module of rank one. Since o_L is itself a free \mathbb{Z}_p -module of rank $d = [L : \mathbb{Q}_p]$, it follows that the underlying \mathbb{Z}_p -module of $T(\mathcal{G}(p)^\vee) \cong \operatorname{Hom}_{\mathbb{Z}_p}(T(\mathcal{G}(p)), \mathbb{Z}_p)$ is free of rank d as a \mathbb{Z}_p -module as well. Since it is also an o_L -module by the functoriality of $\operatorname{Hom}_{\mathbb{Z}_p}(-, \mathbb{Z}_p)$, we see that $T(\mathcal{G}(p)^\vee)$ is also a free o_L -module of rank 1.

On the way to his proof of Theorem 2.4.3, Tate explains how to compute $T(\mathcal{G}(p)^\vee)$: using Cartier duality, on [32, p. 177] he obtains a natural isomorphism of abelian groups

$$(1) \quad T(\mathcal{G}(p)^\vee) \cong \operatorname{Hom}_{p\text{-div}/o_{\mathbb{C}_p}}(\mathcal{G}(p) \times_{o_L} o_{\mathbb{C}_p}, \widehat{\mathbb{G}}_m(p) \times_{o_L} o_{\mathbb{C}_p}).$$

On the other hand, applying Remark 2.4.2 with $R = o_{\mathbb{C}_p}$, we see that the natural map

$$(2) \quad \begin{aligned} \operatorname{Hom}_{\operatorname{fgrp}/o_{\mathbb{C}_p}}(\mathcal{G} \times_{o_L} o_{\mathbb{C}_p}, \widehat{\mathbb{G}}_m \times_{o_L} o_{\mathbb{C}_p}) \\ \rightarrow \operatorname{Hom}_{p\text{-div}/o_{\mathbb{C}_p}}(\mathcal{G}(p) \times_{o_L} o_{\mathbb{C}_p}, \widehat{\mathbb{G}}_m(p) \times_{o_L} o_{\mathbb{C}_p}) \end{aligned}$$

is a bijection. As a consequence, we see that $\operatorname{Hom}_{\operatorname{fgrp}/o_{\mathbb{C}_p}}(\mathcal{G} \times_{o_L} o_{\mathbb{C}_p}, \widehat{\mathbb{G}}_m \times_{o_L} o_{\mathbb{C}_p})$ is free of rank 1 as an o_L -module.

Definition 2.6.1.

- (1) We fix a generator t'_o for $T(\mathcal{G}(p)^\vee)$ as an o_L -module.
- (2) We let $F_{t'_o}$ be the generator for the o_L -module

$$\operatorname{Hom}_{\operatorname{fgrp}/o_{\mathbb{C}_p}}(\mathcal{G} \times_{o_L} o_{\mathbb{C}_p}, \widehat{\mathbb{G}}_m \times_{o_L} o_{\mathbb{C}_p}),$$

which corresponds to t'_o along the isomorphism

$$T(\mathcal{G}(p)^\vee) \xrightarrow{\cong} \operatorname{Hom}_{\operatorname{fgrp}/o_{\mathbb{C}_p}}(\mathcal{G} \times_{o_L} o_{\mathbb{C}_p}, \widehat{\mathbb{G}}_m \times_{o_L} o_{\mathbb{C}_p})$$

obtained by combining (1) and (2).

- (3) We let $\tau : G_L \rightarrow o_L^\times$ be the character afforded by the free rank 1 o_L -module $T(\mathcal{G}(p)^\vee)$:

$$\sigma(t'_o) = \tau(\sigma)t'_o \quad \text{for all } \sigma \in G_L.$$

The morphism of formal groups $F_{t'_o} : \mathcal{G} \times_{o_L} o_{\mathbb{C}_p} \rightarrow \widehat{\mathbb{G}}_m \times_{o_L} o_{\mathbb{C}_p}$ is an element of

$$F_{t'_o}(Z) \in \mathcal{O}(\mathcal{G} \times_{o_L} o_{\mathbb{C}_p}) = o_{\mathbb{C}_p}[[Z]].$$

Then $1 + F_{t'_o}(Z)$ is “grouplike” in the topological Hopf algebra $o_{\mathbb{C}_p}[[Z]]$: it satisfies the relation

$$1 + F_{t'_o}(Z_1 \oplus Z_2) = (1 + F_{t'_o}(Z_1))(1 + F_{t'_o}(Z_2)).$$

When we further base change the formal group $\mathcal{G} \times_{o_L} o_{\mathbb{C}_p}$ to \mathbb{C}_p , it becomes isomorphic to the additive formal group. It follows from this that $\log F_{t'_o}(Z)$ is necessarily “primitive” in the topological Hopf algebra $\mathbb{C}_p[[Z]]$: it satisfies the relation

$$(3) \quad \log(1 + F_{t'_o}(Z_1 \oplus Z_2)) = \log(1 + F_{t'_o}(Z_1)) + \log(1 + F_{t'_o}(Z_2)).$$

Since the *logarithm* $\log_{\text{LT}}(Z)$ of the formal group \mathcal{G} spans the space of primitive elements in $\mathbb{C}_p[[Z]]$, it follows that there exists a unique element $\Omega \in \mathbb{C}_p$ such that

$$1 + F_{t'_o}(Z) = \exp(\Omega \log_{\text{LT}}(Z)).$$

Definition 2.6.2. The element Ω is called the *period* of the dual p -divisible group $\mathcal{G}(p)^\vee$.

Let $I_L \subseteq G_L$ denote the inertia subgroup.

Lemma 2.6.3. *If $L \neq \mathbb{Q}_p$, then the character $\tau : I_L \rightarrow o_L^\times$ has an open image.*

Proof. Let χ_π be the character describing the G_L -action on the Tate module T of \mathcal{G} . By local class field theory we know that on I_L , $\text{Norm}_{L/\mathbb{Q}_p} \circ \chi_\pi = \chi_{\text{cyc}}$, the cyclotomic character. From Definition 2.6.1(2), we have $\tau = \chi_\pi^{-1} \cdot \chi_{\text{cyc}}$. Hence $\tau : I_L \rightarrow o_L^\times$ is the composition of the surjective map $\chi_\pi : I_L \rightarrow o_L^\times$ and of the map given by $x \mapsto \prod_{\sigma: L \rightarrow \overline{\mathbb{Q}_p}, \sigma \neq \text{Id}} \sigma(x)$.

On the Lie algebra L of o_L^\times , the derivative of the above map is given by $U = \text{Tr}_{L/\mathbb{Q}_p} - \text{Id}$. We prove that $U : L \rightarrow L$ is injective, hence surjective, which implies the lemma. If $U(x) = 0$, then $x = (U + \text{Id})x = \text{Tr}_{L/\mathbb{Q}_p}(x) \in \mathbb{Q}_p$ and hence $U(x) = ([L : \mathbb{Q}_p] - 1)x$ so that $x = 0$. \square

For future use, we record here the more precise result (pointed out to us by B. Xie) which gives a sufficient criterion for τ to be surjective.

Lemma 2.6.4. *If $d-1$ and $(p-1)p$ are coprime, then $\tau : I_L \rightarrow o_L^\times$ is surjective.*

Proof. Since $\tau = \chi_\pi^{-1} \cdot \chi_{\text{cyc}}$ and $\chi_{\text{cyc}} = \text{Norm}_{L/\mathbb{Q}_p} \circ \chi_\pi$, we have

$$\tau(g) = \chi_\pi(g)^{-1} \text{Norm}_{L/\mathbb{Q}_p}(\chi_\pi(g)) \quad \text{for any } g \in I_L.$$

Note also that the restriction to I_L of the totally ramified surjective character $\chi_\pi \rightarrow o_L^\times$ is still surjective. Let now $u \in o_L^\times$ be any fixed element.

We first show that there is an $a \in \mathbb{Z}_p^\times$ such that $a^{d-1} = \text{Norm}_{L/\mathbb{Q}_p}(u)$. Let $v := \text{Norm}_{L/\mathbb{Q}_p}(u)$ and let \bar{v} denote its image in \mathbb{F}_p^\times . By our assumption the

polynomial $Z^{d-1} - \bar{v}$ is separable over \mathbb{F}_p and has a root in \mathbb{F}_p^\times . Hence Hensel's Lemma implies that the polynomial $Z^{d-1} - v$ has a root $a \in \mathbb{Z}_p^\times$.

Choosing now a $g \in I_L$ such that $\chi_\pi(g) = au^{-1}$ we deduce that

$$\tau(g) = (au^{-1})^{-1} \text{Norm}_{L/\mathbb{Q}_p}(au^{-1}) = ua^{-1}a^d \text{Norm}_{L/\mathbb{Q}_p}(u^{-1}) = u. \quad \square$$

2.7. The Amice-Katz transform. With the period $\Omega \in \mathbb{C}_p$ in hand, we now recall some constructions from p -adic Fourier Theory [28]. For each $a \in o_L$, define

$$\Delta_a := 1 + F_{at'_o}(Z) = \exp(a\Omega \log_{LT}(Z)) \in \mathbb{C}_p[[Z]]^\times.$$

The map $(o_L, +) \rightarrow (\mathbb{C}_p[[Z]]^\times, \times)$ which sends $a \in o_L$ to Δ_a is a group homomorphism. The fundamental property of these power series is that their coefficients all lie in $o_{\mathbb{C}_p}$:

$$\Delta_a \in o_{\mathbb{C}_p}[[Z]]^\times \quad \text{for all } a \in o_L.$$

This follows from the fact that for each $a \in o_L$, $F_{at'_o} : \mathcal{G} \times_{o_L} o_{\mathbb{C}_p} \rightarrow \widehat{\mathbb{G}}_m \times_{o_L} o_{\mathbb{C}_p}$ is a homomorphism of formal groups defined over $o_{\mathbb{C}_p}$; see also [28, Lemma 4.2(5)].

Definition 2.7.1.

- (1) Let L_∞ be the closure in \mathbb{C}_p of the subfield $L(\Omega)$ of \mathbb{C}_p generated by L and Ω .
- (2) Let $L_\tau := L_\infty \cap \bar{L}$.
- (3) Let $o_\infty := L_\infty \cap o_{\mathbb{C}_p}$.
- (4) Let $o_\tau := L_\tau \cap o_{\mathbb{C}_p}$.

Lemma 2.7.2. *We have $L_\infty = \mathbb{C}_p^{\ker \tau}$ and $o_\infty = o_{\mathbb{C}_p}^{\ker \tau}$.*

Proof. From the relation appearing in Definition 2.6.1(3), we deduce

$$\sigma(\Omega) = \tau(\sigma)\Omega \quad \text{for all } \sigma \in G_L.$$

This immediately implies that $L_\infty \subseteq \mathbb{C}_p^{\ker \tau}$. Let $H := \text{Gal}(\bar{L}/L_\tau)$, a closed subgroup of G_L , and let $g \in H$. Then g extends to a unique continuous L_τ -linear automorphism g of \mathbb{C}_p . Now L_∞ is the closure of L_τ in \mathbb{C}_p , so g fixes $\Omega \in L_\infty$. Hence $\tau(g) = 1$ by the above relation. Hence $H \leq \ker \tau$ which implies that $\mathbb{C}_p^{\ker \tau} \leq \mathbb{C}_p^H$. But \bar{L}^H is dense in \mathbb{C}_p^H by the Ax-Sen-Tate theorem, [9, Proposition 2.1.2], and $\bar{L}^H = L_\tau$ by infinite Galois theory. Hence L_τ is dense in \mathbb{C}_p^H , so \mathbb{C}_p^H is contained in the closure of L_τ in \mathbb{C}_p , namely L_∞ . Hence $\mathbb{C}_p^{\ker \tau} \leq L_\infty$.

The second statement follows from the first by intersecting $L_\infty = \mathbb{C}_p^{\ker \tau}$ with $o_{\mathbb{C}_p}$. \square

It is clear from the definition of Δ_a that in fact $\Delta_a \in o_\infty[[Z]]^\times$ for all $a \in o_L$.

Definition 2.7.3. We write $o_L[[o_L]]$ for the completed group ring of the abelian group o_L with coefficients in o_L . The *Amice-Katz transform* is the unique extension to a continuous o_L -algebra homomorphism

$$\mu : o_L[[o_L]] \rightarrow \mathcal{O}(\mathcal{G} \times_{o_L} o_\infty) = o_\infty[[Z]]$$

of the group homomorphism $o_L \rightarrow o_{\mathbb{C}_p}[[Z]]^\times$ which sends $a \in o_L$ to $\Delta_a \in o_\infty[[Z]]^\times$.

2.8. The Schneider-Teitelbaum uniformisation. At this point, rigid analytic geometry enters the picture. Let \mathbf{B} be the rigid L_∞ -analytic open disc of radius one, with local coordinate Z . By definition, \mathbf{B} is the colimit of the rigid L_∞ -analytic closed discs $\mathbf{B}(r)$ of radius $r < 1$, as $r \in |L_\infty^\times|$ approaches 1 from below:

$$\mathbf{B} = \operatorname{colim}_{r < 1} \mathbf{B}(r), \quad \mathbf{B}(r) = \operatorname{Sp} L_\infty \langle Z/\dot{r} \rangle$$

where \dot{r} is any choice of an element of L_∞^\times such that $|\dot{r}| = r$. Choosing, for convenience, any strictly increasing sequence $r_1 < r_2 < r_3 < \dots$ of real numbers in $|L_\infty| \cap (0, 1)$ approaching 1 from below, we have a descending chain of L_∞ -algebras, each one containing $o_\infty[[Z]]$:

$$\begin{aligned} L_\infty \langle Z/\dot{r}_1 \rangle \supsetneq L_\infty \langle Z/\dot{r}_2 \rangle \supsetneq L_\infty \langle Z/\dot{r}_3 \rangle \supsetneq \dots \supsetneq \bigcap_{n=1}^{\infty} L_\infty \langle Z/\dot{r}_n \rangle \\ = \mathcal{O}(\mathbf{B}) \supseteq o_\infty[[Z]] \otimes_{o_L} L. \end{aligned}$$

With this notation in place, it follows from one of Schneider-Teitelbaum's main results, [28, Theorem 3.6], that the o_L -algebra homomorphism $\mu : o_L[[o_L]] \rightarrow o_\infty[[Z]]$ extends to a continuous *isomorphism* of L -Fréchet algebras

$$\mu_{\text{rig}} : D^{L\text{-an}}(o_L, L_\infty) \xrightarrow{\cong} \mathcal{O}(\mathbf{B})$$

which makes the following diagram commutative:

$$\begin{array}{ccc} o_L[[o_L]] \otimes_{o_L} L & \xrightarrow{\mu} & o_\infty[[Z]] \otimes_{o_L} L \\ \downarrow & & \downarrow \\ D^{L\text{-an}}(o_L, L_\infty) & \xrightarrow[\mu_{\text{rig}}]{\cong} & \mathcal{O}(\mathbf{B}) \end{array}$$

The vertical arrow on the left is the natural restriction map $o_L[[o_L]] \otimes_{o_L} L$ into $D^{L\text{-an}}(o_L, L)$, witnessing the fact that every locally L -analytic function on o_L is continuous, and hence that every continuous distribution on o_L restricts to a locally L -analytic distribution on o_L ; see [29] for more details. The vertical arrow on the right is the inclusion $o_\infty[[Z]] \otimes_{o_L} L \subset \mathcal{O}(\mathbf{B})$ from the above discussion. Combining the isomorphism μ_{rig} with the Fourier transform $\mathcal{F} : D^{L\text{-an}}(o_L, L_\infty) \rightarrow \mathcal{O}(\mathfrak{X} \times_L L_\infty)$, we obtain an isomorphism of L_∞ -Fréchet algebras

$$\mu_{\text{rig}} \circ \mathcal{F}^{-1} : \mathcal{O}(\mathfrak{X} \times_L L_\infty) \xrightarrow{\cong} \mathcal{O}(\mathbf{B}).$$

Since $\mathfrak{X} \times_L L_\infty$ and \mathbf{B} are both Stein rigid analytic varieties over L_∞ , this isomorphism determines, and is completely determined by, an isomorphism

$$\kappa := \mathrm{Sp}(\mu_{\mathrm{rig}} \circ \mathcal{F}^{-1}) : \mathbf{B} \xrightarrow{\cong} \mathfrak{X} \times_L L_\infty.$$

This is a version of [28, Theorem 3.6]: the base-change of the character variety \mathfrak{X} to L_∞ is isomorphic to the rigid L_∞ -analytic open disc of radius one, so κ can be viewed as giving a *uniformisation* of $\mathfrak{X} \times_L L_\infty$ by \mathbf{B} . Schneider and Teitelbaum also show that the morphism κ is given on \mathbb{C}_p -points by the following rule: for each $z \in \mathbf{B}(\mathbb{C}_p)$ we can evaluate the power series $\Delta_a \in o_\infty[[Z]]$ at $Z = z$ to obtain an element $\Delta_a(z) \in o_{\mathbb{C}_p}^\times$, and the locally L -analytic character $\kappa(z) : o_L \rightarrow \mathbb{C}_p$ is given by

$$\kappa(z)(a) = \Delta_a(z) \quad \text{for all } a \in o_L.$$

2.9. $\Lambda_L(\mathfrak{X})$ and the twisted G_L -action on $\mathbb{C}_p[[Z]]$. It is natural to enquire, in the light of the Schneider-Teitelbaum isomorphism

$$\kappa : \mathbf{B} \xrightarrow{\cong} \mathfrak{X} \times_L L_\infty$$

how far the character variety \mathfrak{X} is itself from being isomorphic to an open rigid L -analytic unit disc. For general reasons, $\mathfrak{X} \times_L L_\infty$ carries a natural action of the Galois group G_L , acting on the second factor, giving an isomorphism of L -Fréchet algebras

$$\mathcal{O}(\mathfrak{X}) \cong \mathcal{O}(\mathfrak{X} \times_L L_\infty)^{G_L}.$$

Definition 2.9.1. The *twisted G_L -action* on $\mathcal{O}(\mathbf{B})$ is given as follows:

$$\sigma * F(Z) := (\sigma F)([\tau(\sigma)^{-1}](Z)) \quad \text{for all } F(Z) \in \mathcal{O}(\mathbf{B}), \sigma \in G_L.$$

Here $F \mapsto \sigma F$ is the “coefficient-wise” G_L -action on $\mathbb{C}_p[[Z]] \supset \mathcal{O}(\mathbf{B})$, given explicitly by $\sigma(\sum_{n=0}^\infty a_n Z^n) = \sum_{n=0}^\infty \sigma(a_n) Z^n$ for all $\sigma \in G_L$.

Schneider and Teitelbaum showed that this twisted G_L -action on $\mathcal{O}(\mathbf{B})$ in fact comes from the following twisted G_L -action on the set of \mathbb{C}_p -points $\mathbf{B}(\mathbb{C}_p)$:

$$\sigma * z = \kappa^{-1}(\sigma \circ \kappa(z)) \quad \text{for all } z \in \mathbf{B}(\mathbb{C}_p), \sigma \in G_L.$$

From the proof of [28, Corollary 3.8], we can also deduce the following

Proposition 2.9.2. *The algebra isomorphism*

$$\kappa^* = \mu_{\mathrm{rig}} \circ \mathcal{F}^{-1} : \mathcal{O}(\mathfrak{X} \times_L L_\infty) \xrightarrow{\cong} \mathcal{O}(\mathbf{B})$$

is equivariant with respect to the natural G_L -action on the source, and the twisted G_L -action on the target.

Corollary 2.9.3. *The map μ_{rig} restricts to give an isomorphism of o_L -algebras*

$$(\mu_{\mathrm{rig}} \circ \mathcal{F}^{-1})^\circ : \mathcal{O}^\circ(\mathfrak{X}) \xrightarrow{\cong} o_\infty[[Z]]^{G_L,*}.$$

Proof. Applying the functor \mathcal{O}° to the isomorphism of rigid L_∞ -analytic varieties $\kappa : \mathbf{B} \rightarrow \mathfrak{X} \times_L L_\infty$, we see that $\mu_{\text{rig}} \circ \mathcal{F}^{-1}$ restricts to an o_∞ -algebra isomorphism

$$\mathcal{O}(\mathfrak{X} \times_L L_\infty)^\circ \xrightarrow{\cong} \mathcal{O}(\mathbf{B})^\circ.$$

It is well known that $\mathcal{O}(\mathbf{B})^\circ = o_\infty[[Z]]$ and that $\Lambda_L(\mathfrak{X}) = \mathcal{O}(\mathfrak{X})^\circ = (\mathcal{O}(\mathfrak{X} \times_L L_\infty)^\circ)^{G_L}$. The result follows by passing to G_L -invariants and applying Proposition 2.9.2. \square

Consequently, the image of the Amice-Katz transform $\mu : o_L[[o_L]] \rightarrow o_\infty[[Z]]$ lands in the subring of twisted G_L -invariants. One of our main goals in this paper is to study the following

Question 2.9.4. Is the Amice-Katz transform $\mu : o_L[[o_L]] \rightarrow o_\infty[[Z]]^{G_L,*}$ an isomorphism?

2.10. Some properties of $\Lambda_L(\mathfrak{X})$. In this section, we identify (through the LT-isomorphism) the ring $\Lambda_L(\mathfrak{X}) = \mathcal{O}(\mathfrak{X})^\circ$ with the ring $o_\infty[[Z]]^{G_L,*}$. From [5] we know that $\Lambda_L(\mathfrak{X})$ is an integral domain and that the norm $\| \cdot \|_{\mathfrak{X}} = \| \cdot \|_1$ on $\Lambda_L(\mathfrak{X})$ is multiplicative. Let k_K denote the residue field of K .

Lemma 2.10.1. *If $L \neq \mathbb{Q}_p$ and if K is a finite extension of L , then $\bar{k}[[Z]]^{G_K,*} = k_K$.*

Proof. If $g \in I_K$, then g acts trivially on \bar{k} , so that the $G_{L,*}$ action of $g \in I_K$ on $\bar{k}[[Z]]$ is given by $g : \sum_{n \geq 0} a_n Z^n \mapsto \sum_{n \geq 0} a_n ([\tau(g)^{-1}]Z)^n$. The character $\tau : I_K \rightarrow o_L^\times$ has an open image by Lemma 2.6.3. This image therefore contains $\chi_\pi(I_M)$ where $M \subset L_\infty$ is some finite extension of L , and $\bar{k}[[Z]]^{I_K,*} = \bar{k}[[Z]]^{I_M}$ where I_M acts on $\bar{k}[[Z]]$ via $g : \sum_{n \geq 0} a_n Z^n \mapsto \sum_{n \geq 0} a_n ([\chi_\pi(g)]Z)^n$. We know from the theory of the field of norms that $\bar{k}[[Z]]$ with that action of I_M embeds into $\tilde{\mathbf{E}}^+ \simeq \varprojlim_{(-)^q} o_{\mathbb{C}_p}$ in an I_M -equivariant way. Let $P := \mathbb{C}_p^{I_M}$. We have $(\tilde{\mathbf{E}}^+)^{I_M} \simeq \varprojlim_{(-)^q} o_P = \bar{k}$ since P/\mathbb{Q}_p is finitely ramified. Hence $\bar{k}[[Z]]^{I_M} = \bar{k}$ and $\bar{k}[[Z]]^{I_K,*} = \bar{k}$. The Lemma then follows from the fact that on \bar{k} , the twisted G_L -action coincides with the usual G_L -action, so that $\bar{k}^{G_K,*} = k_K$. \square

Let χ_{triv} denote the character $o_L \rightarrow \mathbb{C}_p^\times$ given by $\chi_{\text{triv}}(a) = 1$ for all $a \in o_L$. Note that $\chi_{\text{triv}} = \kappa(0)$. We have a surjective map $\Lambda_L(\mathfrak{X}) \rightarrow k$ given by $f \mapsto f(\chi_{\text{triv}}) \bmod \mathfrak{m}_L$. Its kernel $\mathfrak{m}(\mathfrak{X}) := \{f \in \Lambda_L(\mathfrak{X}) : f(\chi_{\text{triv}}) \in \mathfrak{m}_L\}$ is a maximal ideal of $\Lambda_L(\mathfrak{X})$, with residue field k . Lemma 2.10.1 above implies that $\mathfrak{m}(\mathfrak{X}) = \mathfrak{m}_{\mathbb{C}_p}[[Z]]^{G_L,*}$.

Lemma 2.10.2. *The ring $\Lambda_L(\mathfrak{X})$ is a local ring.*

Proof. We have to show that $\mathfrak{m}(\mathfrak{X})$ is the unique maximal ideal, i.e., that f is a unit in $\Lambda_L(\mathfrak{X})$ if and only if $f(\chi_{\text{triv}}) \in o_L^\times$. The direct implication is obvious. We therefore assume that $f(\chi_{\text{triv}}) \in o_L^\times$. The image $F(Z) \in o_{\mathbb{C}_p}[[Z]]$ of f under the LT-isomorphism then satisfies $F(0) \in o_L^\times$ and hence is a unit in $o_{\mathbb{C}_p}[[Z]]$. We deduce that f is a unit in $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$. Since the twisted G_L -action must fix

with f also its inverse we obtain that f is a unit in $\mathcal{O}_L(\mathfrak{X})$ and hence in $\mathcal{O}_L^b(\mathfrak{X})$ by [5] Cor. 1.24. The multiplicativity of the norm $\|\cdot\|_{\mathfrak{X}}$ finally implies that $1 = \|f\|_{\mathfrak{X}} = \|f^{-1}\|_{\mathfrak{X}}$. \square

The \mathcal{O}_L -algebra $\Lambda_L(\mathfrak{X})$ carries two natural topologies. One is the p -adic topology which is induced by the norm $\|\cdot\|_{\mathfrak{X}}$. The other is the topology induced by the Fréchet topology of $\mathcal{O}_L(\mathfrak{X})$. We will call the latter the weak topology on $\Lambda_L(\mathfrak{X})$.

Remark 2.10.3. The weak topology on $\Lambda_L(\mathfrak{X})$ is coarser than the p -adic topology.

Proof. Let $\mathfrak{X} = \bigcup_{n \geq 1} \mathfrak{X}_n$ be a Stein covering by affinoid subdomains \mathfrak{X}_n (cf. [5] §1.3). The Fréchet topology of $\mathcal{O}_L(\mathfrak{X})$ is the projective limit of the Banach topologies on the affinoid algebras $\mathcal{O}_L(\mathfrak{X}_n)$. Since \mathfrak{X} is reduced these Banach topologies are defined by the respective supremum norm (cf. [7, Thm. 6.2.4/1]). Therefore the Banach topology on $\mathcal{O}_L(\mathfrak{X}_n)$ induces on its unit ball with respect to the supremum norm the p -adic topology. It follows that the natural maps $\Lambda_L(\mathfrak{X}) \rightarrow \mathcal{O}_L(\mathfrak{X}_n)$ are continuous for the p -adic topology on the source and the Banach topology on the target. Therefore the inclusion $\Lambda_L(\mathfrak{X}) \subseteq \mathcal{O}_L(\mathfrak{X})$ is continuous for the p -adic topology on the source and the Fréchet topology on the target. \square

Lemma 2.10.4. The \mathcal{O}_L -module $\Lambda_L(\mathfrak{X})$ is p -adically separated and complete.

Proof. We show that, for any reduced rigid analytic variety \mathfrak{Y} over L , the ring $\mathcal{O}_L^{\leq 1}(\mathfrak{Y})$ of holomorphic functions bounded by 1 is p -adically separated and complete. Let $\mathfrak{Y} = \bigcup_{i \in I} \mathfrak{Y}_i$ be an admissible covering by affinoid subdomains. Since \mathfrak{Y} is assumed to be reduced, the supremum seminorm on each $\mathcal{O}_L(\mathfrak{Y}_i)$ is a norm and defines its affinoid Banach topology (cf. [5, §1.3]). Hence $\|\cdot\|_{\mathfrak{Y}}$ is a norm on $\mathcal{O}_L^b(\mathfrak{Y})$ and defines the p -adic topology on $\mathcal{O}_L^{\leq 1}(\mathfrak{Y})$. In particular, the p -adic topology on $\mathcal{O}_L^{\leq 1}(\mathfrak{Y})$ is separated. Now let $(f_n)_n$ be a Cauchy sequence for $\|\cdot\|_{\mathfrak{Y}}$ in $\mathcal{O}_L^{\leq 1}(\mathfrak{Y})$. It restricts to a Cauchy sequence in $\mathcal{O}_L^{\leq 1}(\mathfrak{Y}_i)$ for each $i \in I$ which converges to a function $g_i \in \mathcal{O}_L^{\leq 1}(\mathfrak{Y}_i)$. Obviously the g_i glue to a function $g \in \mathcal{O}_L^{\leq 1}(\mathfrak{Y})$. We have to show that the sequence $(f_n)_n$ converges to g with respect to $\|\cdot\|_{\mathfrak{Y}}$. Let $\epsilon > 0$ be arbitrary. First we find an integer $N > 0$ such that $\|f_m - f_n\|_{\mathfrak{Y}} < \epsilon$ for all $m, n > N$. Secondly, for any $i \in I$, we have $\|g - f_m\|_{\mathfrak{Y}_i} < \epsilon$ for all sufficiently large (depending on i) m . It follows that $\|g - f_n\|_{\mathfrak{Y}_i} \leq \max(\|g - f_m\|_{\mathfrak{Y}_i}, \|f_m - f_n\|_{\mathfrak{Y}_i}) \leq \max(\|g - f_m\|_{\mathfrak{Y}_i}, \|f_m - f_n\|_{\mathfrak{Y}}) < \epsilon$ for any $n > N$ and any $i \in I$. Hence $\|g - f_n\|_{\mathfrak{Y}} \leq \epsilon$ for any $n > N$. \square

Proposition 2.10.5. The \mathcal{O}_L -module $\Lambda_L(\mathfrak{X})$ is compact in the weak topology.

Proof. According to [13, Prop. 6.4.5] the space \mathfrak{X} is strictly quasi-Stein. This means that a Stein covering $\mathfrak{X} = \bigcup_{n \geq 1} \mathfrak{X}_n$ can be chosen such that the inclusion maps $\mathfrak{X}_n \subseteq \mathfrak{X}_{n+1}$ are relatively compact. By loc. cit. Prop. 2.1.16 this implies that the restriction maps $\mathcal{O}_L(\mathfrak{X}_{n+1}) \rightarrow \mathcal{O}_L(\mathfrak{X}_n)$, which we simply view as inclusions, are compact maps between Banach spaces. Working over a locally

compact field we deduce (cf. [27, Remark 16.3] and [24, Cor. 6.1.14]) that the closure C_n of $\mathcal{O}_L^{\leq 1}(\mathfrak{X}_{n+1})$ in $\mathcal{O}_L(\mathfrak{X}_n)$ is compact. We, of course, have $\Lambda_L(\mathfrak{X}) \subseteq \mathcal{O}_L^{\leq 1}(\mathfrak{X}_{n+1}) \subseteq C_n$. Therefore, if $\mathcal{L}_n \subseteq \mathcal{O}_L(\mathfrak{X}_n)$ is any open lattice, then the \mathcal{O}_L -modules $\Lambda_L(\mathfrak{X})/\Lambda_L(\mathfrak{X}) \cap \mathcal{L}_n \subseteq C_n/C_n \cap \mathcal{L}_n$ are finite. It is straightforward to see that then $\Lambda_L(\mathfrak{X})/\Lambda_L(\mathfrak{X}) \cap \mathcal{L}$ must be finite for any open lattice $\mathcal{L} \subseteq \mathcal{O}_L(\mathfrak{X})$. On the other hand $\Lambda_L(\mathfrak{X})$ is weakly closed in $\mathcal{O}_L(\mathfrak{X})$ and hence is weakly complete. It follows [27, Cor. 7.6] that $\Lambda_L(\mathfrak{X})$ with its weak topology is the projective limit of the finite groups $\Lambda_L(\mathfrak{X})/\Lambda_L(\mathfrak{X}) \cap \mathcal{L}$ and hence is compact. \square

Lemma 2.10.6.

- (1) Any open neighbourhood of zero for the weak topology on $\Lambda_L(\mathfrak{X})$ contains a power of the maximal ideal $\mathfrak{m}(\mathfrak{X})$.
- (2) If the ideal $\mathfrak{m}(\mathfrak{X})$ is finitely generated then the weak topology on $\Lambda_L(\mathfrak{X})$ coincides with the $\mathfrak{m}(\mathfrak{X})$ -adic topology.

Proof. We have $\mathfrak{m}(\mathfrak{X}) = \pi_L \Lambda_L(\mathfrak{X}) + \mathfrak{n}$, where \mathfrak{n} denotes the ideal of all functions in $\Lambda_L(\mathfrak{X})$ which vanish in χ_{triv} . We consider the divisor Δ on \mathfrak{X} which maps χ_{triv} to 1 and all other points to zero. For any integer $m \geq 1$ we have the ideal $I_{m\Delta} \subseteq \mathcal{O}_L(\mathfrak{X})$ corresponding to the divisor $m\Delta$. As a consequence of [5, Prop. 1.4] these ideals are closed in $\mathcal{O}_L(\mathfrak{X})$ and satisfy $\bigcap_m I_m = \{0\}$. Hence the ideals $I_m \cap \Lambda_L(\mathfrak{X})$ are closed in $\Lambda_L(\mathfrak{X})$ with zero intersection. Let now $U \subseteq \Lambda_L(\mathfrak{X})$ be any fixed open neighbourhood of zero for the weak topology. Suppose that $I_m \cap \Lambda_L(\mathfrak{X}) \not\subseteq U$ for any $m \geq 1$. We then may pick, for any $m \geq 1$, a function $f_m \in (I_m \cap \Lambda_L(\mathfrak{X})) \setminus U$. According to Proposition 2.10.5 the weak topology on $\Lambda_L(\mathfrak{X})$ is compact. Hence the sequence $(f_m)_m$ has a convergent subsequence with a limit $f \in \Lambda_L(\mathfrak{X})$. On the one hand we have $f_n \in I_m \cap \Lambda_L(\mathfrak{X})$ for any $n \geq m$. Since $I_m \cap \Lambda_L(\mathfrak{X})$ is closed it follows that $f \in I_m \cap \Lambda_L(\mathfrak{X})$ for any $m \geq 1$. Therefore $f = 0$. But on the other hand all the f_m and hence f lie in the closed complement of the open subset U . This is a contradiction. We conclude that $\mathfrak{n}^m \subseteq I_m \cap \Lambda_L(\mathfrak{X}) \subseteq U$ for any sufficiently large m . As a consequence of Remark 2.10.3 we also have $\pi_L^m \Lambda_L(\mathfrak{X}) \subseteq U$ for any sufficiently large m . Hence $\mathfrak{m}(\mathfrak{X})^{2m} \subseteq \pi_L^m \Lambda_L(\mathfrak{X}) + \mathfrak{n}^m \subseteq U$ for large m . This proves (1).

We have to show that the ideals $\mathfrak{m}(\mathfrak{X})^m$ are open for the weak topology. Under our assumption all ideals $\mathfrak{m}(\mathfrak{X})^m$, for $m \geq 1$, are finitely generated. Hence all $\mathfrak{m}(\mathfrak{X})^{m+1}/\mathfrak{m}(\mathfrak{X})^m$ are finite dimensional k -vector spaces. We see that each quotient $\Lambda_L(\mathfrak{X})/\mathfrak{m}(\mathfrak{X})^m$, for $m \geq 1$, is a finite \mathcal{O}_L -module. Hence it suffices to show that the ideal $\mathfrak{m}(\mathfrak{X})^m$ is closed for the weak topology. Let f_1, \dots, f_r be generators of $\mathfrak{m}(\mathfrak{X})^m$. Then $\mathfrak{m}(\mathfrak{X})^m$ is the image of the map $\Lambda_L(\mathfrak{X})^r \rightarrow \Lambda_L(\mathfrak{X})$ sending (h_1, \dots, h_r) to $\sum_i h_i f_i$, which is a continuous map between compact spaces by Proposition 2.10.5. This proves (2). \square

Remark 2.10.7. Any $f \in \mathfrak{m}(\mathfrak{X})$ satisfies $\|f\|_{\mathfrak{X}_n} < 1$ for all n .

Proof. If $\|f\|_{\mathfrak{X}_n} = 1$ then the maximum modulus principle for the affinoid \mathfrak{X}_n implies that there is a point $z \in \mathfrak{X}_n$ such that $|f(z)| = 1$. By considering f as an element of $\mathcal{O}_{\mathbb{C}_p}[[T]]$, we see that $f(0)$ is a unit so that f is not in $\mathfrak{m}(\mathfrak{X})$. \square

Next we consider the injective map

$$\Lambda(o_L) = o_L[[o_L]] \longrightarrow \Lambda_L(\mathfrak{X}) ,$$

which we treat as an inclusion. More explicitly, let a_1, \dots, a_d be a basis of o_L as a \mathbb{Z}_p -module. Then the image of the above map is the ring of formal power series $o_L[[\delta_{a_1} - \delta_0, \dots, \delta_{a_d} - \delta_0]]$ inside $\Lambda_L(\mathfrak{X})$. We immediately conclude from Lemma 2.10.1 that

$$\mathfrak{m}(\mathfrak{X}) \cap o_L[[o_L]] = \langle \pi_L, \delta_{a_1} - \delta_0, \dots, \delta_{a_d} - \delta_0 \rangle \subseteq o_L[[o_L]] .$$

Lemma 2.10.8. *We have $\mathcal{O}_L^{\leq 1}(\mathfrak{X}) \cap o_L[[o_L]] = \pi_L o_L[[o_L]]$.*

Proof. We have $\pi_L o_L[[o_L]] \subseteq P := \mathcal{O}_L^{\leq 1}(\mathfrak{X}) \cap o_L[[o_L]]$. It follows that $\overline{P} := P/\pi_L o_L[[o_L]]$ is a “canonical” prime ideal in the formal power series ring $k[[o_L]]$: in particular, it is invariant for the o_L^\times action on the mod- p Iwasawa algebra $k[[o_L]]$. Suppose for a contradiction that \overline{P} is the (unique) maximal ideal of $k[[o_L]]$. Then P would contain $\delta_{a_1} - \delta_0$ and we would have $\|\delta_{a_1} - \delta_0\|_{\mathfrak{X}} < 1$. For each $n \geq 0$, let $\zeta_{p^n} \in o_{\mathbb{C}_p}$ denote a primitive p^n -root of unity, and let $\chi_n : o_L \rightarrow \mathbb{C}_p^\times$ be the unique torsion (hence locally L -analytic) character of o_L that sends a_1 to ζ_{p^n} and a_i to 1 for all $i > 1$. Then $\chi_n \in \mathfrak{X}(\mathbb{C}_p)$, and $|(\delta_{a_1} - \delta_0)(\chi_n)| \leq \|\delta_{a_1} - \delta_0\|_{\mathfrak{X}} < 1$ for all $n \geq 0$. However, $|(\delta_{a_1} - \delta_0)(\chi_n)| = |\zeta_{p^n} - 1| = |p|^{\frac{1}{p^n-1}(p-1)}$ tends to 1 from below, which is a contradiction. Hence \overline{P} is not the maximal ideal of $k[[o_L]]$.

In this situation, [1, Corollary 8.1(b)] implies that \overline{P} must be the zero ideal, provided we can show that the open subgroup $1 + po_L \subset o_L^\times$ acts *rationally irreducibly* on o_L .

We have to show that every non-trivial $1 + po_L$ -stable subgroup of o_L is open in o_L . But such a subgroup contains $(1 + po_L)a - a = pa o_L$ for some $0 \neq a \in o_L$, and is therefore open in o_L . \square

Corollary 2.10.9. *The restriction of the norm $\|\cdot\|_{\mathfrak{X}}$ on $\Lambda_L(\mathfrak{X})$ to $o_L[[o_L]]$ coincides with the π -adic norm on $o_L[[o_L]]$: for any $x \in \pi^n o_L[[o_L]] \setminus \pi^{n+1} o_L[[o_L]]$ we have*

$$\|x\|_{\mathfrak{X}} = |\pi^n|.$$

Proof. Since $\|\pi^n y\|_{\mathfrak{X}} = |\pi^n| \cdot \|y\|_{\mathfrak{X}}$ for any $y \in o_L[[o_L]]$, we may assume that $n = 0$. But now since $x \notin \pi o_L[[o_L]]$, Lemma 2.10.8 tells us that $\|x\|_{\mathfrak{X}} = 1$. \square

Corollary 2.10.10. *The o_L -module $\Lambda_L(\mathfrak{X})/o_L[[o_L]]$ is torsionfree.*

Proof. Suppose that $f \in \Lambda_L(\mathfrak{X})$ is such that $\pi^n f \in o_L[[o_L]]$ for some $n \geq 0$. Choose n least possible and suppose for a contradiction that $n \geq 1$. Then $\pi^n f \in o_L[[o_L]] \setminus \pi o_L[[o_L]]$, else otherwise we would be able to deduce that $\pi^{n-1} f \in o_L[[o_L]]$. Hence $\|\pi^n f\| = 1$ by Corollary 2.10.9, which implies that $|\pi|^{-n} = \|f\| \leq 1$. Hence $n = 0$. \square

Corollary 2.10.11. *We have $\Lambda_L(\mathfrak{X}) \cap (L \otimes_{o_L} o_L[[o_L]]) = o_L[[o_L]]$.*

3. THE KATZ ISOMORPHISM

3.1. The ψ_q -operator. Recall that we denote by \oplus the formal group law of \mathcal{G} . Furthermore let \mathcal{G}_1 denote the group of π -torsion points of \mathcal{G} . Its cardinality is q . It coincides with the set of zeros of the Frobenius power series $[\pi](Z) = \varphi(Z)$.

We fix a π -adically complete and flat o_L -algebra S in what follows and define an injective S -algebra endomorphism $\varphi : S[[Z]] \rightarrow S[[Z]]$ by setting

$$\varphi(F)(Z) := F([\pi](Z)) \quad \text{for all } F(Z) \in S[[Z]].$$

Lemma 3.1.1.

- (1) For any $F \in S[[Z]]$ there is a unique $F_0 \in S[[Z]]$ and a unique polynomial $F_1 \in S[Z]$ of degree $< q$ such that $F = \varphi(Z)F_0 + F_1$.
 (2) $\{F \in S[[Z]] : F(\zeta) = 0 \text{ for any } \zeta \in \mathcal{G}_1\} = \varphi(Z)S[[Z]]$.

Proof. (1). This is a form of Weierstrass division. Since $\varphi(Z) \equiv Z^q \pmod{\pi o_L[[Z]]}$, the proof of [8, VII.3.8 Prop. 5] goes through by replacing the maximal ideal of S in the argument with the ideal πS .

(2). Since $\varphi(Z)$ vanishes on \mathcal{G}_1 , the inclusion \supseteq is clear. If $F \in S[[Z]]$ vanishes on \mathcal{G}_1 then using (1) we may assume that $F \in S[Z]$ with $\deg F < q$. But then $F = 0$, which gives the other inclusion. \square

Using the above Lemma the proof of [10, Lemma 3] remains valid for S and gives

$$\varphi(S[[Z]]) = \{F \in S[[Z]] : F(Z) = F(\zeta \oplus Z) \text{ for all } \zeta \in \mathcal{G}_1\}.$$

Since the map φ is injective, this description of the image of φ implies the existence of a unique S -linear endomorphism ψ_{Col} of $S[[Z]]$ such that

$$\varphi(\psi_{\text{Col}}(F)(Z)) = \sum_{\zeta \in \mathcal{G}_1} F(\zeta \oplus Z) \quad \text{for any } F \in S[[Z]] \quad .$$

Definition 3.1.2. Let $S[[Z]]_L := S[[Z]] \otimes_{o_L} L$. The ψ_q -operator is defined by

$$\psi_q := \frac{1}{q} \psi_{\text{Col}} : S[[Z]]_L \rightarrow S[[Z]]_L.$$

Note that ψ_{Col} (respectively, ψ_q) preserves $S'[[Z]]$ (respectively, $S'[[Z]]_L$) for any intermediate π -adically complete and flat o_L -subalgebra S' of S . These operators satisfy the following useful Projection Formula.

Lemma 3.1.3. For any $F, G \in S[[Z]]$ we have $\psi_q(F\varphi(G)) = \psi_q(F)G$.

Proof. We may instead establish the analogous formula for ψ_{Col} . Note that $[\pi](\zeta \oplus Z) = [\pi](\zeta) \oplus [\pi](Z) = [\pi](Z)$ for any $\zeta \in \mathcal{G}_1$, since $[\pi](\zeta) = \varphi(\zeta) = 0$. Therefore

$$\begin{aligned} \varphi(\psi_{\text{Col}}(F\varphi(G))) &= \sum_{\zeta \in \mathcal{G}_1} (F\varphi(G))(\zeta \oplus Z) = \sum_{\zeta} F(\zeta \oplus Z)G([\pi](\zeta \oplus Z)) \\ &= \sum_{\zeta} F(\zeta \oplus Z)G([\pi](Z)) = \sum_{\zeta} F(\zeta \oplus Z)\varphi(G) \\ &= \varphi(\psi_{\text{Col}}(F))\varphi(G) = \varphi(\psi_{\text{Col}}(F)G) \quad . \end{aligned}$$

The result follows because φ is injective. \square

Corollary 3.1.4. *We have the fundamental equation $\psi_q \circ \varphi = 1_{S[[Z]]_L}$.*

Proof. Note that $\varphi(\psi_{\text{Col}}(1)) = q1$, so $\varphi(\psi_q(1)) = 1$ and hence $\psi_q(1) = 1$. Now set $F = 1$ in Lemma 3.1.3. \square

Next, we remind the reader what the operators φ and ψ_q do to the special power series $\Delta_a = \exp(a\Omega \log_{\text{LT}}(Z)) \in o_\infty[[Z]]$ from §2.7.

Lemma 3.1.5. *Assume that S is an o_∞ -algebra and take $a \in o_L$.*

- (1) $\varphi(\Delta_a) = \Delta_{\pi a}$.
- (2) $\psi_q(\Delta_a) = \delta_{a \in \pi o_L} \Delta_{a/\pi}$.

Proof. (1) More generally, whenever $a, b \in o_L$ we have

$$\Delta_a([b](Z)) = \exp(a\Omega \log_{\text{LT}}([b](Z))) = \exp(ab\Omega \log_{\text{LT}}(Z)) = \Delta_{ab}(Z).$$

Hence $\varphi(\Delta_a) = \Delta_a([\pi](Z)) = \Delta_{\pi a}$ as claimed.

(2) Using the fact that \log_{LT} is a formal homomorphism from \mathcal{G} to the formal additive group we compute

$$\begin{aligned} \varphi(\psi_{\text{Col}}(\Delta_a)) &= \sum_{\zeta \in \mathcal{G}_1} \Delta_a(\zeta \oplus Z) = \sum_{\zeta} \exp(a\Omega \log_{\text{LT}}(\zeta \oplus Z)) \\ &= \sum_{\zeta} \exp(a\Omega(\log_{\text{LT}}(\zeta) + \log_{\text{LT}}(Z))) \\ &= \left(\sum_{\zeta \in \mathcal{G}_1} \Delta_a(\zeta) \right) \Delta_a. \end{aligned}$$

Under the Schneider-Teitelbaum isomorphism κ , the group \mathcal{G}_1 corresponds to the group of characters χ of the finite group $o_L/\pi_L o_L$, and, if ζ corresponds to χ , then $\Delta_a(\zeta) = \text{ev}_{\bar{a}}(\chi) = \chi(\bar{a})$, where $\bar{a} := a + \pi o_L$. Hence

$$\varphi(\psi_{\text{Col}}(\Delta_a)) = \left(\sum_{\chi} \chi(\bar{a}) \right) \Delta_a.$$

By column orthogonality of characters of the finite group o_L/π_L , we have $\sum_{\chi} \chi(\bar{a}) = q\delta_{\bar{a}, \bar{0}} = q\delta_{a \in \pi o_L}$. Hence using part (1): $q\varphi(\psi_q(\Delta_a)) = q\delta_{a \in \pi o_L} \Delta_a = q\delta_{a \in \pi o_L} \varphi(\Delta_{a/\pi})$. Since φ is injective, we deduce that $\psi_q(\Delta_a) = \delta_{a \in \pi o_L} \Delta_{a/\pi}$ as required. \square

Write $\mathfrak{m} := \langle \pi, Z \rangle$ and $A := S[[Z]]$.

Lemma 3.1.6. *The operators φ and ψ_{Col} on A are \mathfrak{m} -adically continuous.*

Proof. Since $\varphi(Z) \in \langle Z \rangle$, we see that $\varphi(\mathfrak{m}^n) \subseteq \langle \pi, \varphi(Z) \rangle^n \subseteq \mathfrak{m}^n$ for all $n \geq 0$. This implies the \mathfrak{m} -adic continuity of φ .

Suppose first that \mathcal{G}_1 is contained in S . Then the S -linear maps $A \rightarrow A$ sending $F(Z)$ to $F(Z \oplus \zeta)$ are continuous with respect to \mathfrak{m} -adic topology for each $\zeta \in \mathcal{G}_1$; hence $\varphi \circ \psi_{\text{Col}}$ is also \mathfrak{m} -adically continuous in this case. Let $L_1 = L(\mathcal{G}_1)$, a finite extension of L and let $S_1 := o_{L_1} \otimes_{o_L} S$. Since o_{L_1} is

a free \mathcal{O}_L -module of finite rank, S_1 is still a π -adically complete and flat \mathcal{O}_L -algebra, so letting $A_1 = S_1[[Z]]$, we see that $\varphi \circ \psi_{\text{Col}} : A_1 \rightarrow A_1$ is $\mathfrak{m}A_1$ -adically continuous. It follows that $\varphi \circ \psi_{\text{Col}} : A \rightarrow A$ is also \mathfrak{m} -adically continuous.

Let $n \geq 0$ be given. Since $\varphi(Z) \equiv Z^q \pmod{\pi A}$, we have $\mathfrak{m}^{qn} = \langle \pi, Z \rangle^{qn} \subseteq \langle \pi, Z^q \rangle^n = \langle \pi, \varphi(Z) \rangle^n = A\varphi(\mathfrak{m}^n)$. Therefore $\mathfrak{m}^{qn} \cap \varphi(A) \subseteq A\varphi(\mathfrak{m}^n) \cap \varphi(A) = \varphi(\mathfrak{m}^n)$ where this last equation follows from the fact that $\varphi(A)$ admits a direct complement in A as a $\varphi(A)$ -module. However since $\varphi \circ \psi_{\text{Col}}$ is continuous, $\varphi\psi_{\text{Col}}(\mathfrak{m}^m) \subseteq \mathfrak{m}^{qn}$ for some $m \geq 0$. Hence

$$\varphi\psi_{\text{Col}}(\mathfrak{m}^m) \subseteq \mathfrak{m}^{qn} \cap \varphi(A) \subseteq \varphi(\mathfrak{m}^n).$$

The \mathfrak{m} -adic continuity of ψ_{Col} now follows from the injectivity of φ . \square

Lemma 3.1.7. *We have $\varphi^n(a_n) \rightarrow 0$ in the \mathfrak{m} -adic topology on A , for any sequence of elements (a_n) contained in ZA .*

Proof. Since $\varphi(Z) \in \mathcal{G}$ we see that $\varphi(Z) \in Z\mathfrak{m}$. Assume inductively that $\varphi^n(Z) \in Z\mathfrak{m}^n$; then $\varphi^{n+1}(Z) \in \varphi(Z\mathfrak{m}^n) \subseteq \varphi(Z)\mathfrak{m}^n \subseteq Z\mathfrak{m}^{n+1}$, completing the induction. Write $a_n = Zb_n$ for some $b_n \in A$; then $\varphi^n(a_n) = \varphi^n(Z)\varphi(b_n) \in Z\mathfrak{m}^n \subseteq \mathfrak{m}^{n+1}$ for all $n \geq 0$, so $\varphi^n(a_n) \rightarrow 0$. \square

We specialize to the case $S = \mathcal{O}_\infty$ until the end of §3.1.

Proposition 3.1.8. *Let $f \in D^{L-\text{an}}(\mathcal{O}_L, L)$ be such that $\mathcal{F}(f) \in \mathcal{O}(\mathfrak{X})^\circ$. Suppose that $\psi_q^n(\mu_{\text{rig}}(f)\Delta_a) \in \mathcal{O}_\infty[[Z]]$ for all $a \in \mathcal{O}_L$ and $n \geq 0$. Then $f \in \mathcal{O}_L[[\mathcal{O}_L]]$.*

Proof. We will show that $|f(\mathbf{1}_{a+\pi^n\mathcal{O}_L})| \leq 1$ for all $a \in \mathcal{O}_L$ and $n \geq 0$.

By [28, Lemma 4.6(4)], we have

$$f(\mathbf{1}_{a+\pi^n\mathcal{O}_L}) = (f\delta_{-a})(\mathbf{1}_{\pi^n\mathcal{O}_L}).$$

The orthogonality of columns in the character table of the finite group $\mathcal{O}_L/\pi^n\mathcal{O}_L$ implies that

$$\mathbf{1}_{\pi^n\mathcal{O}_L} = \frac{1}{q^n} \sum_{[\pi^n](z)=0} \kappa_z.$$

Hence by *ibid.*, $(f\delta_{-a})(\mathbf{1}_{\pi^n\mathcal{O}_L}) = \frac{1}{q^n} \sum_{[\pi^n](z)=0} f(z)\Delta_{-a}(z)$. We now observe that

$$\frac{1}{q^n} \sum_{[\pi^n](z)=0} f(z)\Delta_{-a}(z) = \psi_q^n(\mu_{\text{rig}}(f)\Delta_{-a})(0).$$

Since $\psi_q^n(\mu_{\text{rig}}(f)\Delta_{-a}) \in \mathcal{O}_\infty[[Z]]$ by assumption, we have $|f(\mathbf{1}_{a+\pi^n\mathcal{O}_L})| \leq 1$ for all $a \in \mathcal{O}_L$ and $n \geq 0$, as claimed.

Therefore there exists $g \in \mathcal{O}_L[[\mathcal{O}_L]]$ such that $f = g$ on all $\mathbf{1}_{a+\pi^n\mathcal{O}_L}$. The function $f - g$ is zero on all locally constant functions, and hence on all torsion characters, so that it is divisible by $\log(1 + Z)$. Since $f - g$ is bounded and $\log(1 + Z)$ is unbounded, this implies that $f = g$. \square

We can now prove Theorem 1.4.1 from the introduction.

Theorem 3.1.9. *We have $\Lambda_L(\mathfrak{X}) = \mathcal{O}_L[[\mathcal{O}_L]]$ if and only if $\psi_q(\Lambda_L(\mathfrak{X})) \subset \Lambda_L(\mathfrak{X})$.*

Proof. The forward implication is clear in view of Lemma 3.1.5(1). For the reverse implication, the given condition means that the image $o_\infty[[Z]]^{G_L,*}$ of $\Lambda_L(\mathfrak{X}) = \mathcal{O}(\mathfrak{X})^\circ$ under the bijection $(\mu_{\text{rig}} \circ \mathcal{F}^{-1})^\circ$ from Corollary 2.9.3 is stable under the ψ_q -operator. Now the result follows from Proposition 3.1.8. \square

3.2. The covariant bialgebra of \mathcal{G} . Katz [20, §1] talks about the “algebra $\text{Diff}(\mathcal{G})$ of all \mathcal{G} -invariant o_L -linear differential operators from $\mathcal{O}(\mathcal{G})$ into itself”. Because we are not aware of any place in the literature which adequately deals with invariant differential operators on formal groups, we will instead use the *covariant bialgebra* of \mathcal{G} which will turn out to be isomorphic to Katz’s $\text{Diff}(\mathcal{G})$.

Definition 3.2.1.

- (1) Let $Z_1 \oplus Z_2 \in o_L[[Z_1, Z_2]]$ denote the formal group law defining the formal group \mathcal{G} .
- (2) Let $U(\mathcal{G})$ denote the set of all o_L -linear maps from $\mathcal{O}(\mathcal{G}) = o_L[[Z]]$ to o_L that vanish on some power of the augmentation ideal $Zo_L[[Z]]$. In other words,

$$U(\mathcal{G}) = \varinjlim \text{Hom}_{o_L}(\mathcal{O}(\mathcal{G})/Z^n\mathcal{O}(\mathcal{G}), o_L).$$

We will often use the abbreviation $U := U(\mathcal{G})$.

- (3) For each $f, g \in U(\mathcal{G})$, define the product $f \cdot g$ by the formula

$$(f \cdot g)(F(Z)) = (f \hat{\otimes} g)(F(Z_1 \oplus Z_2)) \quad \text{for all } F(Z) \in o_L[[Z]].$$
- (4) With this product, $U(\mathcal{G})$ is the *covariant bialgebra* of \mathcal{G} , defined at [17, 36.1.8].
- (5) For each $m \geq 0$, let $u_m \in U(\mathcal{G})$ be the unique o_L -linear map that satisfies

$$u_m(Z^n) = \delta_{mn} \quad \text{for all } n \geq 0.$$

- (6) Let $\langle -, - \rangle : U(\mathcal{G}) \times \mathcal{O}(\mathcal{G}) \rightarrow o_L$ be the evaluation pairing:

$$\langle f, F \rangle := f(F).$$

This covariant bialgebra is also known as the *hyperlgebra* or the *distribution algebra* of \mathcal{G} . Note that $U(\mathcal{G})$ is a commutative ring: this follows directly from Definition 3.2.1(3), as the formal group law $Z_1 \oplus Z_2$ on the Lubin-Tate formal group \mathcal{G} is commutative. We will now explain the link with Katz’s work, using his notation.

Lemma 3.2.2.

- (1) $\{u_n : n \geq 0\}$ is an o_L -module basis for $U(\mathcal{G})$.
- (2) Let $i \geq 0$ and write $(Z_1 \oplus Z_2)^i = \sum_{\substack{n, m \geq 0 \\ n+m \geq i}}^\infty \lambda(n, m; i) Z_1^n Z_2^m$ for some $\lambda(n, m; i) \in o_L$. Then for all $n, m \geq 0$ we have

$$u_n \cdot u_m = \sum_{k=0}^{n+m} \lambda(n, m; k) u_k.$$

(3) Let s be a variable. The map $L[s] \rightarrow U(\mathcal{G}) \otimes_{o_L} L$ which sends s to $u_1 \otimes 1$ is an isomorphism of positively filtered L -algebras.

Proof. (1) This is clear because $Z^n o_L[[Z]] = o_L Z^n \oplus Z^{n+1} o_L[[Z]]$ for any $n \geq 0$.

(2) We compute that for every $n, m, i \geq 0$ we have

$$\begin{aligned} (u_n \cdot u_m)(Z^i) &= (u_n \widehat{\otimes} u_m)((Z_1 \oplus Z_2)^i) \\ &= (u_n \widehat{\otimes} u_m) \left(\sum_{\substack{a, b \geq 0 \\ a+b \geq i}}^{\infty} \lambda(a, b; i) Z_1^a Z_2^b \right) = \lambda(n, m; i). \end{aligned}$$

Because $\sum_{k=0}^{n+m} \lambda(n, m; k) u_k$ also sends Z^i to $\lambda(n, m; i)$, it must be equal to $u_n \cdot u_m$.

(3) From (2) we see that the o_L -submodule $U(\mathcal{G})_n$ of $U(\mathcal{G})$ generated by $\{u_i : 0 \leq i \leq n\}$ defines an algebra filtration on $U(\mathcal{G})$:

$$U(\mathcal{G})_n \cdot U(\mathcal{G})_m \subseteq U(\mathcal{G})_{n+m} \quad \text{for all } n, m \geq 0.$$

The associated graded ring is the free o_L -module with basis $\{\text{gr } u_n : n \geq 0\}$. Since $Z_1 \oplus Z_2 \equiv Z_1 + Z_2 \pmod{(Z_1, Z_2)^2}$ by part (2), we see that for any $i \geq 0$ we have

$$\begin{aligned} \sum_{\substack{n, m \geq 0 \\ n+m \geq i}}^{\infty} \lambda(n, m; i) Z_1^n Z_2^m &= (Z_1 \oplus Z_2)^i \\ &\equiv (Z_1 + Z_2)^i = \sum_{n+m=i} \binom{i}{n} Z_1^n Z_2^m \pmod{(Z_1, Z_2)^{i+1}}. \end{aligned}$$

Equating the coefficient of $Z_1^n Z_2^{i-n}$ shows that $\lambda(n, i-n; i) = \binom{i}{n}$ whenever $0 \leq n \leq i$:

$$\lambda(n, m; n+m) = \binom{n+m}{n} \quad \text{for any } n, m \geq 0.$$

Hence from (2) we see that the multiplication in $\text{gr } U(\mathcal{G})$ is given by

$$(\text{gr } u_n) \cdot (\text{gr } u_m) = \binom{n+m}{n} \text{gr } u_{n+m}.$$

The same formulas hold in $\text{gr}(U(\mathcal{G}) \otimes_{o_L} L)$. Induction on n shows that we have $(\text{gr } u_1)^n = n! \text{gr } u_n$ for all $n \geq 0$. Since L has characteristic zero, we see that $\text{gr}(U(\mathcal{G}) \otimes_{o_L} L)$ is generated by $\text{gr } u_1$ as an L -algebra. The result follows. \square

We will henceforth identify $U(\mathcal{G}) \otimes_{o_L} L$ with the polynomial ring $L[s]$. Recall the polynomials $P_n(Y) \in L[Y]$ from [28, Definition 4.1], which are defined by the following formal expansion:

$$\exp(Y \log_{\text{LT}}(Z)) = \sum_{m=0}^{\infty} P_m(Y) Z^m.$$

Lemma 3.2.3. *For every $n \geq 0$, we have $u_n = P_n(u_1)$ inside $U(\mathcal{G}) \otimes_{o_L} L$.*

Proof. The structure constants of Katz's algebra $\text{Diff}(\mathcal{G})$ are the same as the ones in $U(\mathcal{G})$ by [20, (1.2)] and Lemma 3.2.2(2). So the o_L -linear map that sends $D(n) \in \text{Diff}(\mathcal{G})$ to $u_n \in U(\mathcal{G})$ is an o_L -algebra isomorphism. Comparing [20, Corollary 1.8] with [28, Definition 4.1] shows that $D(n) = P_n(D(1))$ in $\text{Diff}(\mathcal{G}) \otimes_{o_L} L$ for all $n \geq 0$. The result follows by applying the algebra isomorphism $\text{Diff}(\mathcal{G}) \rightarrow U(\mathcal{G})$ established above. \square

Of course in the context of affine group schemes, this isomorphism between the algebra of left-invariant differential operators on the group scheme and the distribution algebra of the group scheme is the well known 'Invariance Theorem', [12, Chapter II, §4, Theorem 6.6].

Next, we consider the action of the monoid o_L on the formal group \mathcal{G} . The covariant bialgebra construction is functorial in \mathcal{G} : if $\varphi : \mathcal{G} \rightarrow \mathcal{H}$ is a morphism of formal groups, then $U(\varphi) : U(\mathcal{G}) \rightarrow U(\mathcal{H})$ is the morphism of o_L -bialgebras which is the transpose to the o_L -algebra homomorphism $\varphi^* : \mathcal{O}(\mathcal{H}) \rightarrow \mathcal{O}(\mathcal{G})$ induced by φ . Using the evaluation pairing, we have the following formula which defines this action:

$$(4) \quad \langle U(\varphi)(f), F \rangle = \langle f, \varphi^*(F) \rangle \quad \text{for all } f \in U(\mathcal{G}), F \in \mathcal{O}(\mathcal{G}).$$

Definition 3.2.4. Take $a \in o_L$.

- (1) Let $[a] : \mathcal{G} \rightarrow \mathcal{G}$ be the action of a on \mathcal{G} .
- (2) Write $a \cdot f := U([a])(f)$ for all $f \in U(\mathcal{G})$.

The o_L -algebra endomorphism $U([a])$ of $U(\mathcal{G})$ extends to an L -algebra endomorphism $U([a]) \otimes 1$ of $U(\mathcal{G}) \otimes_{o_L} L = L[s]$. What does this action do to the generator s of $L[s]$?

Lemma 3.2.5. We have $a \cdot s = as$ for all $a \in o_L$.

Proof. We know that $[a](Z) \equiv aZ \pmod{Z^2 o_L[[Z]]}$. Hence

$$\langle U([a])(u_1), Z^n \rangle = \langle u_1, [a](Z)^n \rangle = a\delta_{n,1} = \langle au_1, Z^n \rangle \quad \text{for all } n \geq 0$$

using Definition 3.2.1(5). Hence $a \cdot u_1 = au_1$ and so $a \cdot s = as$. \square

Corollary 3.2.6. For each $j \geq i \geq 0$ and $a \in o_L$ there exists $\sigma_{ij}(a) \in o_L$ such that

$$a \cdot u_j = P_j(as) = \sum_{i=0}^j \sigma_{ij}(a) P_i(s) = \sum_{i=0}^j \sigma_{ij}(a) u_i.$$

Proof. It follows from Lemma 3.2.5 that the L -algebra endomorphisms of $L[s]$ given by $s \mapsto as$ preserve the o_L -subalgebra $U(\mathcal{G}) \subset L[s]$. Hence $a \cdot u_j = P_j(as)$ lies in $U(\mathcal{G})$ for all $a \in o_L$ and all $j \geq 0$. But $U(\mathcal{G})$ has $\{u_i : i \geq 0\}$ as an o_L -module basis by Lemma 3.2.2(a), so $P_j(as)$ must be an o_L -linear combination of these u_i 's. On the other hand, $P_j(s)$ is a polynomial of degree j in s , therefore so is $P_j(as)$; because $\deg P_i = i$ for each i it follows that $P_j(as)$ is an L -linear combination of $P_0(s), \dots, P_j(s)$ only. \square

We now introduce a coefficient ring S , which we assume to be a π -adically complete o_L -algebra. For every S -module M , let $M^* := \text{Hom}_S(M, S)$ be the S -module of S -linear functionals on M . We will need to work with a larger class of S -linear functionals on $S[[Z]]$ than those arising from $U(\mathcal{G})$, namely the *continuous* ones.

Definition 3.2.7. We say that $\lambda \in S[[Z]]^*$ is *continuous* if it is continuous with respect to the $\langle \pi, Z \rangle$ -adic topology on $S[[Z]]$, and the π -adic topology on S . Let $S[[Z]]_{\text{cts}}^*$ denote the set of these continuous S -linear functionals on $S[[Z]]$.

Explicitly $\lambda \in S[[Z]]^*$ is continuous if and only if for all $n \geq 0$ there exists $m \geq 0$ such that $\lambda(\langle \pi, Z \rangle^m) \subseteq \pi^n S$.

Consider now the base change $U(\mathcal{G}_S) := U(\mathcal{G}) \otimes_{o_L} S$, and its π -adic completion

$$\widehat{U(\mathcal{G}_S)} = \varprojlim U(\mathcal{G}) \otimes_{o_L} (S/\pi^n S).$$

Since $\{u_m : m \geq 0\}$ is an o_L -module basis for $U(\mathcal{G})$ by Lemma 3.2.2(1), we see that $\widehat{U(\mathcal{G}_S)}$ has the following description:

$$(5) \quad \widehat{U(\mathcal{G}_S)} = \left\{ \sum_{m=0}^{\infty} a_m u_m : a_m \in S, \lim_{m \rightarrow \infty} a_m = 0 \right\}.$$

Here we equip S with the π -adic topology.

Lemma 3.2.8.

(1) The pairing $\langle -, - \rangle : U(\mathcal{G}) \times o_L[[Z]] \rightarrow o_L$ extends to an S -bilinear pairing

$$\langle -, - \rangle : \widehat{U(\mathcal{G}_S)} \times S[[Z]] \rightarrow S.$$

(2) For each $u \in \widehat{U(\mathcal{G}_S)}$, the S -linear map $\langle u, - \rangle : S[[Z]] \rightarrow S$ is continuous.

(3) The map $\widehat{U(\mathcal{G}_S)} \rightarrow S[[Z]]_{\text{cts}}^*$, $u \mapsto \langle u, - \rangle$, is an S -linear bijection.

(4) The map $S[[Z]] \rightarrow \widehat{U(\mathcal{G}_S)}^*$, $F \mapsto \langle -, F \rangle$, is an S -linear bijection.

Proof. (1) Let $u = \sum_{m=0}^{\infty} a_m u_m \in \widehat{U(\mathcal{G}_S)}$, $F = \sum_{n=0}^{\infty} F_n Z^n \in S[[Z]]$ and define

$$\langle u, F \rangle = \sum_{m=0}^{\infty} a_m F_m. \quad \text{This series converges in } S \text{ because } a_m \rightarrow 0 \text{ as } m \rightarrow \infty$$

and because S is assumed to be π -adically complete.

(2) Let $n \geq 0$ and write $u = \sum_{m=0}^{\infty} a_m u_m$ with $a_m \rightarrow 0$. Then for some $r \geq 0$, $a_m \in \pi^n S$ for all $m \geq r$. Hence $\langle u, - \rangle$ sends the ideal $\langle \pi^n, Z^r \rangle$ of $S[[Z]]$ into $\pi^n S$. Since $\langle \pi, Z \rangle^{n+r} \subseteq \langle \pi^n, Z^r \rangle$, we conclude that $\langle u, - \rangle$ is $\langle \pi, Z \rangle$ -adically continuous.

(3) The injectivity of $u \mapsto \langle u, - \rangle$ follows by evaluating on each Z^n . Now let $\lambda \in S[[Z]]_{\text{cts}}^*$ and define $a_m := \lambda(Z^m) \in S$ for each $m \geq 0$. Since λ is $\langle \pi, Z \rangle$ -adically continuous, for each $n \geq 0$ we can find some $r \geq 0$ such that $\lambda(\langle \pi, Z \rangle^r) \subseteq \pi^n S$. Then $a_m \in \pi^n S$ for all $m \geq r$ which implies that $a_m \rightarrow 0$

as $m \rightarrow \infty$. Hence $u := \sum_{m=0}^{\infty} a_m u_m$ is an element of $\widehat{U(\mathcal{G}_S)}$ and $\langle u, - \rangle - \lambda$ vanishes on $S[Z]$ by construction. Since this difference is continuous and since $S[Z]$ is dense in $S[[Z]]$ with respect to the $\langle \pi, Z \rangle$ -adic topology, we conclude that $\lambda = \langle u, - \rangle$.

(4) Again, the injectivity of $F \mapsto \langle -, F \rangle$ follows from $\langle u_m, F \rangle = F_m$. Given an S -linear map $\lambda : U(\mathcal{G}_S) \rightarrow S$, let $F := \sum_{n=0}^{\infty} \lambda(u_n) Z^n$. Then $\langle u_m, F \rangle = \lambda(u_m)$ for all $m \geq 0$. Since the u_m span $U(\mathcal{G}_S)$ as an S -module, $\lambda = \langle -, F \rangle$. \square

As an immediate consequence of Lemma 3.2.8, we have the following

Corollary 3.2.9.

(1) For every continuous S -linear $\alpha : S[[Z]] \rightarrow S[[Z]]$ there exists a unique S -linear map $\alpha^* : \widehat{U(\mathcal{G}_S)} \rightarrow \widehat{U(\mathcal{G}_S)}$ such that

$$\langle \alpha^* u, F \rangle = \langle u, \alpha F \rangle \quad \text{for all } u \in \widehat{U(\mathcal{G}_S)}, F \in S[[Z]].$$

(2) For every S -linear $\beta : \widehat{U(\mathcal{G}_S)} \rightarrow \widehat{U(\mathcal{G}_S)}$ there exists a unique S -linear map $\beta^* : S[[Z]] \rightarrow S[[Z]]$ such that

$$\langle u, \beta F \rangle = \langle \beta^* u, F \rangle \quad \text{for all } u \in \widehat{U(\mathcal{G}_S)}, F \in S[[Z]].$$

We also extend this S -linear pairing to an $S_L := S \otimes_{o_L} L$ -linear pairing

$$\langle -, - \rangle : \widehat{U(\mathcal{G}_S)}_L \times S[[Z]]_L \rightarrow S_L$$

which we will use without further mention. Observe that there is a natural o_L -linear map $\widehat{U} \rightarrow \widehat{U(\mathcal{G}_S)}$ for any o_L -algebra S .

Lemma 3.2.10. *The restriction map $\widehat{U(\mathcal{G}_S)}^* \rightarrow \text{Hom}_{o_L}(\widehat{U}, S)$ is an S -linear isomorphism.*

Proof. Let $\lambda : \widehat{U(\mathcal{G}_S)} \rightarrow S$ be an S -linear map whose restriction to \widehat{U} is zero. Then in particular $\lambda(u_m) = 0$ for all $m \geq 0$, so λ vanishes on all finite sums of the form $\sum_{m=0}^n a_m u_m \in \widehat{U(\mathcal{G}_S)}$ with $a_m \in S$. These sums are π -adically dense in $\widehat{U(\mathcal{G}_S)}$ in view of (5), so for any $x \in \widehat{U(\mathcal{G}_S)}$, $\lambda(x) \in \bigcap_{n=0}^{\infty} \pi^n S$. Since we're assuming that S is π -adically complete, this intersection is zero, so $\lambda = 0$ and the restriction map in question is injective.

Suppose now $\lambda : \widehat{U} \rightarrow S$ is an o_L -linear map. Using the description of $\widehat{U(\mathcal{G}_S)}$ given in (5), we extend it to an S -linear map $\tilde{\lambda} : \widehat{U(\mathcal{G}_S)} \rightarrow S$ by setting for every zero-sequence (a_m) in S

$$\tilde{\lambda} \left(\sum_{m=0}^{\infty} a_m u_m \right) := \sum_{m=0}^{\infty} a_m \lambda(u_m).$$

Since $\lim_{m \rightarrow \infty} a_m = 0$ in S , the series on the right hand side converges in S because S is assumed to be π -adically complete. So, $\tilde{\lambda}$ is a well-defined S -linear map extending λ . \square

3.3. Gal-continuous functions. Let $\mathcal{C}^0(o_L, \mathbb{C}_p)$ be the \mathbb{C}_p -Banach space of all continuous \mathbb{C}_p -valued functions on o_L , equipped with the supremum norm. The unit ball of this \mathbb{C}_p -Banach space is the $o_{\mathbb{C}_p}$ -submodule $\mathcal{C}^0(o_L, o_{\mathbb{C}_p})$ of continuous $o_{\mathbb{C}_p}$ -valued functions.

Definition 3.3.1. A function $f \in \mathcal{C}^0(o_L, \mathbb{C}_p)$ is said to be *Gal-continuous* if

$$\sigma(f(a)) = f(a\tau(\sigma)) \quad \text{for all } a \in o_L, \sigma \in G_L.$$

We write $C := \mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$ for the set of all Gal-continuous \mathbb{C}_p -valued functions.

Evidently $\mathcal{C} := \mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}) = C \cap \mathcal{C}^0(o_L, o_{\mathbb{C}_p})$ forms an o_L -lattice in C .

Lemma 3.3.2. *Let $f \in C$. Then $\text{im } f \subseteq L_\infty$, and $\text{im } f \subseteq o_\infty$ if $f \in \mathcal{C}$.*

Proof. By Definition 3.3.1, we have $\text{im } f \subseteq \mathbb{C}_p^{\ker \tau}$ for all $f \in C$, and $\text{im } f \subseteq o_{\mathbb{C}_p}^{\ker \tau}$ for all $f \in \mathcal{C}$. But $\mathbb{C}_p^{\ker \tau} = L_\infty$ and $o_{\mathbb{C}_p}^{\ker \tau} = o_\infty$ by Lemma 2.7.2. \square

Lemma 3.3.3. *For each $u \in \widehat{U}$, the function $a \mapsto \mathcal{K}(u)(a) := \langle u, \Delta_a \rangle$ on o_L is Gal-continuous.*

Proof. By definition, $\mathcal{K}(u)$ is the composition of $\mu|_{o_L} : o_L \rightarrow o_\infty[[Z]]^\times$ with the restriction of the linear functional $\langle u, - \rangle : o_\infty[[Z]] \rightarrow o_\infty$ to $o_\infty[[Z]]^\times$. This linear functional is continuous by Lemma 3.2.8(3), so to establish the continuity of $\mathcal{K}(u)$ it remains to show that $\mu|_{o_L}$ is continuous. Since $\mu|_{o_L}$ is a group homomorphism, it is enough to show that it is continuous at the identity element 0 of o_L . Let $n > 0$ and consider the basic open neighbourhood $1 + \langle \pi, Z \rangle^n$ of $1 \in o_\infty[[Z]]^\times$. Since $\varphi^n(Z) \rightarrow 0$ as $n \rightarrow \infty$ in $o_\infty[[Z]]$ by Lemma 3.1.7, we can find $m \geq 0$ such that $\varphi^m(Z) \in \langle \pi, Z \rangle^n$. Hence for any $a \in o_L$, using Lemma 3.1.5 we calculate

$$\Delta_{\pi^m a} - \Delta_0 = \varphi^m(\Delta_a - 1) \in \varphi^m(Z o_\infty[[Z]]) \subseteq \varphi^m(Z) o_\infty[[Z]] \subseteq \langle \pi, Z \rangle^n.$$

Hence $\mu|_{o_L}$ is continuous as required.

Now let $\sigma \in G_L$; since $\Delta_a \in o_\infty[[Z]]$ is invariant for the $*$ -action of G_L on $o_\infty[[Z]]$, we know that $\sigma(\Delta_a) = \Delta_a([\tau(\sigma)](Z)) = \Delta_{a\tau(\sigma)}$ for any $a \in o_L$. Since $u \in \widehat{U}$, we have for any $a \in o_L$

$$\sigma(\mathcal{K}(u)(a)) = \sigma(\langle u, \Delta_a \rangle) = \langle u, \sigma(\Delta_a) \rangle = \langle u, \Delta_{a\tau(\sigma)} \rangle = \mathcal{K}(u)(a\tau(\sigma)).$$

Hence $\mathcal{K}(u)$ is indeed Gal-continuous. \square

Definition 3.3.4.

(1) Define the *Katz map* $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ as follows:

$$\mathcal{K}(u)(a) = \langle u, \Delta_a \rangle \quad \text{for any } u \in \widehat{U}, a \in o_L.$$

- (2) Define $\mathcal{K}_1 : \widehat{U} \rightarrow o_\infty$ by $\mathcal{K}_1 = \text{ev}_1 \circ \mathcal{K}$.
 (3) Define $\psi_C : C \rightarrow C$ by the rule

$$\psi_C(f)(a) = \delta_{a \in \pi o_L} f(a/\pi) \quad \text{for all } a \in o_L.$$

The operator $\psi_C : \mathcal{C} \rightarrow \mathcal{C}$ is by definition the restriction of ψ_C to \mathcal{C} .

- (4) Define $\varphi_C : C \rightarrow C$ by the rule

$$\varphi_C(f)(a) = f(\pi a) \quad \text{for all } a \in o_L.$$

The operator $\varphi_C : \mathcal{C} \rightarrow \mathcal{C}$ is by definition the restriction of φ_C to \mathcal{C} .

Using his notation, Katz has already observed [19, p. 99] the following fact.

Lemma 3.3.5. *The map $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ is an o_L -algebra homomorphism.*

Proof. By the definition of the Lubin-Tate logarithm, we have

$$\log_{LT}(Z_1 \oplus Z_2) = \log_{LT}(Z_1) + \log_{LT}(Z_2)$$

from which it follows that for all $a \in o_L$ we have

$$\Delta_a(Z_1 \oplus Z_2) = \Delta_a(Z_1)\Delta_a(Z_2).$$

Using Definition 3.2.1(3), we can then compute that for any $u, v \in U$ and any $a \in o_L$ we have

$$\begin{aligned} \mathcal{K}(u \cdot v)(a) &= \langle u \cdot v, \Delta_a(Z) \rangle = (u \widehat{\otimes} v)(\Delta_a(Z_1 \oplus Z_2)) \\ &= (u \widehat{\otimes} v)(\Delta_a(Z_1)\Delta_a(Z_2)) = \mathcal{K}(u)(a)\mathcal{K}(v)(a). \end{aligned}$$

Hence $\mathcal{K}(u \cdot v) = \mathcal{K}(u)\mathcal{K}(v)$ for all $u, v \in U$ and the result follows easily. \square

Now we recall the coefficient ring S that was introduced before Definition 3.2.7. Applying the S -linear duality functor

$$(-)^* := \text{Hom}_{o_L}(-, S)$$

to the Katz map $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ gives us the *dual Katz map*

$$\mathcal{K}^* : \mathcal{C}^* \rightarrow \widehat{U}^*$$

defined on the space of S -valued Galois measures $\mathcal{C}^* = \text{Hom}_{o_L}(\mathcal{C}, S)$. We identify $\widehat{U}^* = \text{Hom}_{o_L}(\widehat{U}, S)$ with $S[[Z]]$ using Lemma 3.2.10 and Lemma 3.2.8(4); then $\mathcal{K}^* : \mathcal{C}^* \rightarrow S[[Z]]$ is given explicitly by

$$(6) \quad \langle u_m, \mathcal{K}^*(\lambda) \rangle = \lambda(P_m(-\Omega)) \quad \text{for all } \lambda \in \mathcal{C}^*, m \geq 0.$$

After Lemma 3.1.6 and Corollary 3.2.9 applied with $S = o_L$, we have at our disposal the dual o_L -linear endomorphisms ψ_{Col}^* and φ^* of \widehat{U} .

Lemma 3.3.6. *We have $\mathcal{K}\varphi^* = \varphi_C\mathcal{K}$ and $\mathcal{K}\psi_{\text{Col}}^* = q\psi_C\mathcal{K}$.*

Proof. Let $u \in \widehat{U}_L$ and $a \in o_L$. Then using Lemma 3.1.5, we have

$$\begin{aligned} \mathcal{K}(\psi_{\text{Col}}^*(u))(a) &= \langle \psi_{\text{Col}}^*(u), \Delta_a \rangle = \langle u, \psi_{\text{Col}}(\Delta_a) \rangle = \langle u, q\psi_q(\Delta_a) \rangle \\ &= q\langle u, \delta_{a \in \pi o_L} \Delta_a/\pi \rangle = q\delta_{a \in \pi o_L} \mathcal{K}(u)(a/\pi) = q\psi_C(\mathcal{K}(u))(a) \end{aligned}$$

which gives the second equation. The first equation is proved in a similar manner. \square

Corollary 3.3.7. *We have $\mathcal{K}^* \varphi_{\mathcal{C}}^* = \varphi \mathcal{K}^*$ and $\mathcal{K}^* \psi_{\mathcal{C}}^* = \psi_q \mathcal{K}^*$.*

Proof. We apply the S -linear duality functor $(-)^* = \text{Hom}_{o_L}(-, S)$ to the equations from Lemma 3.3.6. Using Lemma 3.2.8, we see that

$$\mathcal{K}^* \varphi_{\mathcal{C}}^* = (\varphi_{\mathcal{C}} \mathcal{K})^* = (\mathcal{K} \varphi^*)^* = \varphi^{**} \mathcal{K}^* = \varphi \mathcal{K}^*,$$

and similarly,

$$q \mathcal{K}^* \psi_{\mathcal{C}}^* = (q \psi_{\mathcal{C}} \mathcal{K})^* = (\mathcal{K} \psi_{\text{Col}}^*)^* = \psi_{\text{Col}}^* \mathcal{K}^* = q \psi_q \mathcal{K}^*.$$

Now divide both sides by q . \square

Lemma 3.3.8. *We have $\psi_q \circ \mathcal{K}_1^* = 0$.*

Proof. Corollary 3.3.7 gives $\psi_q \mathcal{K}_1^* = \psi_q \mathcal{K}^* \text{ev}_1^* = \mathcal{K}^* \psi_{\mathcal{C}}^* \text{ev}_1^* = (\text{ev}_1 \psi_{\mathcal{C}} \mathcal{K})^*$. But $\text{ev}_1 \psi_{\mathcal{C}}(f) = \psi_{\mathcal{C}}(f)(1) = 0$ for any $f \in \mathcal{C}$ by Definition 3.3.4(3), because $\delta_{1 \in \pi o_L} = 0$. \square

Proposition 3.3.9. *If \mathcal{K} is injective and τ is surjective, then $q \ker \mathcal{K}_1 \subseteq \psi_{\text{Col}}^*(\widehat{U})$.*

Proof. In this proof we may assume $S = o_L$. Suppose that $\text{ev}_1 \circ \mathcal{K}(u) = 0$ for some $u \in \widehat{U}$. Then $\mathcal{K}(u)$ is zero on o_L^\times because τ is surjective and because $\mathcal{K}(u)$ is Gal-continuous by Lemma 3.3.3. Hence $\mathcal{K}(u) = \psi_{\mathcal{C}} \varphi_{\mathcal{C}} \mathcal{K}(u)$. But $q \psi_{\mathcal{C}} \varphi_{\mathcal{C}} \mathcal{K}(u) = q \psi_{\mathcal{C}} \mathcal{K} \varphi^*(u) = \mathcal{K} \psi_{\text{Col}}^* \varphi^*(u)$ by Lemma 3.3.6, so $\mathcal{K}(qu - \psi_{\text{Col}}^* \varphi^*(u)) = 0$. Since \mathcal{K} is injective by assumption, $qu = \psi_{\text{Col}}^* \varphi^*(u) \in \psi_{\text{Col}}^*(\widehat{U})$. \square

Proposition 3.3.10. *Suppose that $\tau : G_L \rightarrow o_L^\times$ and $\mathcal{K}_1 : \widehat{U} \rightarrow o_\infty$ are both surjective, and that $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ is injective. Then*

$$\mathcal{K}_1^* : o_\infty^* \rightarrow S[[Z]]^{\psi_q=0}$$

is an S -linear bijection.

Proof. The image of $\mathcal{K}^* : o_\infty^* \rightarrow S[[Z]]$ is contained in $S[[Z]]^{\psi_q=0}$ by Lemma 3.3.8. If $\mathcal{K}_1^*(\ell) = 0$ for some $\ell \in o_\infty^*$, then $\ell \circ \mathcal{K}_1 = 0$ so $\ell(\mathcal{K}_1(\widehat{U})) = 0$. But $\mathcal{K}_1(\widehat{U}) = o_\infty$ by assumption, so $\ell = 0$. Hence \mathcal{K}_1^* is injective and it remains to prove it is also surjective.

Take some $F \in S[[Z]]^{\psi_q=0}$ and let $\ell := \langle -, F \rangle \in \widehat{U(\mathcal{G}_S)}^* \cong \widehat{U}^*$ be the S -valued o_L -linear functional on \widehat{U} given by Lemma 3.2.10 and Lemma 3.2.8(4). Then since $\psi_{\text{Col}}(F) = q \psi_q(F) = 0$,

$$0 = \langle u, \psi_{\text{Col}}(F) \rangle = \langle \psi_{\text{Col}}^*(u), F \rangle = \ell(\psi_{\text{Col}}^*(u)) \quad \text{for all } u \in \widehat{U}.$$

So, ℓ vanishes on $\psi_{\text{Col}}^*(\widehat{U})$ and hence also on $q \ker \mathcal{K}_1$ by Proposition 3.3.9. Since o_L has no q -torsion, we see that ℓ is zero on $\ker \mathcal{K}_1$. Hence ℓ descends to an S -valued o_L -linear functional on $\widehat{U} / \ker \mathcal{K}_1$. But this quotient is isomorphic

to o_∞ by assumption. So, we get a well-defined o_L -linear form $\bar{\ell} : o_\infty \rightarrow S$ such that $\bar{\ell}(\mathcal{K}_1(u)) = \ell(u)$ for all $u \in \widehat{U}$. Then

$$\langle u, \mathcal{K}_1^*(\bar{\ell}) \rangle = \bar{\ell}(\mathcal{K}_1(u)) = \ell(u) = \langle u, F \rangle \quad \text{for all } u \in \widehat{U}$$

which implies that $F = \mathcal{K}_1^*(\bar{\ell})$ by Lemma 3.2.8(4). Hence \mathcal{K}_1^* is surjective. \square

We make the following tentative

Conjecture 3.3.11. The map $\mathcal{K}_1 : \widehat{U} \rightarrow o_\infty$ is surjective and the map $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ is injective whenever τ is surjective.

3.4. The largest ψ_q -stable o_L -submodule of $o_L[[Z]]$. For brevity, we will write

$$A := S[[Z]]$$

in this subsection. The ψ_q -operator is only defined on A_L and it does *not* preserve A , in general.

Definition 3.4.1. Let $A^{\psi_q\text{-int}}$ be the largest S -submodule of A stable under ψ_q .

Remark 3.4.2. We have $A^{\psi_q\text{-int}} = \{F \in A : \psi_q^n(F) \in A \text{ for all } n \geq 0\}$.

Lemma 3.4.3. The image of $\mathcal{K}^* : \mathcal{C}^* \rightarrow A$ is contained in $A^{\psi_q\text{-int}}$.

Proof. Let $\lambda \in \mathcal{C}^*$. By Corollary 3.3.7, $\psi_q^n(\mathcal{K}^*(\lambda)) = \mathcal{K}^*((\psi_{\mathcal{C}}^*)^n(\lambda))$ lies in A for all $n \geq 0$. Now use Remark 3.4.2. \square

Clearly, $A^{\psi_q=0}$ is contained in $A^{\psi_q\text{-int}}$; moreover this last is φ -stable in view of Remark 3.4.2 and the fact that $\psi_q \circ \varphi = 1_A$ by Corollary 3.1.4. Therefore

$$S + \sum_{n=0}^{\infty} \varphi^n(A^{\psi_q=0}) \subseteq A^{\psi_q\text{-int}}.$$

Our next result makes this relation more precise; first we need some more notation.

Definition 3.4.4. We have the following *truncation operators*:

- (1) $s : \mathcal{C} \rightarrow \mathcal{C}$, given by $s(f) = f - f(0)1$, and
- (2) $t : A \rightarrow A$, given by $t(a) = a - a(0)1$.

It will be helpful to observe that $t\varphi = \varphi t$ as S -linear endomorphisms of A .

Proposition 3.4.5. There is a well-defined o_L -linear bijection

$$1 \oplus \sum_{n=0}^{\infty} \varphi^n t : o_L \oplus \prod_{n=0}^{\infty} A^{\psi_q=0} \xrightarrow{\cong} A^{\psi_q\text{-int}}.$$

Proof. Given any $(a_n)_n \in \prod_{n=0}^{\infty} A^{\psi_q=0}$, Lemma 3.1.7 implies that $\varphi^n(t(a_n)) \rightarrow 0$ as $n \rightarrow \infty$, because $t(a_n) \in ZA$ for all $n \geq 0$. Hence

$$(z, (a_n)_n) \mapsto z + \sum_{n=0}^{\infty} \varphi^n(t(a_n))$$

is a well-defined S -linear map $\gamma : S \oplus \prod_{n=0}^{\infty} A^{\psi_q=0} \rightarrow A$. Now $A^{\psi_q\text{-int}}$ is a t -stable S -submodule of A since $\psi_q(1) = 1$. Because $a_n \in A^{\psi_q=0}$, this implies that $\varphi^n(t(a_n)) = t\varphi^n(a_n) \in t(A^{\psi_q\text{-int}}) \subseteq A^{\psi_q\text{-int}}$ for any $n \geq 0$. Since $\psi_{\text{Col}} : A \rightarrow A$ is continuous by Lemma 3.1.6 and since $A^{\psi_q\text{-int}} = \{a \in A : \psi_{\text{Col}}^n(a) \in q^n A \text{ for all } n \geq 0\}$ by Remark 3.4.2, we see that $A^{\psi_q\text{-int}}$ is a closed S -submodule of A with respect to the $\langle \pi, Z \rangle$ -adic topology on $A = S[[Z]]$. Hence the image of γ is contained in $A^{\psi_q\text{-int}}$, and it remains to show that γ is bijective.

Suppose that $\gamma(z, (a_n)_n) = 0$ so that $z = -\sum_{n=0}^{\infty} \varphi^n(t(a_n))$. Since ZA is closed in A , this infinite sum lies in ZA . Since $S \cap ZA = 0$, we conclude that $z = 0$. Hence $a_0 = -\sum_{n=1}^{\infty} \varphi^n(t(a_n)) \in \varphi(A)$. But $a_0 \in A^{\psi_q=0}$ by definition, and

$$A^{\psi_q=0} \cap \varphi(A) = 0$$

because $\psi_q \circ \varphi = 1_A$ by Corollary 3.1.4. Hence $a_0 = 0$. Proceeding inductively on n , we quickly deduce that $a_n = 0$ for all $n \geq 0$ in a similar manner. Hence γ is injective.

Now let $a \in A^{\psi_q\text{-int}}$; then by definition, $\psi_q^n(a) \in A$ for all $n \geq 0$, so we can define

$$a_n := \psi_q^n(a) - \varphi\psi_q^{n+1}(a) \in A.$$

Since $\psi_q \circ \varphi = 1_A$ by Corollary 3.1.4, we see that $a_n \in A^{\psi_q=0}$ for all $n \geq 0$. Since $t\varphi = \varphi t$,

$$\sum_{n=0}^m \varphi^n(t(a_n)) = t\left(\sum_{n=0}^m \varphi^n(\psi_q^n(a) - \varphi\psi_q^{n+1}(a))\right) = t(a - \varphi^{m+1}\psi_q^{m+1}(a))$$

for any $m \geq 0$. Since $t\varphi^{m+1}\psi_q^{m+1}(a) = \varphi^{m+1}(t\psi_q^{m+1}(a)) \rightarrow 0$ as $m \rightarrow \infty$ by Lemma 3.1.7,

$$\gamma(a(0), (a_n)_n) = a(0) + t(a) - \lim_{m \rightarrow 0} \varphi^{m+1}(t\psi_q^{m+1}(a)) = a.$$

Hence γ is surjective. □

Lemma 3.4.6. *For each $n \geq 0$, there is a commutative diagram*

$$\begin{array}{ccccc} o_{\infty}^* & \xrightarrow{\text{ev}_{\pi^n}^*} & \mathcal{C}^* & \xrightarrow{s^*} & \mathcal{C}^* \\ \mathcal{K}_1^* \downarrow & & \downarrow \mathcal{K}^* & & \downarrow \mathcal{K}^* \\ A^{\psi_q=0} & \xrightarrow{\varphi^n} & A^{\psi_q\text{-int}} & \xrightarrow{t} & A^{\psi_q\text{-int}}. \end{array}$$

Proof. To see that the square on the left commutes, we use Corollary 3.3.7:

$$\varphi^n \mathcal{K}_1^* = \varphi^n \mathcal{K}^* \text{ev}_1^* = \mathcal{K}^* \varphi_{\mathcal{C}}^* \text{ev}_1^* = \mathcal{K}^* (\text{ev}_1 \varphi_{\mathcal{C}})^* = \mathcal{K}^* \text{ev}_{\pi^n}^*.$$

Hence in view of Lemma 3.2.8(4), it remains to show that

$$\langle u_m, \mathcal{K}^*(s^*(\lambda)) \rangle = \langle u_m, t(\mathcal{K}_1^*(\lambda)) \rangle \quad \text{for all } m \geq 0, \lambda \in \mathcal{C}^*.$$

Since t kills the constant term of a power series in A , we have

$$\langle u_m, t(a) \rangle = \delta_{m \geq 1} \langle u_m, a \rangle \quad \text{for all } a \in A.$$

Now $\mathcal{K}(u_m)(0) = P_m(0) = \delta_{m,0}$ by [28, Lemma 4.2] and $\mathcal{K}(u_0) = \mathcal{K}(1) = 1$, so

$$\begin{aligned} \langle u_m, \mathcal{K}^*(s^*(\lambda)) \rangle &= \lambda(s(\mathcal{K}(u_m))) \\ &= \lambda(\mathcal{K}(u_m) - \mathcal{K}(u_m)(0)1) = \delta_{m \geq 1} \lambda(\mathcal{K}(u_m)) = \langle u_m, t(\mathcal{K}^*(\lambda)) \rangle. \end{aligned}$$

The result follows. \square

$$\text{Let } c_0(o_\infty) := \{(x_n)_n \in \prod_{n=0}^{\infty} o_\infty : \lim_{n \rightarrow \infty} x_n = 0\}.$$

Lemma 3.4.7. *Suppose that τ is surjective. Then the map*

$$\eta : \mathcal{C} \rightarrow o_L \oplus c_0(o_\infty)$$

given by $\eta(f) = (f(0), (f(\pi^n) - f(0))_n)$ is an o_L -linear bijection.

Proof. Recall that any $f \in \mathcal{C}$ takes values in o_∞ by Lemma 3.3.2. Since $\pi^n \rightarrow 0$ as $n \rightarrow \infty$ in o_L and since f is continuous, $f(\pi^n) - f(0) \rightarrow 0$ as $n \rightarrow \infty$ in o_∞ . Thus η is well-defined.

Suppose $\eta(f) = 0$ for some $f \in \mathcal{C}$. Then $f(0) = 0$ and $f(\pi^n) = 0$ for all $n \geq 0$. Hence $f(\pi^n \tau(\sigma)) = \sigma(f(\pi^n)) = 0$ for all $\sigma \in G_L$, so f also vanishes on $\pi^n \tau(G_L)$ for each $n \geq 0$. Since τ is surjective, f vanishes on $\bigcup_{n=0}^{\infty} \pi^n o_L^\times \cup \{0\} = o_L$, so $f = 0$. Hence η is injective.

To show η is surjective, let $(z, (z_n)_n) \in o_L \oplus c_0(o_\infty)$ and define $f : o_L \rightarrow o_\infty$ by setting $f(0) = z$ and $f(\pi^n \tau(\sigma)) := z + \sigma(z_n)$ for all $n \geq 0$ and all $\sigma \in G_L$. This makes sense because τ is surjective, and if $\tau(\sigma) = \tau(\sigma')$ for some $\sigma, \sigma' \in G_L$ then $\sigma^{-1}\sigma' \in \ker \tau$ fixes o_∞ by Lemma 2.7.2, so $\sigma'(z_n) = \sigma(\sigma^{-1}\sigma'(z_n)) = \sigma(z_n)$ for any $n \geq 0$. It is easy to see that $f : o_L \rightarrow o_\infty$ is Gal-continuous and that $\eta(f) = (z, (z_n)_n)$. Hence η is surjective. \square

Lemma 3.4.7 allows us to give an explicit description of the space of Galois measures \mathcal{C}^* .

Corollary 3.4.8. *Suppose τ is surjective. Then*

$$\eta^* : o_L \oplus \prod_{n=0}^{\infty} o_\infty^* \rightarrow \mathcal{C}^*$$

is an o_L -linear bijection.

Proof. The functor $(-)^* = \text{Hom}_{o_L}(-, S)$ from o_L -modules to S -modules commutes with finite direct sums and sends $c_0(o_\infty)$ to $\prod_{n=0}^{\infty} o_\infty^*$. Now apply this functor to the isomorphism $\eta : \mathcal{C} \xrightarrow{\cong} o_L \oplus c_0(o_\infty)$ from Lemma 3.4.7. \square

Theorem 3.4.9. *Suppose that τ is surjective and that $\mathcal{K}_1^* : o_\infty^* \rightarrow A^{\psi_q=0}$ is an isomorphism. Then $\mathcal{K}^* : \mathcal{C}^* \rightarrow A^{\psi_q\text{-int}}$ is an isomorphism as well.*

Proof. Using Corollary 3.4.8 and Proposition 3.4.5, we can build the following diagram:

$$\begin{array}{ccc}
 S \oplus \prod_{n=0}^{\infty} o_{\infty}^* & \xrightarrow{\eta^*} & \mathcal{C}^* \\
 \downarrow 1 \oplus \prod_{n=0}^{\infty} \mathcal{K}_1^* & & \downarrow \mathcal{K}^* \\
 S \oplus \prod_{n=0}^{\infty} A^{\psi_q=0} & \xrightarrow{1 \oplus \sum_{n=0}^{\infty} \varphi^n t} & A^{\psi_q\text{-int}}.
 \end{array}$$

Note that we can write $\eta = \text{ev}_0 \oplus (\text{ev}_{\pi^n} \circ s)_n$. Lemma 3.4.6 implies that

$$\mathcal{K}^*(\text{ev}_{\pi^n} \circ s)^* = \mathcal{K}^* s^* \text{ev}_{\pi^n}^* = t \varphi^n \mathcal{K}_1^* = \varphi^n t \mathcal{K}_1^* \quad \text{for any } n \geq 0.$$

Using $P_m(0) = \delta_{m,0}$ again together with (6), we also have

$$\mathcal{K}^*(\eta^*(1, (0)_n)) = \mathcal{K}^*(\text{ev}_0^*(1)) = \sum_{m=0}^{\infty} \text{ev}_0^*(1)(P_m(-\Omega))Z^m = \sum_{m=0}^{\infty} P_m(0)Z^m = 1.$$

So the diagram is commutative. Now η^* is an isomorphism by Corollary 3.4.8, and the bottom map is an isomorphism by Proposition 3.4.5. Since \mathcal{K}_1^* is an isomorphism by assumption, the vertical map on the left is an isomorphism. Hence \mathcal{K}^* is also an isomorphism by the commutativity of the diagram. \square

Corollary 3.4.10. *Let S be any π -adically complete o_L -algebra. The dual Katz map*

$$\mathcal{K}^* : \mathcal{C}^* \rightarrow S[[Z]]^{\psi_q\text{-int}}$$

is an isomorphism if $\tau : G_L \rightarrow o_L^\times$ and $\mathcal{K}_1 : \widehat{U} \rightarrow o_\infty$ are surjective, and $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ is injective.

Proof. Apply Theorem 3.4.9 together with Proposition 3.3.10. \square

Remark 3.4.11. Katz claims on [19, p. 60] that it is easy to show that the map that he denotes by $(**)$ on p.59, is injective. In our notation, this map is \mathcal{K}^* ; at least when τ is surjective, the proof of Theorem 3.4.9 shows that its injectivity is equivalent to the injectivity of \mathcal{K}_1^* , which is equivalent to $\mathcal{K}_1(\widehat{U}) \cdot L = L_\infty$ in view of the proof of Proposition 3.3.10. We were only able to establish the equality $\mathcal{K}_1(\widehat{U}) = o_\infty$ in the case where $L = \mathbb{Q}_{p^2}$ by carrying out an explicit computation — see Proposition 3.6.5 below.

3.5. The Newton polygon of $\Delta_1(Z) - 1$. In this section, we obtain some estimates on $v_\pi(P_k(\Omega))$, $k \geq 1$. Recall that d and e and f denote the degree and ramification and inertia indices of L/\mathbb{Q}_p , respectively.

Lemma 3.5.1. *If $k \geq 0$ and $1 \leq r \leq e$, then we have an isomorphism of abelian groups*

$$o_L/\pi^{ek+r}o_L \cong (\mathbb{Z}/p^k\mathbb{Z})^{f(e-r)} \oplus (\mathbb{Z}/p^{k+1}\mathbb{Z})^{fr}.$$

Proof. Note that $po_L = \pi^e o_L \subseteq \pi^r o_L$ since $e \geq r$ by assumption, so $o_L/\pi^r o_L$ is an elementary abelian p -group of order $|o_L/\pi^r o_L| = p^{fr}$. Hence, using the elementary divisors theorem, we can find $v_1, \dots, v_d \in o_L$ such that

$$o_L = \mathbb{Z}_p v_1 \oplus \dots \oplus \mathbb{Z}_p v_d \quad \text{and} \quad \pi^r o_L = \bigoplus_{i=1}^s \mathbb{Z}_p v_i \oplus \bigoplus_{i=s+1}^d \mathbb{Z}_p p v_i$$

for some integer s with $1 \leq s \leq d$. We deduce that $fr = d - s$, so $s = f(e - r)$. Since $\pi^{ek+r} o_L = p^k \pi^r o_L$, the result now follows easily. \square

Lemma 3.5.2. *In $o_L/\pi^{ek+r} o_L$, the image of 1 has order p^{k+1} .*

Proof. This can be proved directly as $p^k \cdot 1 \in \pi^{ek} \cdot o_L^\times \neq 0$ in $o_L/\pi^{ek+r} o_L$. \square

Definition 3.5.3. Let $m \geq 0$.

- (1) Let $k_m = \lfloor (m-1)/e \rfloor$, so that $m = ek_m + r$ with $1 \leq r \leq e$.
- (2) Define $x_m := q^m/p^{k_m+1}$.
- (3) Define

$$y_0 = \frac{e}{p-1} - \frac{1}{q-1} \quad \text{and} \quad y_m = \frac{e}{p-1} - \sum_{j=1}^{m-1} \frac{1}{p^{k_j+1}} - \frac{q}{p^{k_m+1}(q-1)}.$$

For example, $x_0 = 1$ and $x_1 = q/p$. Note that if $m = en + r$ with $1 \leq r \leq e$, then

$$y_{en+r} = \frac{e}{p^n(p-1)} - \frac{r}{p^{n+1}} - \frac{1}{(q-1)p^{n+1}}.$$

Theorem 3.5.4. *The vertices of the Newton polygon of $\Delta_1(Z) - 1$ (using the valuation v_π , and excluding the point $(0, +\infty)$) are the points (x_m, y_m) for $m \geq 0$.*

Proof. Via the Schneider-Teitelbaum isomorphism $\kappa : \mathbf{B}(\mathbb{C}_p) \xrightarrow{\cong} \mathfrak{X}(\mathbb{C}_p)$, the zeroes of the power series

$$\Delta_1(Z) - 1 = \sum_{m=1}^{\infty} P_m(\Omega) Z^m \in o_{\mathbb{C}_p}[[Z]]$$

are the $z \in \mathfrak{m}_{\mathbb{C}_p}$ such that $\kappa(z)$ is an L -analytic character satisfying $\kappa(z)(1) = 1$. These characters are torsion³, and correspond to some of the torsion points of the Lubin-Tate group \mathcal{G} . There are precisely q^m points in $\mathcal{G}[\pi^m]$, and the common valuation of each point $z \in \mathcal{G}[\pi^m] \setminus \mathcal{G}[\pi^{m-1}]$ is $v_\pi(z) = 1/q^{m-1}(q-1)$.

If we write $m = ek + r$ as above, then in view of Lemma 3.5.1 and Lemma 3.5.2 there are $x_m = q^m/p^{k_m+1}$ elements $z \in \mathcal{G}[\pi^m]$ such that $\kappa(z)(1) = 1$.

Let $((x'_m, y'_m))_{m=0}^\infty$ be the vertices of the Newton polygon, so that the first vertex is $(x'_0, y'_0) = (1, v_\pi(\Omega)) = (x_0, y_0)$. The slope of the line segment between (x'_{m-1}, y'_{m-1}) and (x'_m, y'_m) is minus the common valuation of the elements of

³Suppose that $\lambda : o_L \rightarrow \mathbb{C}_p^\times$ is a locally L -analytic character such that $\lambda(1) = 1$. Then $\lambda(a) = 1$ for all $a \in \mathbb{Z}_p$. Hence $\lambda'(1) = 0$. Since λ is locally L -analytic, λ' is L -linear, and hence $\lambda' = 0$ so that λ is locally constant, and hence torsion.

$z \in \mathcal{G}[\pi^m] \setminus \mathcal{G}[\pi^{m-1}]$ satisfying $\kappa(z) = 1$, that is $1/q^{m-1}(q-1)$. Hence $x'_m = x_m$ for all $m \geq 0$. Using the definitions of x_m and y_m , we have the formula

$$y_m = y_0 - \frac{x_1 - x_0}{q^{1-1}(q-1)} - \cdots - \frac{x_m - x_{m-1}}{q^{m-1}(q-1)}$$

which implies that $y'_m = y_m$ for all $m \geq 0$. \square

Remark 3.5.5. Note that $y_m \rightarrow 0$ as $m \rightarrow +\infty$. This is consistent with the fact that $\|\Delta_1(Z) - 1\| = 1$.

Corollary 3.5.6. *We have the following formulas for $v_\pi(P_k(\Omega))$.*

- (1) For all $m \geq 0$, we have $v_\pi(P_{x_m}(\Omega)) = y_m$.
- (2) For all $n \geq 0$, we have $v_\pi(P_{p^{n(d-1)}}(\Omega)) = 1/p^n \cdot v_\pi(\Omega)$.

Proof. Item (1) follows immediately from Theorem 3.5.4. Item (2) follows from item (1) with $m = en$. Indeed, $x_{en} = q^{en}/p^n = p^{n(d-1)}$ and

$$y_{en} = \frac{e}{p-1} - \frac{e}{p} - \frac{e}{p^2} - \cdots - \frac{e}{p^{n-1}} - \frac{e-1}{p^n} - \frac{q}{p^n(q-1)} = \frac{1}{p^n} \cdot \left(\frac{e}{p-1} - \frac{1}{q-1} \right). \quad \square$$

Remark 3.5.7. If L/\mathbb{Q}_p is unramified, then item (2) of Corollary 3.5.6 gives all the valuations of the $P_k(\Omega)$ that can be computed using the Newton polygon. For $n \geq 0$, we get

$$\text{val}_p(P_{p^{n(d-1)}}(\Omega)) = 1/p^n \cdot v_\pi(\Omega) = \frac{1}{p^{n-1}(p-1)} \cdot \frac{q/p-1}{q-1}.$$

Corollary 3.5.8. *Suppose that $L = \mathbb{Q}_{p^2}$ and $\pi = p$. Then we have*

$$\text{val}_p(P_{p^k}(\Omega)) = \frac{1}{p^{k-1}(q-1)} \quad \text{for all } k \geq 1,$$

and if $k \geq 1$ and $p^{k-1} \leq m \leq p^k$, then

$$\text{val}_p(P_m(\Omega)) \geq \frac{1}{p^{k-1}(q-1)} + \frac{p^k - m}{q^{k-1}(q-1)} = \frac{1}{p^{k-2}(q-1)} - \frac{m - p^{k-1}}{q^{k-1}(q-1)}.$$

3.6. Verifying Conjecture 3.3.11 in a special case.

Definition 3.6.1. Fix $m \geq 1$.

- (1) Let $\mathcal{G}_m = \mathcal{G}[\pi^m]$ be the finite flat o_L -group scheme of π^m -torsion points in the Lubin-Tate formal group \mathcal{G} .
- (2) Let \mathcal{G}'_m be the Cartier dual of \mathcal{G}_m .
- (3) Let $U(m) := \mathcal{O}(\mathcal{G}'_m) = \text{Hom}_{o_L}(o_L[[Z]]/\langle \varphi^m(Z) \rangle, o_L)$.
- (4) Let $\mathcal{G}' := \text{colim } \mathcal{G}'_m$ be the dual p -divisible group to the p -divisible group defined by the formal group \mathcal{G} .

Recall that by Cartier duality — see [32, p. 177] — the period $\Omega \in \mathbb{C}_p$ corresponds to a choice of generator $t' \in T_p \mathcal{G}' = T_\pi \mathcal{G}'$ as an o_L -module. We recall how this correspondence works. First, the element

$$\Delta_1 = \sum_{n=0}^{\infty} P_n(\Omega) Z^n \in o_{\mathbb{C}_p}[[Z]]$$

gives a compatible system of group-like elements $(\Delta_1(m))_{m=1}^\infty \in \prod_{m=1}^\infty \mathcal{O}(\mathcal{G}_m)$, where $\Delta_1(m)$ is the image of Δ_1 in $\mathcal{O}(\mathcal{G}_m \times_{o_L} o_{\mathbb{C}_p}) = o_{\mathbb{C}_p}[[Z]]/\langle \varphi^m(Z) \rangle$ under the natural surjective homomorphism of $o_{\mathbb{C}_p}$ -algebras $o_{\mathbb{C}_p}[[Z]] \twoheadrightarrow \mathcal{O}(\mathcal{G}_m \times_{o_L} o_{\mathbb{C}_p})$. Since $\mathcal{O}(\mathcal{G}_m \times_{o_L} o_{\mathbb{C}_p})$ can be identified with $\text{Hom}_{o_{\mathbb{C}_p}}(\mathcal{O}(\mathcal{G}'_m \times_{o_L} o_{\mathbb{C}_p}), o_{\mathbb{C}_p})$, $\Delta_1(m)$ can be viewed as an $o_{\mathbb{C}_p}$ -linear map $U(m) \otimes_{o_L} o_{\mathbb{C}_p} \rightarrow o_{\mathbb{C}_p}$ which is in fact an $o_{\mathbb{C}_p}$ -algebra homomorphism because $\Delta_1(m)$ is group-like. This map is determined by its restriction to $U(m)$; this restriction is an o_L -algebra homomorphism $t'_m : U(m) \rightarrow o_{\mathbb{C}_p}$ and is therefore an element of $\mathcal{G}'_m(\mathbb{C}_p)$.

Finally, the multiplication-by- π -maps $\mathcal{G}'_{m+1}(\mathbb{C}_p) \rightarrow \mathcal{G}'_m(\mathbb{C}_p)$ in the inverse system defining the Tate module $T_\pi \mathcal{G}'$ are induced by the inclusions of o_L -algebras $U(m) \hookrightarrow U(m+1)$, so $t'_{m+1|U(m)} = t'_m$ for all $m \geq 1$, and the generator $t' \in T_\pi \mathcal{G}'$ is given by $t' = (t'_m)_{m=1}^\infty \in \prod_{m=1}^\infty \mathcal{G}'_m(\mathbb{C}_p)$.

Lemma 3.6.2. *Let $m \geq 1$. The restriction of \mathcal{K}_1 to $U(m) \subset \widehat{U}$ is equal to t'_m .*

Proof. Recall that we have identified \widehat{U} with $o_L[[Z]]_{\text{cts}}^*$ using Lemma 3.2.8(3). Let $u \in U(m)$ and let $\tilde{u} \in \widehat{U}$ be the corresponding o_L -linear map $o_L[[Z]] \rightarrow o_L$ which kills $\langle \varphi^m(Z) \rangle$. Then

$$t'_m(u) = \Delta_1(m)(u) = \langle \tilde{u}, \Delta_1 \rangle = \mathcal{K}(\tilde{u})(1) = \mathcal{K}_1(\tilde{u})$$

and the result follows. \square

For each $m \geq 1$, let L_m be the finite Galois extension of L contained in $L_\infty = \mathbb{C}_p^{\ker \tau}$ defined by $\text{Gal}(L_\infty/L_m) = \tau^{-1}(1 + \pi^m o_L)$.

Lemma 3.6.3. *Let $m \geq 1$. Then $t'_m(U(m)) \subseteq o_{L_m}$.*

Proof. Let $\sigma \in \text{Gal}(L_\infty/L_m)$ so that $\tau(\sigma) \in 1 + \pi^m o_L$. Then by definition of the character τ , σ acts trivially on $\mathcal{G}'_m(\mathbb{C}_p)$. In other words, $\sigma(t'_m(u)) = t'_m(u)$ for all $u \in U(m)$ and hence $t'_m(U(m)) \subseteq L_\infty^{\text{Gal}(L_\infty/L_m)} = L_m$. But $U(m)$ is a finitely generated o_L -module so $t'_m(U(m))$ is integral over o_L and is therefore contained in o_{L_m} . \square

Recall from Definition 3.2.1(2) that $U = U(\mathcal{G})$ is the covariant bialgebra of the formal group \mathcal{G} .

Definition 3.6.4. For each $m \geq 1$, let $U(m)_k := \text{im}(U(m) \rightarrow \widehat{U}/\pi\widehat{U})$.

We will identify $U_k := U/\pi U$ with $\widehat{U}/\pi\widehat{U}$ via the natural map $U/\pi U \rightarrow \widehat{U}/\pi\widehat{U}$ and we regard $U(m)_k$ as being naturally embedded into $U(m+1)_k$.

Proposition 3.6.5. *Suppose that $t'_m(U(m)) = o_{L_m}$ for all $m \geq 1$. Then $\mathcal{K}_1 : \widehat{U} \rightarrow o_\infty$ is surjective.*

Proof. Consider $o_\tau := \overline{L} \cap o_\infty$. Since o_τ is π -adically dense in o_∞ , to prove that $\mathcal{K}_1(\widehat{U})$ contains o_∞ , it is enough to prove that it contains o_τ . Fix $m \geq 1$. By Lemma 3.6.2, the restriction of $\mathcal{K}_1 : \widehat{U} \rightarrow o_{\mathbb{C}_p}$ to $U(m)$ is equal to t'_m .

Hence by assumption $o_{L_m} = t'_m(U(m)) = \mathcal{K}_1(U(m))$, so $o_r = \bigcup_{m \geq 1} o_{L_m}$ is also contained in $\mathcal{K}_1(\widehat{U})$. \square

Lemma 3.6.6. *For each $m \geq 1$, we have $U(m) + \pi\widehat{U} = \sum_{r=0}^{q^m-1} o_L u_r + \pi\widehat{U}$.*

Proof. Let $u \in U(m)$ and let $\tilde{u} : o_L[[Z]] \rightarrow o_L$ be the corresponding o_L -linear form which vanishes on $\langle \varphi^m(Z) \rangle$. Consider $v := \tilde{u} - \sum_{r=0}^{q^m-1} \tilde{u}(Z^r) u_r \in \widehat{U}$. For each $r < q^m$, u_r sends $\langle \varphi^m(Z) \rangle$ into πo_L because $\varphi^m(Z) \equiv Z^{q^m} \pmod{\pi o_L[[Z]]}$. Since \tilde{u} kills $\langle \varphi^m(Z) \rangle$, we see that v also sends $\langle \varphi^m(Z) \rangle$ into πo_L . By construction, v is zero on $1, Z, \dots, Z^{q^m-1}$. Since

$$(7) \quad o_L 1 \oplus o_L Z \oplus \dots \oplus o_L Z^{q^m-1} \oplus \langle \varphi^m(Z) \rangle = o_L[[Z]],$$

we conclude that $v(o_L[[Z]]) \subseteq \pi o_L$ and hence $v = \pi w$ for some o_L -linear form $w : o_L[[Z]] \rightarrow o_L$. Since $v : o_L[[Z]] \rightarrow o_L$ is continuous for the weak topology on $o_L[[Z]]$, so is w . Hence $w \in \widehat{U}$ and hence $\tilde{u} \in \sum_{r=0}^{q^m-1} o_L u_r + \pi\widehat{U}$. This shows that \subseteq holds.

For the reverse containment, it is enough to show that $u_r \in U(m) + \pi\widehat{U}$ for each $r = 0, \dots, q^m - 1$. Using (7), define an o_L -linear form $w_r : o_L[[Z]] \rightarrow o_L$ which is zero on $\langle \varphi^m(Z) \rangle$ and which sends Z^i to $\delta_{i,r}$ for each $0 \leq i < q^m$. Since u_r sends $\langle \varphi^m(Z) \rangle$ into πo_L , the same is true of $u_r - w_r$. Since $u_r - w_r$ is zero on $1, Z, \dots, Z^{q^m-1}$ by construction, we see that $u_r - w_r$ sends all of $o_L[[Z]]$ into πo_L . Hence $u_r - w_r = \pi v_r$ for some o_L -linear form $v_r : o_L[[Z]] \rightarrow o_L$. Since $u_r - w_r$ is continuous for the weak topology on $o_L[[Z]]$, so is v_r . Because w_r is zero on $\langle \varphi^m(Z) \rangle$, it lies in $U(m)$ and hence $u_r = w_r + \pi v_r \in U(m) + \pi\widehat{U}$. \square

Proposition 3.6.7. *If $L = \mathbb{Q}_{p^2}$, then $t'_m(U(m)) = o_{L_m}$ for all $m \geq 1$.*

Proof. Fix $m \geq 1$. By Lemma 3.6.6, for each $0 \leq r < q^m$ we can find $w_r \in U(m)$ such that $w_r - u_r \in \pi\widehat{U}$. Set $r := p^{2m-1} = pq^{m-1} < q^m$. Note that $\mathcal{K}_1(u_r) = \mathcal{K}(u_r)(1) = \langle u_r, \Delta_1 \rangle = P_r(\Omega)$. Since $L = \mathbb{Q}_{p^2}$, Corollary 3.5.8 applied with $k = 2m - 1$ tells us that

$$\text{val}_p(\mathcal{K}_1(u_r)) = \text{val}_p(P_r(\Omega)) = \frac{1}{p^{2m-2}(q-1)} = \frac{1}{q^{m-1}(q-1)} = [L_m : L]^{-1} < 1.$$

Now $\pi o_L = po_L$ since $L = \mathbb{Q}_{p^2}$, so $\mathcal{K}_1(u_r - w_r) \in \mathcal{K}_1(\pi\widehat{U}) \subseteq po_{\mathbb{C}_p}$ since \mathcal{K}_1 takes values in $o_{\mathbb{C}_p}$. Hence $\text{val}_p(\mathcal{K}_1(u_r) - \mathcal{K}_1(w_r)) \geq 1$ and $\text{val}_p(\mathcal{K}_1(w_r)) = \text{val}_p(\mathcal{K}_1(u_r)) = [L_m : L]^{-1}$. Therefore $\mathcal{K}_1(w_r)$ is a uniformiser in L_m and the result follows. \square

Now we start to explore the injectivity of $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$.

Lemma 3.6.8. *For each $m \geq 1$, we have $U(m) \cap \pi\widehat{U} = \pi U(m)$.*

Proof. Let $g = \pi h \in U(m)$ for some $h \in \widehat{U}$. Then $\pi\langle h, F \rangle = \langle \pi h, F \rangle = 0$ for any $F \in \langle \varphi^m(Z) \rangle$. Hence $\langle h, F \rangle = 0$ for all such F as well, so $h \in U(m)$ and $g \in \pi U(m)$. \square

Corollary 3.6.9. *The map $\mathcal{O}(\mathcal{G}'_m \times_{o_L} k) = U(m)/\pi U(m) \rightarrow U(m)_k$ is an isomorphism.*

Since \mathcal{G}' forms a p -divisible group, we have a closed immersion $\mathcal{G}'_m \rightarrow \mathcal{G}'_{m+1}$ for each $m \geq 1$. The comorphism of this map $\mathcal{O}(\mathcal{G}'_{m+1}) \rightarrow \mathcal{O}(\mathcal{G}'_m)$ is the dual of the o_L -Hopf algebra map $\mathcal{O}(\mathcal{G}_m) \rightarrow \mathcal{O}(\mathcal{G}_{m+1})$ induced by $\varphi : \mathcal{O}(\mathcal{G}) \rightarrow \mathcal{O}(\mathcal{G})$. Using Corollary 3.6.9, we obtain connecting maps $\varphi_k^* : U(m+1)_k \rightarrow U(m)_k$.

Lemma 3.6.10. *The comorphisms $\varphi_k^* : U(m+1)_k \rightarrow U(m)_k$ are surjective for all $m \geq 1$.*

Proof. By Corollary 3.6.9, $U(m)_k$ is isomorphic as a k -vector space to

$$\mathcal{O}(\mathcal{G}'_m \times_{o_L} k) = \text{Hom}_k(\mathcal{O}(\mathcal{G}_m \times_{o_L} k), k).$$

Since $\varphi(Z) \equiv Z^q \pmod{\pi o_L \llbracket Z \rrbracket}$, we have $\mathcal{O}(\mathcal{G}_m \times_{o_L} k) = k \llbracket Z \rrbracket / \langle Z^{q^m} \rangle$ and the k -algebra homomorphism $\varphi_k : k \llbracket Z \rrbracket / \langle Z^{q^m} \rangle \rightarrow k \llbracket Z \rrbracket / \langle Z^{q^{m+1}} \rangle$ which sends Z to Z^q is *injective*. Hence the dual map

$$\varphi_k^* : \text{Hom}_k(k \llbracket Z \rrbracket / \langle Z^{q^{m+1}} \rangle, k) \rightarrow \text{Hom}_k(k \llbracket Z \rrbracket / \langle Z^{q^m} \rangle, k)$$

is surjective and the result follows. \square

Next we consider an ideal I of U_k and we set $I(m) := I \cap U(m)$ for all $m \geq 1$. We assume that I is φ^* -stable, in the sense that $\varphi^*(I) \subseteq I$.

Proposition 3.6.11. *Suppose that I is a φ^* -stable ideal of U_k such that $\varprojlim \frac{U(m)_k}{I(m)}$ is finite dimensional over k . Then $U_k/I = \text{colim} \frac{U(m)}{I(m)}$ is also finite dimensional over k .*

Proof. Let $m \geq 1$ and consider the short exact sequence

$$0 \rightarrow I(m) \rightarrow U(m)_k \rightarrow U(m)_k/I(m) \rightarrow 0.$$

Since I is φ^* -stable by assumption, we get a short exact sequence of towers of finite-dimensional k -vector spaces. Passing to the inverse limit therefore gives an exact sequence

$$0 \rightarrow I(\infty) := \varprojlim I(m) \rightarrow \varprojlim U(m)_k \rightarrow \varprojlim \frac{U(m)_k}{I(m)} \rightarrow 0.$$

By assumption, the term on the right is a finite dimensional k -vector space. We see from Lemma 3.6.10 that the connecting maps $U(m+1)_k/I(m+1) \rightarrow U(m)_k/I(m)$ induced by φ^* are *surjective*. Therefore, for large m , all of these maps are necessarily isomorphisms, and hence there exists $m_0 \geq 1$ such that

$$\dim \frac{U(m+1)_k}{I(m+1)} = \dim \frac{U(m)_k}{I(m)} \quad \text{for all } m \geq m_0.$$

Now the definition of $I(m)$ shows that the natural connecting maps in the opposite direction $U(m)_k/I(m) \rightarrow U(m+1)_k/I(m+1)$ is *injective* for any $m \geq 1$. So they are isomorphisms whenever $m \geq m_0$. The result follows. \square

Proposition 3.6.12. *Let $J = \ker \mathcal{K}$ and let $I := (J + \pi\widehat{U})/\pi\widehat{U}$ be its image in U_k . Then I is a φ^* -stable ideal in U_k such that $\dim U_k/I = \infty$.*

Proof. Since \mathcal{K} is an o_L -algebra homomorphism by Lemma 3.3.5, J is an ideal in \widehat{U} . Since $\mathcal{K}\varphi^* = \varphi_{\mathcal{C}}\mathcal{K}$ by Lemma 3.3.6, this ideal is in fact φ^* -stable. Hence its image I in U_k is also φ^* -stable.

Suppose that $h \in \widehat{U}$ and $r \geq 1$ are such that $\pi^r h \in J$. Then $\mathcal{K}(\pi^r h) = 0$ in \mathcal{C} , so $\mathcal{K}(h) = 0$ as well. So $J \cap \pi^r \widehat{U} = \pi^r J$ for all $r \geq 1$. Now consider the short exact sequence

$$0 \rightarrow J \rightarrow \widehat{U} \rightarrow \mathcal{K}(U) \rightarrow 0.$$

Equip both \widehat{U} and $\mathcal{K}(U)$ with the π -adic filtrations. Then the above shows that the subspace filtration on J induced by the π -adic filtration on \widehat{U} coincides with the π -adic filtration on J . Therefore we get a short exact sequence of $\text{gr } o_L$ -modules

$$0 \rightarrow \text{gr } J \rightarrow \text{gr } \widehat{U} \rightarrow \text{gr } \mathcal{K}(U) \rightarrow 0.$$

So, if $\dim U_k/I < \infty$, then $\text{gr } \widehat{U}/\text{gr } J \cong (U_k/I)[\text{gr } \pi]$ is a finitely generated module over $\text{gr } o_L$, so $\text{gr } \mathcal{K}(U)$ is a finitely generated $\text{gr } o_L$ -module. The π -adic filtration on \mathcal{C} is separated, hence the π -adic filtration on $\mathcal{K}(U)$ is also separated. Therefore $\mathcal{K}(U)$ is a finitely generated o_L -module by [22, Chapter I, Theorem 5.7]. Hence $\dim_L \mathcal{K}(U[1/\pi]) < \infty$. But this contradicts [28, Theorem 4.7]: the space of locally L -analytic Gal-continuous functions is not finite dimensional over L since it contains the subspace of locally constant Gal-continuous functions, which is infinite dimensional over L . \square

Corollary 3.6.13. *If $d := [L : \mathbb{Q}_p] = 2$, then $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ is injective.*

Proof. By Proposition 3.6.12, $I = (\ker \mathcal{K} + \pi\widehat{U})/\pi\widehat{U}$ is a φ^* -stable ideal in U_k of infinite codimension in U_k . Hence $I(\infty) := \varprojlim (I \cap U(m)_k)$ is an ideal of infinite codimension in $\varprojlim U(m)_k$ by Proposition 3.6.11. By [18, Example 2.5.3], the Dieudonné module $M(\mathcal{G}_k)$ associated with the Lubin-Tate formal group $\mathcal{G}_k = \mathcal{G} \times_{o_L} k$ over the perfect field k has basis $\{\gamma, V\gamma, \dots, V^{d-1}\gamma\}$ over the ring of p -typical Witt vectors $\mathbb{W}(k)$ for a certain element $\gamma \in M(\mathcal{G}_k)$, and satisfies $V^d = p$. Hence the Verschiebung operator V on $M(\mathcal{G}_k)$ is topologically nilpotent. Therefore the Cartier dual \mathcal{G}'_k is connected. Hence $\varprojlim U(m)_k \cong \mathcal{O}(\mathcal{G}' \times_{o_L} k)$ is isomorphic to $k[[X_1, \dots, X_{d-1}]]$ by [32, Propositions 1 and 3]. Since $d = 2$, we conclude that $I(\infty) = 0$. Hence $I(m) = 0$ for all $m \geq 1$ and hence $I = 0$. So $\ker \mathcal{K} = 0$ as well. \square

Theorem 3.6.14. *Suppose that $L = \mathbb{Q}_{p^2}$. Then*

$$\mathcal{K}_1^* : o_{\infty}^* \rightarrow o_L[[Z]]^{\psi_q=0}$$

is an o_L -linear bijection.

Proof. Since $d = 2$, we know that τ is surjective by Lemma 2.6.4. Then $\mathcal{K} : \widehat{U} \rightarrow \mathcal{C}$ is injective by Corollary 3.6.13 and $\mathcal{K}_1 : \widehat{U} \rightarrow o_\infty$ is surjective by Proposition 3.6.5 and Proposition 3.6.7. Now apply Proposition 3.3.10. \square

We can now prove Theorem 1.6.1 from the Introduction. In fact, we prove the following more general version, from which Theorem 1.6.1 follows as a special case by setting $S = o_K$.

Theorem 3.6.15. *Let $L = \mathbb{Q}_{p^2}$ and let S be a π -adically complete o_L -algebra.*

- (1) *The map $\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), S) \rightarrow S[[Z]]$ is injective.*
- (2) *Its image is equal to $S[[Z]]^{\psi_q\text{-int}}$.*

Proof. Since $d = 2$, τ is surjective by Lemma 2.6.4. By Theorem 3.6.14, the map $\mathcal{K}_1^* : o_\infty^* \rightarrow o_L[[Z]]^{\psi_q=0}$ is an isomorphism. Now apply Theorem 3.4.9. \square

4. INTEGER-VALUED POLYNOMIALS

4.1. The algebraic dual of $\mathcal{O}^\circ(\mathfrak{X}_K)$. Recall from §2.1 that our coefficient field K is a complete field extension of L contained in \mathbb{C}_p . Pick a basis $\{v_1, \dots, v_d\}$ for o_L as a \mathbb{Z}_p -module with $v_1 = 1$. We view o_L as a p -valued group with p -valuation ω given by

$$\omega\left(\sum_{i=1}^d \lambda_i v_i\right) = 1 + \min_{1 \leq i \leq d} \text{val}_p(\lambda_i).$$

Let r be a real number in the range $1/p \leq r < 1$. Recall from [29, §4] that $D^{\mathbb{Q}_p\text{-an}}(o_L, K)$ carries a norm $\|\cdot\|_r$ given by

$$(8) \quad \left\| \sum_{\alpha \in \mathbb{N}^d} d_\alpha \mathbf{b}^\alpha \right\|_r = \sup_{\alpha \in \mathbb{N}^d} |d_\alpha| r^{|\alpha|}.$$

where $b_i := \delta_{v_i} - 1 \in D^{\mathbb{Q}_p\text{-an}}(o_L, K)$ for $i = 1, \dots, d$, $\mathbf{b}^\alpha = b_1^{\alpha_1} \dots b_d^{\alpha_d} \in D^{\mathbb{Q}_p\text{-an}}(o_L, K)$ and $|\alpha| = \tau\alpha = \alpha_1 + \dots + \alpha_d$ for all $\alpha \in \mathbb{N}^d$.

Definition 4.1.1. Let $1/p \leq r < 1$.

- (1) Let $D_r^{\mathbb{Q}_p\text{-an}}(o_L, K)$ denote the completion of $D^{\mathbb{Q}_p\text{-an}}(o_L, K)$ with respect to $\|\cdot\|_r$.
- (2) Let $\mathfrak{X}_0(r)_K := \text{Sp } D_r^{\mathbb{Q}_p\text{-an}}(o_L, K)$.
- (3) Let $\mathfrak{X}(r)_K := \mathfrak{X}_K \cap \mathfrak{X}_0(r)_K = \text{Sp } D_r^{L\text{-an}}(o_L, K)$, where $D_r^{L\text{-an}}(o_L, K)$ is the factor algebra of $D_r^{\mathbb{Q}_p\text{-an}}(o_L, K)$ by the ideal generated by

$$u_2 - v_2 u_1, \quad u_3 - v_3 u_1, \quad \dots, \quad u_d - v_d u_1$$

where $u_i := \log(1 + b_i) \in D^{\mathbb{Q}_p\text{-an}}(o_L, K)$.

As r approaches 1 from below, the K -affinoid varieties $\mathfrak{X}(r)_K$ form an increasing family of K -affinoid subvarieties of \mathfrak{X}_K : whenever $1/p \leq r < r' < 1$ we have

$$(9) \quad \mathbb{1} \in \mathfrak{X}(1/p)_K \subset \dots \subset \mathfrak{X}(r)_K \subset \mathfrak{X}(r')_K \subset \dots \subset \mathfrak{X}_K = \bigcup_{1/p \leq r < 1} \mathfrak{X}(r)_K.$$

Here $\mathbb{1} \in \mathfrak{X}_K$ is the *trivial character*: the ideal generated by b_1, \dots, b_d .

Lemma 4.1.2. *The completed local ring $\widehat{\mathcal{O}_{\mathfrak{X}_K, \mathbb{1}}}$ of \mathfrak{X} at $\mathbb{1}$ is isomorphic to a power series ring in one variable $b := b_1$ over K :*

$$\widehat{\mathcal{O}_{\mathfrak{X}, \mathbb{1}}} \cong K[[b]].$$

Proof. We have $\mathcal{O}(\mathfrak{X}_0(1/p)_K) = K\langle b_1/p, \dots, b_d/p \rangle = K\langle u_1/p, \dots, u_d/p \rangle$.

Quotienting out by the ideal generated by the elements $u_i - v_i u_1$ shows that $\mathcal{O}(\mathfrak{X}(1/p)_K) = K\langle u_1/p \rangle = K\langle b/p \rangle$. So $\mathfrak{X}(1/p)_K$ is isomorphic to the closed disc of radius $|p| = 1/p$ with local coordinate b ; it is well known that the completed local ring at $b = 0$ of such a disc is $K[[b]]$. The result follows since $\mathbb{1} \in \mathfrak{X}(1/p)_K$ implies that $\widehat{\mathcal{O}_{\mathfrak{X}_K, \mathbb{1}}} = \widehat{\mathcal{O}_{\mathfrak{X}(1/p)_K, \mathbb{1}}} = K[[b]]$. \square

Applying the functor \mathcal{O}° to the increasing chain of rigid K -varieties (9) and using Lemma 4.1.2 yields a decreasing chain of o_K -algebras

$$(10) \quad K[[b]] \supset \mathcal{O}^\circ(\mathfrak{X}(1/p)_K) \supset \dots \supset \mathcal{O}^\circ(\mathfrak{X}(r)_K) \\ \supset \mathcal{O}^\circ(\mathfrak{X}(r')_K) \supset \dots \supset \mathcal{O}^\circ(\mathfrak{X}_K) \supseteq o_K[[o_L]].$$

Definition 4.1.3. Let A be an o_K -subalgebra of $K[[b]]$ and let $m \geq 0$. The m -th *infinitesimal neighbourhood* of $\mathbb{1}$ in A is the image A_m of A in $K[[b]]/b^{m+1}K[[b]]$:

$$A_m := \frac{A + b^{m+1}K[[b]]}{b^{m+1}K[[b]]} \subset \frac{K[[b]]}{b^{m+1}K[[b]]}.$$

Remark 4.1.4. This construction respects inclusions and is compatible with variation in m . More precisely, whenever $A \subseteq B$ are two o_K -subalgebras of $K[[b]]$, for every $n \geq m$ there is a commutative diagram of o_K -algebras

$$\begin{array}{ccc} A_n & \longrightarrow & B_n \\ \downarrow & & \downarrow \\ A_m & \longrightarrow & B_m \end{array}$$

with injective horizontal arrows and surjective vertical arrows.

Definition 4.1.5. Let A be an o_K -subalgebra of $K[[b]]$ and for each $m \geq 0$, let $A_m^* := \text{Hom}_{o_K}(A_m, o_K)$. The *algebraic dual* of A is

$$A_\infty^* := \text{colim}_{m \geq 0} A_m^*.$$

Lemma 4.1.6. *Let $o_K[[o_L]] \subseteq A \subseteq B$ be two o_K -subalgebras of $K[[b]]$ and let $n \geq m \geq 0$.*

(1) *In the commutative square*

$$\begin{array}{ccc} A_n^* & \longleftarrow & B_n^* \\ \uparrow & & \uparrow \\ A_m^* & \longleftarrow & B_m^* \end{array}$$

all arrows are injective.

(2) The map $B_\infty^* \rightarrow A_\infty^*$ is injective.

Proof. (1) The vertical maps $A_m^* \rightarrow A_n^*$ are injective because $A_n \rightarrow A_m$ is surjective. Let C be the cokernel of the map $A_n \rightarrow B_n$. Since A_n contains $o_K[o_L]_n$ which is an o_K -lattice in $K[[b]]_n$, we see that C is a torsion o_K -module. The dual functor $(-)^*$ is left exact, so we have the exact sequence $0 \rightarrow C^* \rightarrow B_n^* \rightarrow A_n^*$. Since C is torsion, $C^* = 0$ which shows the injectivity of the horizontal arrows in our diagram.

(2) This follows by taking the colimit over all of the horizontal maps in part (1) above. \square

Thus we see that the connecting maps appearing in the colimit in Definition 4.1.5 are injective. Applying the contravariant algebraic dual functor $(-)_\infty^*$ to the chain (10) and using Lemma 4.1.6(2) gives us a chain of algebraic duals

$$\begin{aligned} \mathcal{O}^\circ(\mathfrak{X}(1/p)_K)_\infty^* &\subset \cdots \subset \mathcal{O}^\circ(\mathfrak{X}(r)_K)_\infty^* \\ &\subset \mathcal{O}^\circ(\mathfrak{X}(r')_K)_\infty^* \subset \cdots \subset \mathcal{O}^\circ(\mathfrak{X}_K)_\infty^* \subseteq o_K[o_L]_\infty^*. \end{aligned}$$

We can now calculate the largest one of these, namely the algebraic dual of the Iwasawa algebra $o_K[o_L]$, but first we must introduce integer-valued polynomials. Recall the following notion from [6].

Definition 4.1.7. A π -ordering for o_L is a subset $\{\alpha_0, \alpha_1, \alpha_2, \dots\}$ of o_L such that

$$(11) \quad v_\pi \left(\prod_{i=0}^{k-1} (\alpha_k - \alpha_i) \right) = \inf_{s \in o} v_\pi \left(\prod_{i=0}^{k-1} (s - \alpha_i) \right) \quad \text{for all } k \geq 1.$$

Starting from an arbitrary element $\alpha_0 \in o_L$, it is possible to construct a π -ordering $\{\alpha_0, \alpha_1, \dots\}$ of o_L by induction on k , choosing at each stage α_k to minimise the expression appearing on the right hand side of (11). In particular, π -orderings always exist, but are far from unique.

Definition 4.1.8. Let $\{\alpha_0, \alpha_1, \dots\}$ be a π -ordering for o_L .

(1) Define the *Lagrange polynomials* as follows: $f_0(X) := 1$ and

$$f_k(X) := \frac{(X - \alpha_0)(X - \alpha_1) \cdots (X - \alpha_{k-1})}{(\alpha_k - \alpha_0)(\alpha_k - \alpha_1) \cdots (\alpha_k - \alpha_{k-1})} \in L[X] \quad \text{for each } k \geq 1.$$

(2) Suppose that R is an o_L -algebra which embeds into $R_L := R \otimes_{o_L} L$. Then we define the ring of *R -valued polynomials on o_L* as follows:

$$\text{Int}(o_L, R) := \{g(X) \in R_L[X] : g(o_L) \subset R\}$$

(3) For each $m \geq 0$, let $\text{Int}(o_L, R)_m$ denote the R -submodule of $\text{Int}(o_L, R)$ consisting of all R -valued polynomials on o_L of degree at most m .

The following result, closely related to de Shalit's work on Mahler bases [30], explains why we are interested in these Lagrange polynomials.

Lemma 4.1.9. The set $\{f_0, f_1, f_2, \dots\}$ is an R -module basis for $\text{Int}(o_L, R)$.

Proof. It follows directly from Definition 4.1.7 that $v_\pi(f_k(s)) \geq 0$ for all $s \in o_L$ and all $k \geq 0$. Hence $f_k(o_L) \subset o_L \subset R$ for all $k \geq 0$ which implies that

$$(12) \quad Rf_0 + Rf_1 + Rf_2 + \cdots + Rf_n + \cdots \subseteq \text{Int}(o_L, R).$$

If $g \in R_L[X]$ has degree n and leading coefficient λ , then $g - \lambda(\alpha_n - \alpha_0) \cdots (\alpha_n - \alpha_{n-1})f_n$ has degree strictly less than n . This implies that $\{f_0, f_1, f_2, \dots\}$ generates $R_L[X]$ as an R_L -module. Now let $g \in \text{Int}(o_L, R)$ and write $g = \lambda_0 f_0 + \cdots + \lambda_n f_n$ for some $\lambda_0, \dots, \lambda_n \in R_L$ as above. Setting $X = \alpha_0$ shows that $\lambda_0 = g(\alpha_0) \in R$ since $g \in \text{Int}(o_L, R)$. Assume inductively that $\lambda_0, \dots, \lambda_{t-1} \in R$ for some $1 \leq t \leq n$. Setting $X = \alpha_t$ shows that

$$\lambda_t = g(\alpha_t) - \lambda_0 f_0(\alpha_t) - \lambda_1 f_1(\alpha_t) - \cdots - \lambda_{t-1} f_{t-1}(\alpha_t)$$

and this lies in R because $g(\alpha_t) \in R$ and $f_i(\alpha_t) \in R$ for all i . This completes the induction and shows that we have equality in (12). Taking $g = 0$ in the above argument also shows that the sum on the left hand side of (12) is direct. \square

Using Lemma 4.1.9, we obtain the following

Corollary 4.1.10.

(1) *The multiplication map*

$$\text{Int}(o_L, o_L) \otimes_{o_L} o_K \rightarrow \text{Int}(o_L, o_K)$$

is an isomorphism, which sends $\text{Int}(o_L, o_L)_m \otimes_{o_L} o_K$ onto $\text{Int}(o_L, o_K)_m$ for any $m \geq 0$.

(2) *The Lagrange polynomials $\{f_0(Y), \dots, f_m(Y)\}$ associated with a choice of π -ordering for o_L form an o_K -module basis for $\text{Int}(o_L, o_K)_m$.*

Proposition 4.1.11. *The evaluation map $\text{ev} : \text{Int}(o_L, o_K)_m \longrightarrow o_K[o_L]_m^*$ defined by*

$$\text{ev}(f(Y))(\lambda) := \lambda(f(Y))$$

for all $f(Y) \in \text{Int}(o_L, o_K)_m$, $\lambda \in o_K[o_L]_m^$ is an o_K -module isomorphism.*

Proof. This is essentially a complicated-looking tautology, but we try to give the details.

Note that $o_K[o_L]_m$ is an o_K -lattice in $K[b]_m$. We can therefore identify $o_K[o_L]_m^*$ with an o_K -submodule of $V := \text{Hom}_K(K[b]_m, K)$, a K -vector space of dimension $m+1$. The linear functionals $\text{ev}(1), \text{ev}(Y), \dots, \text{ev}(Y^m)$ are linearly independent in V because if $\sum_{i=0}^m c_i \text{ev}(Y^i) = 0$ then $\text{ev}(\sum_{i=0}^m c_i Y^i)(\delta_a) = \sum_{i=0}^m c_i a^i = 0$ for all $a \in o_L$ and this forces $c_0 = \dots = c_m = 0$. It follows that $\text{ev} : K[Y]_m \rightarrow V$ is injective and is therefore an isomorphism by the rank-nullity theorem.

Hence $\text{ev} : \text{Int}(o_L, o_K)_m \rightarrow o_K[o_L]_m^*$ is injective. However if $g \in o_K[o_L]_m^*$ then by the above we can find some $f(Y) \in K[Y]_m$ such that $\text{ev}(f(Y)) = g$. Since $\delta_a \in o_K[o_L]$ for all $a \in o_L$, we see that $f(a) = \text{ev}(f(Y))(\delta_a) = g(\delta_a)$ must lie in o_K for all $a \in o_L$. \square

Corollary 4.1.12. *The map $\text{ev} : \text{Int}(o_L, o_K) \rightarrow o_K[o_L]_\infty^*$ is an isomorphism.*

Proof. This follows immediately from Proposition 4.1.11. \square

Proposition 4.1.13. *Suppose that K is discretely valued. Then*

$$\mathcal{O}^\circ(\mathfrak{X}_K)_\infty^* = \operatorname{colim}_{r < 1} \mathcal{O}^\circ(\mathfrak{X}(r)_K)_\infty^*.$$

Proof. Since colimits commute with colimits, it is enough to show that for every $m \geq 0$,

$$\mathcal{O}^\circ(\mathfrak{X}_K)_m^* = \operatorname{colim}_{r < 1} \mathcal{O}^\circ(\mathfrak{X}(r)_K)_m^*.$$

Fix $m \geq 0$. Then $\mathcal{O}^\circ(\mathfrak{X}(r)_K)_m$ form a decreasing chain of o_K -submodules of the $m + 1$ -dimensional K -vector space $K[[b]]_m$, and all of them contain the o_K -lattice $o_K[[o_L]]_m$. Since K is discretely valued, the o_K -module $(K/o_K)^{m+1}$ satisfies the descending chain condition. Hence there exists $r_0 < 1$ such that

$$(13) \quad \mathcal{O}^\circ(\mathfrak{X}(r)_K)_m = \mathcal{O}^\circ(\mathfrak{X}(r_0)_K)_m \quad \text{whenever} \quad r_0 \leq r < 1.$$

Following an argument of Schmidt [26, proof of Proposition 4.9], we will now show that

$$\mathcal{O}^\circ(\mathfrak{X}_K)_m = \mathcal{O}^\circ(\mathfrak{X}(r_0)_K)_m.$$

The forward inclusion is clear, so fix some $\xi \in \mathcal{O}^\circ(\mathfrak{X}(r_0)_K)_m$, choose a sequence of real numbers $r_0 < r_1 < r_2 < \dots$ approaching 1 and consider the K -Banach space

$$A_j := \mathcal{O}(\mathfrak{X}(r_j)_K).$$

Let $\varphi_j : A_j^\circ \rightarrow K[[b]]_m$ be the obvious o_K -linear map. Using (13) we see that the convex subset

$$\varphi_j^{-1}(\xi) \subset A_j$$

is non-empty. It was recorded in the proof of [29, Lemma 6.1] that the restriction maps $A_{j+1} \rightarrow A_j$ are compact. We may therefore argue as in [16, Proposition V.3.2] that

$$\bigcap_{j=0}^{\infty} \varphi_j^{-1}(\xi) \subseteq \mathcal{O}^\circ(\mathfrak{X}_K)$$

is non-empty. Then any element λ in this intersection satisfies $\lambda_m = \xi$, so $\xi \in \mathcal{O}^\circ(\mathfrak{X}_K)_m$ as required. Hence $\mathcal{O}^\circ(\mathfrak{X}_K)_m^* = \mathcal{O}^\circ(\mathfrak{X}(r)_K)_m^*$ whenever $r_0 \leq r < 1$, and the result follows. \square

4.2. The matrix coefficients $\rho_{i,j}(Y)$. Let $\mathbf{B}_{\mathbb{C}_p}$ be the rigid analytic open unit disc of radius 1 defined over \mathbb{C}_p , with global coordinate function Z . There is a twisted $G_L = \operatorname{Gal}(\mathbb{C}_p/L)$ -action on $\mathcal{O}(\mathbf{B}_{\mathbb{C}_p})$ given by $F \mapsto F^\sigma \circ [\tau(\sigma^{-1})]$, which induces an L -algebra isomorphism

$$\mu : \mathcal{O}(\mathfrak{X}_L) \xrightarrow{\cong} \mathcal{O}(\mathbf{B}_{\mathbb{C}_p})^{G_L,*},$$

see [28, Corollary 3.8]. Inspecting the proof of this result, we see that it extends naturally to give a description of $\mathcal{O}(\mathfrak{X}_K)$ for more general closed coefficient fields $L \subseteq K \subseteq \mathbb{C}_p$ as well:

Lemma 4.2.1. *There is a K -algebra isomorphism*

$$\mu_K : \mathcal{O}(\mathfrak{X}_K) \xrightarrow{\cong} \mathcal{O}(\mathbf{B}_{\mathbb{C}_p})^{G_K,*}.$$

Since $\mathcal{O}^\circ(\mathbf{B}_{\mathbb{C}_p}) = o_{\mathbb{C}_p}[[Z]]$, we deduce the following

Corollary 4.2.2. *There is an isomorphism of o_K -algebras*

$$\mu_K : \mathcal{O}^\circ(\mathfrak{X}_K) \xrightarrow{\cong} o_{\mathbb{C}_p}[[Z]]^{G_K,*}.$$

Until the end of §4.2, we assume that Ω is transcendental over K .

Definition 4.2.3. We call an o_K -subalgebra R of $K[\Omega] \cap o_{\mathbb{C}_p}$ *admissible* if $P_n(\Omega) \in R$ for all $n \geq 0$, and if R is stable under the natural G_L -action on $K[\Omega] \cap o_{\mathbb{C}_p}$ (in particular, K itself is then stable under G_L).

Example 4.2.4. The algebra $K[\Omega] \cap o_{\mathbb{C}_p}$ is itself an admissible o_K -subalgebra of $K[\Omega]$.

Proof. This follows from Corollary 4.2.2 together with [28, Lemma 4.2(5)]. \square

Definition 4.2.5. Let $R \subset K[\Omega]$ be an admissible o_K -subalgebra.

- (1) Let $K[\Omega]_n := \{f(\Omega) \in K[\Omega] : \deg(f) \leq n\}$ for each $n \geq 0$.
- (2) Let $R_n := R \cap K[\Omega]_n$ for each $n \geq 0$.
- (3) $\{b_n(\Omega) : n \geq 0\} \subset R$ is a *regular basis* if

$$b_0(\Omega) = 1, \quad \text{and} \quad R_n = R_{n-1} \oplus o_K b_n(\Omega) \quad \text{for all } n \geq 1.$$

Lemma 4.2.6. *Suppose that K is discretely valued. Then a regular basis exists for every admissible o_K -subalgebra R of $K[\Omega] \cap o_{\mathbb{C}_p}$.*

Proof. Since Ω is assumed to be transcendental over K , the K -vector space $K[\Omega]_n$ has dimension $n+1$. The restriction of the norm $|\cdot|$ on \mathbb{C}_p to $K[\Omega]_n$ turns it into a normed vector space over K and by Definition 4.2.3(1), R_n is contained in the unit ball with respect to this norm. Since any two norms on a finite dimensional K -vector space are equivalent — see [27, Proposition 4.13] — it follows that $R_n \subseteq \pi^{-m} o_K[\Omega]_n$ for sufficiently large m .

Since K is discretely valued, its valuation ring o_K is Noetherian and this forces R_n to be a free o_K -module of rank $n+1$. Because the R_n 's form a nested chain, we can now construct the desired o_K -module basis for R by induction on n . \square

Example 4.2.7. Because Ω is assumed to be transcendental over K , Lemma 3.2.2(1) together with Lemma 3.2.3 implies that $\sum_{n=0}^{\infty} o_K P_n(\Omega)$ is isomorphic to $U(\mathcal{G}) \otimes_{o_L} o_K$ as an o_K -algebra. Abusing notation, we will write $U := \sum_{n=0}^{\infty} o_K P_n(\Omega)$ until the end of §4. Although this conflicts with Definition 3.2.1(2), we hope that no confusion will be caused by this abuse of notation. Then U is an admissible subalgebra of $K[\Omega]$, and $\{P_n(\Omega) : n \geq 0\}$ is a regular basis for R : since $\deg P_j(Y) = j$, an element $f(\Omega)$ of U_n is a K -linear combination of $P_0(\Omega), \dots, P_n(\Omega)$ lying in U , but $\{P_m(\Omega) : m \geq 0\}$ is an o_L -module basis for U so all coefficients of $f(\Omega)$ must in fact lie in o_L .

Until the end of §4.2, we assume that

- K is a *discretely* valued intermediate subfield $L \subseteq K \subseteq \mathbb{C}_p$,
- Ω is transcendental over K ,
- $R \subseteq K[\Omega] \cap o_{\mathbb{C}_p}$ is an admissible o_K -subalgebra, and
- $\{b_n(\Omega) : n \geq 0\}$ is a regular basis for R .

Lemma 4.2.8. *Take $j \geq 0$.*

(1) *There are unique $\rho_{0,j}(Y), \rho_{1,j}(Y), \dots, \rho_{j,j}(Y) \in K[Y]$ such that*

$$P_j(Y\Omega) = \sum_{i=0}^j \rho_{i,j}(Y)b_i(\Omega).$$

(2) *$\deg \rho_{i,j}(Y) \leq j$ whenever $0 \leq i \leq j$.*

(3) *$\deg \rho_{j,j}(Y) = j$.*

(4) *$\rho_{i,j}(a) \in o_K$ whenever $a \in o_L$ and $0 \leq i \leq j$.*

Proof. (1) Ω is transcendental over K , and $\{b_i(\Omega) : i \geq 0\}$ is a K -vector space basis for $K[\Omega]$ with $\deg b_i(\Omega) = i$ for each i . Hence it is also a $K[Y]$ -module basis for the two-variable polynomial algebra $K[\Omega, Y]$, so we can find unique $\rho_{i,j}(Y) \in K[Y]$ such that

$$P_j(Y\Omega) = \sum_{i \geq 0} \rho_{i,j}(Y)b_i(\Omega)$$

where $\rho_{i,j}(Y) = 0$ for sufficiently large i . Now $P_j(s)$ is a polynomial in s of degree j by [28, Lemma 4.2(3)], so Ω^j is the highest degree monomial in Ω appearing in $P_j(Y\Omega)$. Since $\deg b_i(\Omega) = i$, this means $\rho_{i,j}(Y) = 0$ for $i > j$.

(2) Since the highest degree monomial in Y appearing in $P_j(Y\Omega)$ is Y^j , this means that $\deg \rho_{i,j}(Y) \leq j$ for each $i \leq j$.

(3) The monomial $Y^j \Omega^j$ appears in $P_j(Y\Omega)$ with a non-zero coefficient. This monomial does not appear in $\rho_{i,j}(Y)b_i(\Omega)$ for any $i < j$ because $\deg b_i(\Omega) = i$ for all i . So it must appear in $\rho_{j,j}(Y)b_j(\Omega)$, and because of (2), this can only happen if $\deg \rho_{j,j}(Y) = j$.

(4) Let $a \in o_L$. We know that $P_j(a\Omega) \in o_{\mathbb{C}_p}$ by [28, Lemma 4.2(5)]; in fact, $P_j(a\Omega)$ is an o_L -linear combination of the $P_i(\Omega)$ for $0 \leq i \leq j$ by Corollary 3.2.6, so $P_j(a\Omega) \in R$. Setting $Y = a$ in (1) shows that $\rho_{i,j}(a) \in o_K$, since $\{b_i(\Omega) : i \geq 0\}$ is a regular basis for R . \square

Theorem 4.2.9. *For each $\lambda \in D^{L-\text{an}}(o_L, K)$ we have*

$$\mu_K(\lambda) = \sum_{j=0}^{\infty} \sum_{k=0}^j \lambda(\rho_{k,j}(Y))b_k(\Omega)Z^j.$$

In the case when $\lambda = \delta_a$ for some $a \in o_L$, Lemma 4.2.8 implies that

$$\begin{aligned} \mu(\delta_a) &= \sum_{j=0}^{\infty} P_j(a\Omega)Z^j = \sum_{j=0}^{\infty} \left(\sum_{i=0}^j \rho_{i,j}(a)b_i(\Omega) \right) Z^j \\ &= \sum_{j=0}^{\infty} \sum_{k=0}^j \delta_a(\rho_{k,j}(Y))b_k(\Omega)Z^j \end{aligned}$$

which explains where the formula comes from. We will now give a rigorous argument to show that the formula is valid for any $\lambda \in D^{L-\text{an}}(o_L, K)$.

Lemma 4.2.10. *Let $t := \log_{LT}(Z)$ be the Lubin-Tate logarithm. Then*

$$\mu_K(\lambda) = \sum_{k=0}^{\infty} \lambda(Y^k/k!) \Omega^k t^k \quad \text{for all } \lambda \in D^{L-\text{an}}(o_L, K).$$

Proof. We may identify $\mathbb{C}_p[[t]]$ with $\mathbb{C}_p[[Z]]$, and we write $\mu_K(\lambda) = \sum_{m=0}^{\infty} c_{i,m} t^m$ for some $c_{i,m} \in \mathbb{C}_p$. Then applying [28, Lemma 4.6(8)], we have

$$\begin{aligned} \lambda(Y^k/k!) &= \{\mu_K(\lambda), Y^k/k!\} = \frac{(\Omega^{-1}\partial_t)^k}{k!}(\mu_K(\lambda))(0) \\ &= \Omega^{-k} c_{i,k} \quad \text{for all } k \geq 0. \quad \square \end{aligned}$$

Proposition 4.2.11. *Let $\lambda \in \text{Hom}_L(L[Y], K)$. Then in $\mathbb{C}_p[[t]] = \mathbb{C}_p[[Z]]$,*

$$\sum_{k=0}^{\infty} \lambda(Y^k/k!) \Omega^k t^k = \sum_{j=0}^{\infty} \sum_{k=0}^j \lambda(\rho_{k,j}(Y)) b_k(\Omega) Z^j.$$

Proof. For each $k \geq 0$, write $t^k = \sum_{j=k}^{\infty} d_j^{(k)} Z^j \in L[[Z]]$. Substituting this into Lemma 4.2.10 gives

$$\begin{aligned} (14) \quad \sum_{k=0}^{\infty} \lambda(Y^k/k!) \Omega^k t^k &= \sum_{k=0}^{\infty} \lambda(Y^k/k!) \Omega^k \sum_{j=k}^{\infty} d_j^{(k)} Z^j \\ &= \sum_{j=0}^{\infty} \left(\sum_{k=0}^j \frac{1}{k!} d_j^{(k)} \Omega^k \lambda(Y^k) \right) Z^j. \end{aligned}$$

On the other hand, the identity

$$\sum_{j=0}^{\infty} P_j(Y\Omega)Z^j = \exp(Y\Omega t) = \sum_{k=0}^{\infty} \frac{1}{k!} t^k \Omega^k Y^k = \sum_{k=0}^{\infty} \frac{Y^k \Omega^k}{k!} \sum_{j=k}^{\infty} d_j^{(k)} Z^j$$

together with Lemma 4.2.8 shows that for all $j \geq 0$ we have

$$(15) \quad \sum_{k=0}^j \frac{1}{k!} d_j^{(k)} \Omega^k Y^k = P_j(\Omega Y) = \sum_{k=0}^j \rho_{k,j}(Y) b_k(\Omega).$$

Now, the L -linear form $\lambda : L[Y] \rightarrow K$ extends to a $K[\Omega]$ -linear form $K[\Omega, Y] \rightarrow K[\Omega]$. Applying this extension to (15) gives

$$\sum_{k=0}^j \frac{1}{k!} d_j^{(k)} \Omega^k \lambda(Y^k) = \sum_{k=0}^j \lambda(\rho_{k,j}(Y)) b_k(\Omega).$$

Substituting this equation into (14) gives the result. \square

Proof of Theorem 4.2.9. Follows immediately from Lemma 4.2.10 and Proposition 4.2.11. \square

Definition 4.2.12. Let \check{R} be the o_K -linear span of $\{\rho_{k,j}(Y) : j \geq k \geq 0\}$ in the space $I := \text{Int}(o_L, o_K)$ of o_K -valued polynomials on o_L .

We will see shortly that \check{R} does not depend on the choice of regular basis for R .

Corollary 4.2.13. Let $\lambda \in D^{L-\text{an}}(o_L, K)$. Then $\mu_K(\lambda) \in R[[Z]]$ if and only if $\lambda(\check{R}) \subseteq o_K$.

Proof. Theorem 4.2.9 tells us that $\mu_K(\lambda)$ belongs to $R[[Z]]$ if and only if $\sum_{k=0}^j \lambda(\rho_{k,j}(Y)) b_k(\Omega) \in R$ for all $j \geq 0$. Since $\{b_k(\Omega) : k \geq 0\}$ is a regular basis, this is equivalent to $\lambda(\rho_{k,j}(Y)) \in o_K$ for all $j \geq k \geq 0$. \square

Proposition 4.2.14. Let $\lambda \in \text{Hom}_K(K[Y], K)$ be such that $\lambda(\check{R}) \subseteq o_K$. Then there exists $\tilde{\lambda} \in \mu_K^{-1}(R[[Z]]) \subseteq \mathcal{O}^\circ(\mathfrak{X}_K)$ such that $\tilde{\lambda}|_{K[Y]} = \lambda$.

Proof. The twisted G_L -action on $\mathbb{C}_p[[Z]]$ preserves $R[[Z]]$ since we assumed that $R \subseteq K[\Omega] \cap o_{\mathbb{C}_p}$ is G_L -stable in Definition 4.2.3. Therefore $R[[Z]]^{G_L,*}$ makes sense.

Define $F_\lambda := \sum_{j=0}^{\infty} \sum_{k=0}^j \lambda(\rho_{k,j}(Y)) b_k(\Omega) Z^j \in \mathbb{C}_p[[Z]]$. Then $F_\lambda \in K[[\Omega t]] = \mathbb{C}_p[[Z]]^{G_K,*}$ by Proposition 4.2.11 and $F_\lambda \in R[[Z]]$ because $\lambda(\check{R}) \subseteq o_K$. Hence $F_\lambda \in R[[Z]]^{G_K,*} \subseteq o_{\mathbb{C}_p}[[Z]]^{G_K,*}$, so $F_\lambda = \mu_K(\tilde{\lambda})$ for some $\tilde{\lambda} \in \mathcal{O}^\circ(\mathfrak{X}_K)$ by Corollary 4.2.2. In particular, $\tilde{\lambda} \in \mu_K^{-1}(R[[Z]])$.

Next, applying [28, Lemma 4.6(8)] we see that for all $m \geq 0$,

$$\begin{aligned} \tilde{\lambda}(Y^m/m!) &= \{\mu_K(\tilde{\lambda}), Y^m/m!\} = \{F_\lambda, Y^m/m!\} \\ &= \left\{ \sum_{k=0}^{\infty} \lambda(Y^k/k!) \Omega^k t^k, Y^m/m! \right\} = \lambda(Y^m/m!). \end{aligned}$$

Since the $Y^m/m!$ span $K[Y]$ as a K -vector space, we have $\tilde{\lambda}|_{K[Y]} = \lambda$. \square

Recall the isomorphism $\text{ev} : \text{Int}(o_L, o_K) \rightarrow o_K[[o_L]]_\infty^*$ from Corollary 4.1.12.

Theorem 4.2.15. We have $\text{ev}(\check{R}) = \mu_K^{-1}(R[[Z]])_\infty^*$.

Proof. The o_K -module \check{R} contains the o_K -submodule of $K[Y]$ generated by $\{\rho_{j,j}(Y) : j \geq 0\}$ and $\deg \rho_{j,j}(Y) = j$ for each $j \geq 0$ by Lemma 4.2.8(3). Hence \check{R} spans $K[Y]$ as a K -vector space. On the other hand, $\check{R}_n := \check{R} \cap K[Y]_{\leq n}$ is contained in $\text{Int}(o_L, o_K)_n$ by Lemma 4.2.8(4), which is a finitely generated o_K -module by Remark 4.1.10(2). Since K is discretely valued, \check{R}_n is a finitely generated o_K -module for each $n \geq 0$. So we can find an o_K -module basis $\{t_0, t_1, \dots, t_n, \dots\}$ for \check{R} such that $\{t_0, \dots, t_n\}$ is an o_K -module basis for \check{R}_n for each $n \geq 0$. It follows that the natural map $\check{R} \otimes_{o_K} K \rightarrow K[Y]$ is an isomorphism, and we may identify $\text{Hom}_{o_K}(\check{R}, o_K)$ with $\{\phi \in \text{Hom}_K(\check{R}, K) : \phi(\check{R}) \subseteq o_K\}$.

Let $\{t_m^* : m \geq 0\} \subset \text{Hom}_{o_K}(\check{R}, o_K)$ be determined by

$$t_m^*(t_n) = \delta_{m,n} \quad \text{for all } m, n \geq 0.$$

Then by Proposition 4.2.14, t_m^* extends to some $\lambda_m \in \mu_K^{-1}(R[[Z]])$ such that $\lambda_m|_{K[Y]} = t_m^*$. In particular, we have $\lambda_m(t_n) = \delta_{m,n}$ for all $m, n \geq 0$.

Now suppose that $g \in \mu_K^{-1}(R[[Z]])_\infty^* \subseteq o_K[[o_L]]_\infty^*$. Then $g = \text{ev}(h)$ for some $h \in \text{Int}(o_L, o_K)_m$ by Proposition 4.1.11. Since $h \in K[Y]_{\leq m}$ and since $\{t_0, \dots, t_m\}$ is a K -vector space basis for $K[Y]_m$, we can write $h = \sum_{n=0}^m c_n t_n$ for some $c_n \in K$. But then

$$g(\lambda_n) = \text{ev}(h)(\lambda_n) = \lambda_n(h) = t_n^*(h) = c_n \quad \text{for all } n \geq 0.$$

Since $\lambda_n \in \mu_K^{-1}(R[[Z]])$ and $g \in \mu_K^{-1}(R[[Z]])_\infty^*$, we conclude that $g(\lambda_n) \in o_K$ for all $n \geq 0$. Hence $h \in \sum_{n=0}^m o_K t_n \subseteq \check{R}$ and $g = \text{ev}(h) \in \text{ev}(\check{R})$. Hence $\mu_K^{-1}(R[[Z]])_\infty^* \subseteq \text{ev}(\check{R})$.

Conversely, take $\lambda \in \mu_K^{-1}(R[[Z]])$. Then $\lambda(\check{R}) \subseteq o_K$ by Corollary 4.2.13 and thus for all $g \in \check{R}$, $\text{ev}(g)(\lambda) = \lambda(g) \in o_K$. Hence $\text{ev}(\check{R}) \subseteq \mu_K^{-1}(R[[Z]])_\infty^*$. \square

Corollary 4.2.16. *Let $S \subseteq R$ be two admissible subalgebras of $K[\Omega]$. Then $\check{R} \subseteq \check{S}$.*

Proof. We have $\mu_K^{-1}(S[[Z]]) \subseteq \mu_K^{-1}(R[[Z]])$, so $\mu_K^{-1}(R[[Z]])_\infty^* \subseteq \mu_K^{-1}(S[[Z]])_\infty^*$ by Lemma 4.1.6(2). Hence $\text{ev}(\check{R}) \subseteq \text{ev}(\check{S})$ by Theorem 4.2.15. Hence $\check{R} \subseteq \check{S}$ because ev is an isomorphism by Corollary 4.1.12. \square

Note that Theorem 4.2.15 implies that the o_K -module \check{R} depends only on the admissible subalgebra R and not the particular choice of regular basis $\{b_n(\Omega) : n \geq 0\}$ for R .

Lemma 4.2.17. *If $\lambda \in D^{L-\text{an}}(o_L, K)$, then $\lambda \in o_K[[o_L]]$ if and only if we have $\lambda(\text{Int}(o_L, o_K)) \subseteq o_K$.*

Proof. Suppose that $\lambda(\text{Int}(o_L, o_K)) \subseteq o_K$. The π -adic completion of I is naturally isomorphic to the ring $\mathcal{C}^0(o_L, o_K)$ of o_K -valued continuous functions on o_L . Since $\lambda(I) \subseteq o_K$, λ extends to an o_K -linear form $\tilde{\lambda} : \mathcal{C}^0(o_L, o_K) \rightarrow o_K$ which is automatically continuous. View $\tilde{\lambda}$ as an element of $o_K[[o_L]] = D^{\text{cts}}(o_L, K)$. The restrictions of $\tilde{\lambda}$ and of $\lambda \in D^{L-\text{an}}(o_L, K)$ to $K[Y]$ agree by

construction. Since $K[Y]$ is dense in $C^{\text{an}}(o_L, K)$, we conclude that λ lies in $o_K[[o_L]]$.

Conversely, if $\lambda \in o_K[[o_L]] = \mathcal{C}^0(o_L, o_K)^*$, then λ must take integer values on $\text{Int}(o_L, o_K) \subset \mathcal{C}^0(o_L, o_K)$. \square

Theorem 4.2.18. *Let R be an admissible subalgebra of $K[\Omega]$.*

We have $\mu_K^{-1}(R[[Z]]) = o_K[[o_L]]$ if and only if $\tilde{R} = I$.

Proof. (\Leftarrow). Suppose that $\tilde{R} = I$, and take $\lambda \in \mu_K^{-1}(R[[Z]])$. Then $\lambda(\tilde{R}) \subseteq o_K$ by Corollary 4.2.13. Since $\tilde{R} = I$, this means that $\lambda(I) \subseteq o_K$. Hence $\lambda \in o_K[[o_L]]$ by Lemma 4.2.17.

(\Rightarrow). Suppose that $\tilde{R} < I$. Since K is discretely valued, K/o_K is an injective cogenerator of the category of o_K -modules. Hence $\text{Hom}_{o_K}(I/\tilde{R}, K/o_K)$ is non-zero. So there exists an o_K -linear map $\lambda : I \rightarrow K$ such that $\lambda(\tilde{R}) \subseteq o_K$, but $\lambda(I) \not\subseteq o_K$. Regard λ as an element of $\text{Hom}_K(K[Y], K)$; then by Proposition 4.2.14, λ extends to some $\tilde{\lambda} \in \mathcal{O}^\circ(\mathfrak{X}_K)$ such that $\tilde{\lambda}|_{K[Y]} = \lambda$. Since $\lambda(\tilde{R}) \subseteq o_K$, using Theorem 4.2.9 we see that $\mu_K(\tilde{\lambda}) \in R[[Z]]$. However, $\tilde{\lambda} \notin o_K[[o_L]]$ by Lemma 4.2.17 because $\tilde{\lambda}(I) \not\subseteq o_K$, so $\tilde{\lambda} \in \mu_K^{-1}(R[[Z]]) \setminus o_K[[o_L]]$. \square

We will now see what implications the above general results have for particular choices of the admissible subalgebra R . Let $B = K[\Omega] \cap o_{\mathbb{C}_p}$ be the largest possible admissible subalgebra of $K[\Omega]$, and let $U := \sum_{n=0}^{\infty} o_K P_n(\Omega)$ be the smallest possible one. Recall from Example 4.2.7 that $\{P_n(\Omega) : n \geq 0\}$ forms a regular basis for U .

Corollary 4.2.19.

- (1) $\tilde{U} = \text{Int}(o_L, o_K)$ if and only if $\mu_K^{-1}(U[[Z]]) = o_K[[o_L]]$.
- (2) $o_K[[o_L]] = \Lambda_K(\mathfrak{X})$ if and only if $\tilde{B} = \text{Int}(o_L, o_K)$.

Proof. (1) This is an immediate consequence of Theorem 4.2.18 with $R = U$.

(2) Theorem 4.2.18 tells us that $\tilde{B} = I$ if and only if $o_K[[o_L]] = \mu_K^{-1}(B[[Z]])$. However $\mu_K^{-1}(B[[Z]]) = \mu_K^{-1}(\mathbb{C}_p[[Z]]^{G_L,*} \cap B[[Z]])$ since $\mu_K(\mathcal{O}(\mathfrak{X})_K)$ is fixed by the twisted G_L -action on $\mathbb{C}_p[[Z]]$ by Lemma 4.2.1. Hence $\mu_K^{-1}(B[[Z]]) = \mu_K^{-1}(o_{\mathbb{C}_p}[[Z]]^{G_L,*}) = \Lambda_K(\mathfrak{X})$ by Corollary 4.2.2, and the result follows. \square

Recall the matrix coefficients $\sigma_{i,j}(a)$ from Corollary 3.2.6.

Lemma 4.2.20. *Let $R = U$ and let $b_n := P_n$ for each $n \geq 0$. Then*

- (1) $\rho_{ij}(Y) = \sigma_{i,j}(Y)$ for all $j \geq i \geq 0$, and
- (2) $[a](Z)^i = \sum_{j=i}^{\infty} \sigma_{i,j}(a) Z^j$ for any $a \in o_L, i \geq 0$.

Proof. (1) This follows by comparing Corollary 3.2.6 with Lemma 4.2.8(1).

(2) Using Definition 3.2.1(5) and Lemma 3.2.3 we see that $\langle P_k(s), Z^i \rangle = \delta_{ki}$ for all $i, k \geq 0$. By Corollary 3.2.6 we have $P_j(as) = \sum_{k=0}^j \sigma_{kj}(a) P_k(s)$. Fix

$i \geq 0$ and apply $\langle -, Z^i \rangle$ to this equation: using equation (4) we then have

$$\sigma_{i,j}(a) = \left\langle \sum_{k=0}^j \sigma_{kj}(a) P_k(s), Z^i \right\rangle = \langle P_j(as), Z^i \rangle = \langle P_j(s), [a](Z)^i \rangle.$$

Hence $\sigma_{i,j}(a)$ is precisely the coefficient of Z^j in the power series $[a](Z)^i$. \square

This justifies the definition of the polynomials $\sigma_{i,j}(Y)$ which was given in §1.5. We can now give the proof of Theorem 1.5.1 from the Introduction.

Theorem 4.2.21. *If $\Lambda_L(\mathfrak{X}) = o_L[o_L]$, then $\text{Pol} = \text{Int}$.*

Proof. Note that $\text{Pol} = \tilde{U}$, in view of Lemma 4.2.20(1) and Definition 4.2.12. Now $\Lambda_L(\mathfrak{X}) = \mathcal{O}^\circ(\mathfrak{X}_L)$, so if this is equal to $o_L[o_L]$, then $\tilde{B} = \text{Int}(o_L, o_L)$ by Corollary 4.2.19(2). But $U \subseteq B$, so $\tilde{B} \subseteq \tilde{U} \subseteq \text{Int}(o_L, o_L)$ by Corollary 4.2.16. Hence $\tilde{U} = \text{Int}(o_L, o_L)$ as claimed. \square

4.3. Calculating the matrix coefficients $\sigma_{i,j}(Y)$. Here we will assume that the coordinate Z on the Lubin-Tate formal group is chosen in such a way that

$$\log_{LT}(Z) = \sum_{n=0}^{\infty} \frac{Z^{q^n}}{\pi^n}.$$

It turns out that the polynomials $P_j(s)$ are *sparse*: the coefficient of s^i in $P_j(s)$ is non-zero *only* if $i \equiv j \pmod{q-1}$. We will obtain more information about these coefficients; this will require developing some notation to deal with this sparsity. The calculations that follow rest on the following observation.

Proposition 4.3.1. *For every $n \geq 0$, we have*

$$P_n(Y) = \sum_{k_0+qk_1+\dots+q^d k_d=n} \frac{Y^{k_0+\dots+k_d}}{k_0! \dots k_d! \cdot \pi^{1 \cdot k_1 + 2 \cdot k_2 + \dots + d \cdot k_d}}.$$

Proof. If $\log_{LT}(Z) = \sum_{k=0}^{\infty} Z^{q^k} / \pi^k$ and \exp is the usual exponential, then

$$\begin{aligned} \sum_{n=0}^{\infty} P_n(Y) Z^n &= \exp(Y \cdot \log_{LT}(Z)) = \prod_{\ell \geq 0} \exp(Y \cdot Z^{q^\ell} / \pi^\ell) \\ &= \prod_{\ell \geq 0} \sum_{k \geq 0} (Y \cdot Z^{q^\ell} / \pi^\ell)^k / k! \end{aligned}$$

The coefficient of Z^n in this product is the sum of $Y^{k_0+\dots+k_d} / k_0! \dots k_d! \cdot \pi^{1 \cdot k_1 + 2 \cdot k_2 + \dots + d \cdot k_d}$ over all tuples (k_0, \dots, k_d) of positive integers such that $k_0 + qk_1 + \dots + q^d k_d = n$. \square

The following formula for the derivative $\frac{d}{dY} P_n(Y)$ will be very useful in the calculations.

Proposition 4.3.2. *For every $n \geq 0$, we have $\frac{d}{dY} P_n(Y) = \sum_{k \geq 0} \pi^{-k} \cdot P_{n-q^k}(Y)$.*

Proof. We have $P_n(Y + Z) = P_n(Y) + \sum_{j=1}^n P_j(Z)P_{n-j}(Y)$ by [28, Lemma 4.2(4)]. Hence it is enough to determine which $P_j(Z)$ have a term of degree 1 in them, and what the corresponding coefficient is in this case. The answer now follows from Proposition 4.3.1. \square

We fix $m \in \{0, 1, 2, \dots, q-2\}$ from now on. We will use the convenient notation

$$\underline{i} := m + i(q-1) \quad \text{for all } i \geq 0.$$

Definition 4.3.3. For each $j \geq i \geq 0$, we define

$$Q_m(i, j) := \left\{ \mathbf{k} \in \mathbb{N}^\infty : \sum_{\ell=0}^{\infty} k_\ell = \underline{i}, \quad \sum_{\ell=1}^{\infty} k_\ell \left(\frac{q^\ell - 1}{q - 1} \right) = j - i \right\}, \quad \text{and}$$

$$r_{i,j}^{(m)} := \sum_{\mathbf{k} \in Q_m(i,j)} \binom{\underline{i}}{k_0; k_1; k_2; \dots} \cdot \pi^{-\sum_{\ell=1}^{\infty} \ell \cdot k_\ell}.$$

Here $\binom{\underline{i}}{k_0; k_1; k_2; \dots} = \frac{(\underline{i})!}{k_0! \cdot k_1! \cdot k_2! \dots}$ is the multinomial coefficient.

Lemma 4.3.4. We have $r_{jj}^{(m)} = 1$ for all $j \geq 0$.

Proof. If $i = j$, then the second condition on a vector $\mathbf{k} \in \mathbb{N}^\infty$ to lie in $Q_m(i, j)$ forces $k_1 = k_2 = \dots = 0$ because $\frac{q^\ell - 1}{q - 1} > 0$ for all $\ell \geq 1$. But then $k_0 = \underline{j}$ from the first condition, so the formula for $r_{jj}^{(m)}$ collapses to give 1. \square

Proposition 4.3.5. Let $n = \underline{j}$ for some $j \geq 0$. Write

$$P_n(s) = \sum_{k=0}^n b_k^{(n)} s^k$$

with $b_k^{(n)} \in L$ for $k = 0, \dots, n$.

- (1) We have $b_k^{(n)} = 0$ if $k \not\equiv n \pmod{q-1}$.
- (2) For each $0 \leq i \leq j$, we have $b_{\underline{i}}^{(j)} = \frac{r_{i,j}^{(m)}}{\underline{i}!}$.

Proof. By Proposition 4.3.1, the coefficient $b_k^{(n)}$ of s^k in $P_n(s)$, is given by

$$b_k^{(n)} = \sum_{\mathbf{k}} \frac{1}{(k_0! k_1! k_2! \dots) \pi^{0 \cdot k_0 + 1 \cdot k_1 + 2 \cdot k_2 + \dots}},$$

where the sum runs over all possible sequences $\mathbf{k} = (k_0, k_1, k_2, \dots)$ of non-negative integers satisfying the following two conditions:

$$k_0 + k_1 + k_2 + \dots = k, \quad \text{and} \quad k_0 + qk_1 + q^2k_2 + \dots = n.$$

Of course given any such sequence, necessarily k_ℓ must be zero for all sufficiently large ℓ depending only on n and k , and the set of solutions to these equations is always finite, so the sum of all these fractions makes sense.

Next note that if k_0, k_1, \dots satisfies these two conditions, then necessarily

$$n \equiv k \pmod{q-1}.$$

This implies part (1). For part (2), let $k = \underline{i}$ and $n = \underline{j}$, and suppose that the non-negative integers k_0, k_1, \dots satisfy $k_0 + k_1 + \dots = k$; then subtracting gives

$$k_0 + qk_1 + q^2k_2 + \dots = m + (q-1)j \Leftrightarrow (q-1)k_1 + (q^2-1)k_2 + \dots = (q-1)(j-i).$$

In this way, we see that $Q_m(i, j)$ is precisely the set of sequences that contribute to the coefficient of $s^{\underline{i}}$ in $P_{\underline{j}}(s)$. This coefficient is then

$$b_k^{(n)} = \frac{1}{k!} \sum_{\mathbf{k} \in Q_m(i, j)} \frac{k!}{k_0!k_1!\dots} \cdot \pi^{-\sum_{\ell=1}^{\infty} \ell \cdot k_{\ell}} = \frac{r_{i, j}^{(m)}}{k!}. \quad \square$$

Lemma 4.3.6. *Suppose that $j \geq i \geq 0$. Then $r_{ij}^{(m)}$ is the coefficient of $Z^{\underline{j}}$ in $\log_{LT}(Z)^{\underline{i}}$.*

Proof. Write $\log_{LT}(Z)^k = \sum_{n=k}^{\infty} d_n^{(k)} Z^n$. Then

$$\begin{aligned} \sum_{n=0}^{\infty} P_n(Y) Z^n &= \exp(Y \log_{LT}(Z)) = \sum_{k=0}^{\infty} \frac{1}{k!} \log_{LT}(Z)^k Y^k \\ &= \sum_{k=0}^{\infty} \frac{1}{k!} \sum_{n=k}^{\infty} d_n^{(k)} Z^n Y^k. \end{aligned}$$

Equating the coefficient of $Z^n Y^k$ shows that

$$b_k^{(n)} = \frac{1}{k!} d_n^{(k)} \quad \text{for } 1 \leq j \leq n.$$

Applying Proposition 4.3.5(2), we have $r_{i, j}^{(m)} = \underline{i}! b_{\underline{i}}^{(j)} = d_{\underline{j}}^{(\underline{i})}$. \square

Corollary 4.3.7. *Define polynomials $R_j^{(m)}(t) \in L[t]$ for $j \geq 0$ by the formula*

$$R_j^{(m)}(t) := \sum_{i=0}^j \frac{r_{i, j}^{(m)}}{(\underline{i})!} t^{\underline{i}}.$$

Then for all $j \geq 0$ we have $P_{\underline{j}}(s) = s^m \cdot R_{\underline{j}}^{(m)}(s^{q-1})$.

Lemma 4.3.8. *For each $j \geq i \geq 0$ there exist $\sigma_{i, j}(Y) \in \text{Int}(o_L, o_L)$ such that*

$$P_j(Ys) = \sum_{i=0}^j \sigma_{i, j}(Y) P_i(s).$$

Proof. By Example 4.2.7, $\{P_n(\Omega) : n \geq 0\}$ forms a regular basis for the admissible subalgebra $\sum_{n=0}^{\infty} o_K P_n(\Omega)$ of $L[\Omega]$. Apply Lemma 4.2.8 and use the transcendence of Ω over L . \square

Of course this is just another way of rephrasing Corollary 3.2.6. We will now see that the matrix of polynomials $(\sigma_{i,j}(Y))_{i,j}$ is sparse as well.

Proposition 4.3.9. *Let $j \geq 0$ and suppose that $0 \leq k \leq j$.*

- (1) $\sigma_{k,j}(Y) = 0$ if $k \not\equiv m \pmod{q-1}$.
- (2) For each $i = 0, \dots, j$ there exists $\tau_{i,j}^{(m)}(X) \in L[X]$ such that

$$\sigma_{i,j}(Y) = Y^m \cdot \tau_{i,j}^{(m)}(Y^{q-1}).$$

Proof. Using Lemma 4.3.8, we have

$$P_j(Ys) = \sum_{k=0}^j \sigma_{k,j}(Y) P_k(s).$$

Dividing both sides by $Y^m s^m$ we obtain an equality of Laurent polynomials

$$(16) \quad R_j^{(m)}(Y^{q-1}s^{q-1}) = \sum_{k=0}^j Y^{-m} \sigma_{k,j}(Y) \cdot s^{-m} P_k(s).$$

The left hand side of (16) is a polynomial in s^{q-1} with coefficients in $L[Y]$. The Laurent polynomial $s^{-m} P_k(s)$ lies in $s^{k-m} L[s^{q-1}, s^{1-q}]$ by Proposition 4.3.5. Since

$$L[Y][s, s^{-1}] = \bigoplus_{c=0}^{q-2} s^c L[Y][s^{q-1}, s^{1-q}],$$

looking at the component of the right hand side of (16) that lies in the space $s^c L[Y][s^{q-1}, s^{1-q}]$ for $c \in \{1, \dots, q-2\}$ and then looking at the leading coefficient of $s^{-m} P_k(s)$ implies (1).

Using Corollary 4.3.7, we can now rewrite (16) as follows:

$$(17) \quad R_j^{(m)}(Y^{q-1}s^{q-1}) = \sum_{i=0}^j Y^{-m} \sigma_{i,j}(Y) \cdot R_i^{(m)}(s^{q-1}).$$

Since the left hand side of (17) is now a polynomial in Y^{q-1} with coefficients in $L[s^{q-1}]$, we deduce by looking at the right hand side of (17) that the *a priori* Laurent polynomial $Y^{-m} \sigma_{i,j}(Y)$ in Y in fact lies in $L[Y^{q-1}]$. Part (2) follows. \square

Setting $t = s^{q-1}$ and $X = Y^{q-1}$, we deduce the following

Corollary 4.3.10. *The polynomials $R_j^{(m)}(tX)$ satisfy*

$$R_j^{(m)}(tX) = \sum_{i=0}^j \tau_{i,j}^{(m)}(X) R_i^{(m)}(t).$$

Definition 4.3.11. Consider the following infinite upper-triangular matrices.

- (1) $[r^{(m)}]_{ij} = r_{ij}^{(m)}$ for $j \geq i \geq 0$,
- (2) $\mathcal{T}_{ij}^{(m)} = \tau_{i,j}^{(m)}(X)$, and

$$(3) \mathcal{D}_X := \text{diag}(1, X, X^2, \dots).$$

Lemma 4.3.12. *We have the matrix equation*

$$r^{(m)} \cdot \mathcal{T}^{(m)} = \mathcal{D}_X \cdot r^{(m)}.$$

Proof. Note that each matrix appearing on the right hand side has infinitely many rows and columns, but each one is also upper triangular, so matrix multiplication makes sense. Moreover, as $r_{jj}^{(m)} = 1$ for all $j \geq 0$ by Lemma 4.3.4, the matrix $r^{(m)}$ is invertible, with inverse matrix having entries in L .

Substitute the definition of $R_j^{(m)}(t)$ from Corollary 4.3.7 into Corollary 4.3.10 to obtain

$$\sum_{\ell=0}^j \frac{r_{\ell,j}^{(m)}}{(\ell)!} t^\ell X^\ell = \sum_{i=0}^j \tau_{i,j}^{(m)}(X) \sum_{\ell=0}^i \frac{r_{\ell,i}^{(m)}}{(\ell)!} t^\ell.$$

Equate the coefficients of t^ℓ to get

$$r_{\ell,j}^{(m)} X^\ell = \sum_{i=0}^j \tau_{i,j}^{(m)}(X) \cdot r_{\ell,i}^{(m)}.$$

The right hand side is the (ℓ, j) -th entry of $r^{(m)} \cdot \mathcal{T}^{(m)}$. The left hand side is the (ℓ, j) -th entry of $\mathcal{D}_X \cdot r^{(m)}$. The result follows. \square

The following two results on the coefficients $r_{i,j}^{(m)}$ are strictly speaking not needed for the calculations appearing in Appendix A, but they are nevertheless interesting in their own right.

Lemma 4.3.13. *For each $j \geq i \geq 0$, we have*

$$r_{i,j}^{(m)} = \left(\sum_{\mathbf{k} \in Q_m(i,j)} \binom{i}{k_0; k_1; \dots} \pi^{\sum_{\ell=1}^{\infty} k_\ell \left(\frac{q^\ell - 1}{q - 1} - \ell \right)} \right) \cdot \pi^{i-j}.$$

Proof. Let $\mathbf{k} \in Q_m(i, j)$. Then $\sum_{\ell=1}^{\infty} k_\ell \left(\frac{q^\ell - 1}{q - 1} \right) = j - i$, and therefore

$$\pi^{\sum_{\ell=1}^{\infty} k_\ell \left(\frac{q^\ell - 1}{q - 1} - \ell \right)} \cdot \pi^{i-j} = \pi^{j-i} \cdot \pi^{-\sum_{\ell=1}^{\infty} \ell k_\ell} \cdot \pi^{i-j} = \pi^{-\sum_{\ell=1}^{\infty} \ell k_\ell}.$$

The result now follows from Definition 4.3.3. \square

Proposition 4.3.14. *Let $j \geq i \geq 0$. Then*

- (1) $\pi^{j-i} \cdot r_{i,j}^{(m)} \in o_L$, and
- (2) $\pi^{j-i} \cdot r_{i,j}^{(m)} \equiv \binom{i}{j-i} \pmod{\pi^{q-1} o_L}.$

Proof. (1) Note that for every $\ell \geq 1$ we have

$$\begin{aligned} \alpha_\ell := \frac{q^\ell - 1}{q - 1} - \ell &= \frac{(1 + (q-1))^\ell - 1}{q - 1} - \ell = \frac{1 + \ell(q-1) + \binom{\ell}{2}(q-1)^2 + \dots + (q-1)^{\ell-1} - 1}{q - 1} - \ell \\ &= \binom{\ell}{2}(q-1) + \binom{\ell}{3}(q-1)^2 + \dots + (q-1)^{\ell-1}. \end{aligned}$$

Thus $\alpha_\ell \geq 0$ always. Hence the expression in the big brackets in Lemma 4.3.13 lies in o_L .

(2) The exponent of π appearing in the term in the sum corresponding to $\mathbf{k} \in Q_m(i, j)$ is equal to $\sum_{\ell=1}^{\infty} k_\ell \alpha_\ell$. It follows from the formula for α_ℓ established above that $\alpha_1 = 0$. Hence this exponent is a positive multiple of $q - 1$, unless $k_\ell = 0$ for all $\ell \geq 2$. In this case, the exponent is 0 and the corresponding term is equal to $\binom{i}{j-i}$ because in this case $k_1 = \sum_{\ell=1}^{\infty} k_\ell \frac{q^\ell - 1}{q - 1} = j - i$. \square

5. CONSEQUENCES OF THE KATZ ISOMORPHISM

5.1. Equivariant endomorphisms of L_∞ . Throughout this §, we assume that $L = \mathbb{Q}_{p^2}$ and that $\pi = p$. In particular, L_∞ is the completion of $L(\mathcal{G}[p^\infty])$. We recall the statement of the Katz isomorphism (Theorem 3.6.15): if S is a π -adically complete o_L -algebra, then the map

$$\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), S) \rightarrow S[[Z]]^{\psi_q\text{-int}}$$

is an isomorphism. Recall ([19], page 58), that μ is said to be supported in o_L^\times if and only if $\mu(f) = 0$ for all f such that $f = 0$ on o_L^\times . We have the following criterion.

Lemma 5.1.1. *A measure $\mu \in \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), S)$ is supported in o_L^\times if and only if $\psi_q(\mathcal{K}^*(\mu)) = 0$.*

Proof. Note that $f \in \mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p})$ is zero on o_L^\times if and only if $f = \psi_C(g)$ for some $g \in \mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p})$. Indeed, $\psi_C(g)(a) = 0$ if $a \in o_L^\times$ by definition, and if $f = 0$ on o_L^\times , then $f = \psi_C \varphi_C(f)$.

The map \mathcal{K}^* is injective by Theorem 3.6.15. By Corollary 3.3.7, we have $\psi_q \mathcal{K}^* = \mathcal{K}^* \psi_C^*$. Hence $\psi_q \mathcal{K}^*(\mu) = 0 \Leftrightarrow \mathcal{K}^* \psi_C^*(\mu) = 0 \Leftrightarrow \psi_C^*(\mu) = 0 \Leftrightarrow \mu(\psi_C(g)) = 0$ for all $g \Leftrightarrow \mu$ is supported in o_L^\times . \square

There is the usual $G_L, *$ action on $o_{\mathbb{C}_p}[[Z]]$, and on $\text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})$ it is given by $g^*(\mu)(f) = g(\mu(g^{-1}(f))) = g(\mu(a \mapsto f(\tau(g)^{-1} \cdot a)))$ since f is Gal continuous. In particular, Theorem 3.6.15 applied with $S = o_{\mathbb{C}_p}$ implies the following.

Corollary 5.1.2. *We have*

- (1) $\text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})^{G_L, *} = \Lambda_L(\mathfrak{X})^{\psi_q\text{-int}}.$
- (2) $\text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})^{G_L, *} = \Lambda_L(\mathfrak{X})^{\psi_q=0}.$

Since $L = \mathbb{Q}_{p^2}$, the map τ is surjective. Let $\Gamma_L = \text{Gal}(L(\mathcal{G}[p^\infty])/L)$.

Lemma 5.1.3. *The map $\mathcal{C}_{\text{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}) \rightarrow o_\infty$ given by $f \mapsto f(1)$ is an isomorphism of o_L -modules.*

Proof. Since $d = 2$, we know that τ is surjective by Lemma 2.6.4. Now, if $x \in o_\infty$, let $f_x \in \mathcal{C}_{\text{Gal}}^0(o_L^\times, o_{\mathbb{C}_p})$ be given by $f_x(1) = x$ and $f_x(\tau(g)) = g(x)$. Every element of $\mathcal{C}_{\text{Gal}}^0(o_L^\times, o_{\mathbb{C}_p})$ is of this form. \square

Theorem 3.6.15 applied with $S = o_L$ now gives us the following

Theorem 5.1.4. *The map \mathcal{K}^* gives rise to an o_L -linear isomorphism $o_\infty^* \simeq o_L[[Z]]^{\psi_q=0}$.*

Proposition 5.1.5. *The space $\mathrm{Hom}_{o_L}(\mathcal{C}_{\mathrm{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})^{G_L,*}$ is naturally isomorphic to the space of Γ_L -equivariant o_L -linear maps $o_\infty \rightarrow o_\infty$.*

Proof. If $x \in o_\infty$, let $f_x \in \mathcal{C}_{\mathrm{Gal}}^0(o_L^\times, o_{\mathbb{C}_p})$ be as in the proof of Lemma 5.1.3 above. If $\mu \in \mathrm{Hom}_{o_L}(\mathcal{C}_{\mathrm{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})^{G_L,*}$, we define a map $T : o_\infty \rightarrow o_\infty$ by $T(x) = \mu(f_x)$. We have $f_{x+y} = f_x + f_y$ and $f_{ax} = af_x$ if $a \in o_L$ so that T is o_L -linear. In addition, T is Γ_L -equivariant because μ is fixed under the $G_L, *$ -action. Indeed, $g(T(x)) = g(\mu(f_x)) = \mu(g(f_x))$ and $g(f_x)(1) = g(x)$ so that $g(f_x) = f_{g(x)}$. Therefore, $g(T(x)) = T(g(x))$.

Conversely, a Γ_L -equivariant o_L -linear map $T : o_\infty \rightarrow o_\infty$ as above gives an element $\mu \in \mathrm{Hom}_{o_L}(\mathcal{C}_{\mathrm{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})^{G_L,*}$ via $\mu(f_x) = T(x)$. \square

Combining Corollary 5.1.2 and Proposition 5.1.5, we get the following.

Theorem 5.1.6. *We have $\mathrm{End}_{o_L}^{G_L}(o_\infty) \simeq \Lambda_L(\mathfrak{X})^{\psi_q=0}$.*

Corollary 5.1.7. *We have $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if every Γ_L -equivariant o_L -linear map $o_\infty \rightarrow o_\infty$ comes from an element of $o_L[[\Gamma_L]]$.*

Proof. We have $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if $\Lambda_L(\mathfrak{X})^{\psi=0} = \Lambda(o_L^\times)$ by Lemma 5.1.9 below. If $\mu \in \Lambda_L(\mathfrak{X})^{\psi=0}$, then by Corollary 5.1.2 it corresponds to an element of $\mathrm{Hom}_{o_L}(\mathcal{C}_{\mathrm{Gal}}^0(o_L^\times, o_{\mathbb{C}_p}), o_{\mathbb{C}_p})^{G_L,*}$. By Proposition 5.1.5, the element $\mu \in \Lambda_L(\mathfrak{X})^{\psi=0}$ comes from an element $\nu \in o_L[[\Gamma_L]]$. The element μ then corresponds to the image of ν in $\Lambda(o_L^\times)$ via τ . Indeed, if $g \in \Gamma_L$ and T is given by $x \mapsto g(x)$, then it corresponds to $\mu : f_x \mapsto g(x)$ and $g(x) = f_x(\tau(g))$ so that $\mu = \delta_{\tau(g)}$. \square

Using Corollary 5.1.7, we get the following

Theorem 5.1.8. *We have $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$ if and only if every continuous L -linear and G_L -equivariant map $f : L_\infty \rightarrow L_\infty$ comes from the Iwasawa algebra $L \otimes_{o_L} o_L[[\Gamma_L]]$.*

Proof. Indeed, by Corollary 2.10.11, $\Lambda_L(\mathfrak{X}) \cap (L \otimes_{o_L} o_L[[o_L]]) = o_L[[o_L]]$. \square

Lemma 5.1.9. *If $\Lambda_L(\mathfrak{X})^{\psi=0} = \Lambda(o_L^\times)$, then $\Lambda_L(\mathfrak{X}) = o_L[[o_L]]$.*

Proof. If $f \in \Lambda_L(\mathfrak{X})$, then $\delta_1 \cdot \varphi(f) \in \Lambda_L(\mathfrak{X})^{\psi=0}$. So $\varphi(f) \in o_L[[o_L]]$ and $f = \psi_q \varphi(f) \in o_L[[o_L]]$. \square

The following proposition implies that there are no ‘‘Tate trace maps’’ $L_\infty \rightarrow L$ or $L_\infty \rightarrow L_n$ (recall that L_∞ is the completion of $L(\mathcal{G}[p^\infty])$).

Proposition 5.1.10. *Let $f : L_\infty \rightarrow L_\infty$ be a continuous, Γ_L -equivariant and L -linear map. If $f(L_\infty)$ is included in a finite field extension of L , then $f(1) = 0$.*

Proof. We have $\log \Omega \in L_\infty$ and $(g-1)\log \Omega = \log \tau(g)$ if $g \in \Gamma_L$. Hence

$$(g-1)f(\log \Omega) = f((g-1)\log \Omega) = f(\log \tau(g)) = \log \tau(g) \cdot f(1).$$

Hence if $f(1) \neq 0$, then $f(\log \Omega)$ cannot belong to a finite extension of L . \square

Note that a similar result was proved by Fourquaux, see for instance [14, Prop 2.2.1]. Proposition 5.1.10 can be strengthened: almost the same proof gives us the following proposition. Recall that $L_\infty^{\Gamma_{L_n}} = L_n$ by the Ax-Sen-Tate theorem, see Theorem 1 on page 176 of [32].

Proposition 5.1.11. *Let $f : L_\infty \rightarrow L_\infty$ be a continuous, Γ_L -equivariant and L -linear map. If $f \neq 0$, then there exists $a_1 \neq 0$, $a_0 \in L(\mathcal{G}[p^\infty])$ such that $f(L_\infty)$ contains $a_1 \log \Omega + a_0$.*

Proof. We have $\log \Omega \in L_\infty$ and $(g-1)\log \Omega = \log \tau(g)$ if $g \in \Gamma_L$. Take $x \in L(\mathcal{G}[p^\infty])$ such that $f(x) \neq 0$, and choose (note that $f(L_n) \subset L_n$ by the Ax-Sen-Tate theorem) some n such that $x, f(x) \in L_n$. If $g \in \Gamma_n$, then

$$(g-1)f(x \cdot \log \Omega) = f((g-1)(x \cdot \log \Omega)) = f(x \cdot \log \tau(g)) = \log \tau(g) \cdot f(x).$$

Therefore $(g-1)(f(x \cdot \log \Omega) - f(x) \cdot \log \Omega) = 0$ for all $g \in \Gamma_n$, so that $f(x \cdot \log \Omega) - f(x) \cdot \log \Omega \in L_n$ by Ax-Sen-Tate.

We can now take $a_1 = f(x)$ and $a_0 = f(x \cdot \log \Omega) - f(x) \cdot \log \Omega$. \square

This can be strengthened even further. Let L_∞^{alg} denote the vectors in L_∞ that are locally algebraic for the action of the Lie group Γ_L . Let $c(g) = \log \tau(g) = \log \chi_p^\sigma(g)$. The set L_∞^{alg} is the set of $x \in L_\infty$ such that there exists an open subgroup Γ_x of Γ_L and $d \geq 0$ and $x_0 = x, x_1, \dots, x_d \in L_\infty$ such that $g(x) = x_0 + x_1 c(g) + \dots + x_d c(g)^d$ if $g \in \Gamma_x$. Note that technically, these are the locally σ -analytic locally algebraic vectors in L_∞ . However since $L = \mathbb{Q}_{p^2}$, every locally analytic vector of L_∞ is locally σ -analytic (see [4]).

Lemma 5.1.12. *We have $L_\infty^{\text{alg}} = L(\mathcal{G}[p^\infty])[\log \Omega]$.*

Proof. One inclusion is easy. Now take $x \in L_\infty^{\text{alg}}$ and write $g(x) = x_0 + x_1 c(g) + \dots + x_d c(g)^d$ if $g \in \Gamma_x$. On L_∞^{alg} we have the derivative $\nabla : x \mapsto x_1$ and we know (from the theory of locally analytic vectors) that $\nabla^j(x)/j! = x_j$ for all j . In particular, $\nabla(x_d) = 0$, so that $x_d \in L(\mathcal{G}[p^\infty])$. The element $x - x_d \log^d \Omega$ is then in L_∞^{alg} and it is of degree $\leq d-1$, which allows us to prove the Lemma by induction. \square

We see that $\nabla = \frac{d}{d \log \Omega}$. For all n , the map $\nabla : L_n[\log \Omega] \rightarrow L_n[\log \Omega]$ is surjective, and its kernel is L_n . If $f : L_\infty \rightarrow L_\infty$ is a continuous, Γ_L -equivariant and L -linear map, then $f(L_\infty^{\text{alg}}) \subset L_\infty^{\text{alg}}$. In addition, $\nabla = \lim_{g \rightarrow 1} (g-1)/c(g)$ so that $f \circ \nabla = \nabla \circ f$.

Proposition 5.1.13. *Let $f : L_\infty \rightarrow L_\infty$ be a continuous, Γ_L -equivariant and L -linear map. If $f \neq 0$, there exists $n \geq 0$ such that $L_n \cdot f(L_n[\log \Omega])$ contains $L_n[\log \Omega]$.*

Proof. Take $x \in L(\mathcal{G}[p^\infty])$ such that $f(x) \neq 0$ and let $n \geq 0$ be such that $x, f(x) \in L_n$. We prove by induction on d that $L_n \cdot f(L_n[\log \Omega])$ contains $L_n[\log \Omega]_{\deg \leq d}$. In order to do this, we prove that $f(x \cdot \log^d \Omega)$ is a polynomial (in $\log \Omega$) of degree d . The case $d = 0$ follows from the fact that $f(x) \neq 0$. Now assume that the result holds for $d - 1$. We have

$$\nabla f(x \cdot \log^d \Omega) = f(x \cdot \nabla \log^d \Omega) = f(dx \cdot \log^{d-1} \Omega),$$

so that $f(x \cdot \log^d \Omega)$ is a polynomial of degree d . This implies the claim. \square

5.2. The dual of the ring of integers of a p -adic Lie extensions. Recall that $\pi \in o_L$ is a uniformiser and $k_L := o_L/\pi o_L$ is the residue field of L . In this §, L_∞/L is an infinite Galois extension with Galois group $\Gamma = \text{Gal}(L_\infty/L)$. We fix a chain

$$\Gamma \supseteq \Gamma_1 \supseteq \Gamma_2 \supseteq \cdots$$

of open normal subgroups of Γ such that $\bigcap_{n=1}^{\infty} \Gamma_n = 1$.

Definition 5.2.1. Let $n \geq 1$.

- (1) $L_n := L_\infty^{\Gamma_n}$, a finite Galois extension of L with Galois group Γ/Γ_n .
- (2) o_n is the integral closure of o_L in L_n .
- (3) $o_n^* := \text{Hom}_{o_L}(o_n, o_L)$.
- (4) $k_n := o_n/\pi o_n$.
- (5) $k_n^\vee := \text{Hom}_{k_L}(k_n, k_L)$.

Note that o_n and o_n^* are naturally $o_L[\Gamma/\Gamma_n]$ -modules, both free of finite rank as an o_L -module, and k_n and k_n^\vee are $k_L[\Gamma/\Gamma_n]$ -modules, both finite dimensional over k_L .

Remark 5.2.2. Let $n \geq 1$.

- (1) o_n^* can be identified with the *inverse different* $\mathfrak{d}_{L_n/L}^{-1}$ of the extension L_n/L .
- (2) Applying the duality functor $(-)^* = \text{Hom}_{o_L}(-, o_L)$ to the natural inclusion of o_L -modules $o_n \rightarrow o_{n+1}$, we obtain a natural connecting map $o_{n+1}^* \rightarrow o_n^*$. This map is surjective, because the o_{n+1}/o_n is a finitely generated and torsion-free o_L -module.

Lemma 5.2.3. *For each $n \geq 1$, there is a short exact sequence of $o_L[\Gamma/\Gamma_n]$ -modules*

$$0 \rightarrow o_n^* \xrightarrow{\pi} o_n^* \rightarrow k_n^\vee \rightarrow 0.$$

Proof. Let M be an o_L -module and consider the complex of o_L -modules

$$0 \rightarrow M^* \xrightarrow{\pi} M^* \xrightarrow{\eta_M} (M/\pi M)^\vee \rightarrow 0$$

where $M^* := \text{Hom}_{o_L}(M, o_L)$, $(M/\pi M)^\vee = \text{Hom}_{k_L}(M/\pi M, k_L)$ and where $\eta_M(f)(m + \pi M) = f(m) + \pi o_L \in k_L$. This complex commutes with finite direct sums and is exact in the case when $M = o_L$. So the complex is exact whenever M is a finitely generated free o_L -module. If M also happens to be

an $o_L[G]$ -module for some group G , then the maps in the complex are $o_L[G]$ -linear. The result follows when we set $M = o_n$, an $o_L[\Gamma/\Gamma_n]$ -module which is free of finite rank as an o_L -module. \square

We now pass to the limit as $n \rightarrow \infty$.

Definition 5.2.4. Recall the Iwasawa algebras $\Lambda(\Gamma) = \varprojlim o_L[\Gamma/\Gamma_n]$ and $\Omega(\Gamma) = \varprojlim k_L[\Gamma/\Gamma_n]$.

- (1) $o_\infty := \operatorname{colim} o_n$, an $o_L[\Gamma]$ -module.
- (2) $o_\infty^* := \varprojlim o_n^*$, a $\Lambda(\Gamma)$ -module.
- (3) $k_\infty := \operatorname{colim} k_n$, a $k_L[\Gamma]$ -module.
- (4) $k_\infty^\vee := \varprojlim k_n^\vee$, an $\Omega(\Gamma)$ -module.

Lemma 5.2.5. *There is a short exact sequence of $\Lambda(\Gamma)$ -modules*

$$0 \rightarrow o_\infty^* \xrightarrow{\pi} o_\infty^* \rightarrow k_\infty^\vee \rightarrow 0.$$

Proof. The short exact sequences from Lemma 5.2.3 are compatible with variation in n , in other words we get a short exact sequence of towers of $\Lambda(\Gamma)$ -modules. Applying the inverse limit functor gives a long exact sequence

$$0 \rightarrow o_\infty^* \xrightarrow{\pi} o_\infty^* \rightarrow k_\infty^\vee \rightarrow \varprojlim^{(1)} o_n^*.$$

The $\varprojlim^{(1)}$ term on the right vanishes in view of Remark 5.2.2(2), whence the result. \square

Remark 5.2.2(2) also implies that the natural maps $o_\infty^* \rightarrow o_n^*$ are surjective.

Proposition 5.2.6. *The $\Lambda(\Gamma)$ -modules o_∞ and o_∞^* are faithful.*

Proof. Suppose $\xi \in \Lambda(\Gamma)$ kills o_∞ . Then its image $\xi_n \in o[\Gamma/\Gamma_n]$ kills o_n . Therefore $\xi_n \in L[\Gamma/\Gamma_n]$ kills $L_n = o_n \otimes_{o_L} L$. But L_n is a free $L[\Gamma/\Gamma_n]$ -module of rank 1 by the Normal Basis Theorem. So, $\xi_n = 0$ for all $n \geq 0$ and therefore $\xi = 0$ as well.

Suppose now $\xi \in \Lambda(\Gamma)$ kills o_∞^* . Then ξ kills each the quotients o_n^* of o_∞^* . But the action of $\Lambda(\Gamma)$ on o_n^* factors through $o_L[\Gamma/\Gamma_n]$, so the image ξ_n of ξ in $o_L[\Gamma/\Gamma_n]$ kills o_n^* . Since ξ_n also kills $o_n \cong (o_n^*)^*$, we deduce from the above that $\xi_n = 0$ for all n . Hence $\xi = 0$. \square

Proposition 5.2.7. *Suppose that $p \nmid |\Gamma/\Gamma_1|$. Then k_1^\vee is a free $k_L[\Gamma/\Gamma_1]$ -module of rank 1.*

Proof. The field extension L_1/L is tamely ramified by our assumption on $|\Gamma/\Gamma_1|$. Now it follows from Noether's Theorem on rings of integers in tamely ramified extensions that o_1 is a free $o_L[\Gamma/\Gamma_1]$ -module of rank one — see, e.g. [33, Proposition 2.1]. Hence $o_1/\pi o_1$ is a free $k_L[\Gamma/\Gamma_1]$ -module of rank one, and we can apply Lemma 5.2.3 to conclude. \square

Lemma 5.2.8. *Suppose that Γ is a p -adic Lie group. Let $M = \varprojlim M_n$ be an inverse limit of a tower of $\Omega(\Gamma)$ -modules, where each M_n is finite dimensional over k_L . Then the natural map on Γ -coinvariants*

$$M_\Gamma \rightarrow \varprojlim (M_n)_\Gamma$$

is an isomorphism.

Proof. The Iwasawa algebra $\Omega(\Gamma)$ is Noetherian, so its augmentation ideal $J = (\Gamma - 1)\Omega(\Gamma)$ is finitely generated. Let $u_1, \dots, u_r \in J$ be generators and let N be an $\Omega(\Gamma)$ -module; then

$$N_\Gamma = N/(\Gamma - 1) \cdot N = N/JN = N/(u_1N + \dots + u_rN).$$

In other words, we have the short exact sequence of k_L -vector spaces

$$(18) \quad N^r \xrightarrow{(u_1, \dots, u_r)} N \rightarrow N_\Gamma \rightarrow 0.$$

Applying this to each M_n , we obtain an exact sequence of towers of $\Omega(\Gamma)$ -modules

$$M_n^r \xrightarrow{(u_1, \dots, u_r)} M_n \rightarrow (M_n)_\Gamma \rightarrow 0$$

where each term is a finite dimensional k_L -vector space. The inverse limit functor is exact on such towers, since they all satisfy the Mittag-Leffler condition. So passing to the inverse limit we obtain the exact sequence of k_L -vector spaces

$$M^r \xrightarrow{(u_1, \dots, u_r)} M \rightarrow \varprojlim (M_n)_\Gamma \rightarrow 0.$$

Comparing this with (18) applied with $N = M$ gives the result. \square

Theorem 5.2.9. *Suppose that*

- Γ is abelian,
- $p \nmid |\Gamma/\Gamma_1|$,
- Γ_1 is a torsionfree pro- p group of finite rank.

Then o_∞^ is a free $\Lambda(\Gamma)$ -module of rank 1 if and only if the map $k_1 \rightarrow k_\infty^{\Gamma_1}$ is an isomorphism.*

Proof. (\Leftarrow) Note that the connecting maps $k_n \rightarrow k_{n+1}$ in the colimit $k_\infty := \operatorname{colim} k_n$ are injective: if $x + \pi o_n \in k_n$ maps to zero in k_{n+1} then there is $y \in o_{n+1}$ such that $x = \pi y$; but then $y \in L_n \cap o_{n+1} = o_n$ and hence $x = \pi y \in \pi o_n$. Under our hypothesis that $k_1 \rightarrow k_\infty^{\Gamma_1}$ is an isomorphism, it follows that for each $n \geq 1$, the map $k_n^{\Gamma_1} \rightarrow k_{n+1}^{\Gamma_1}$ is an isomorphism. Applying the $(-)^{\vee} = \operatorname{Hom}_{k_L}(-, k_L)$ functor, we deduce that for each $n \geq 1$, the map on Γ_1 -coinvariants

$$(k_{n+1}^{\vee})_{\Gamma_1} \rightarrow (k_n^{\vee})_{\Gamma_1}$$

is an isomorphism. Now, Lemma 5.2.8 tells us that

$$(k_\infty^{\vee})_{\Gamma_1} \cong \varprojlim (k_n^{\vee})_{\Gamma_1}.$$

Since the maps in the tower of Γ_1 -coinvariants are all isomorphisms, we conclude that the natural map of $k[\Gamma/\Gamma_1]$ -modules

$$(k_\infty^\vee)_{\Gamma_1} \rightarrow k_1^\vee$$

must be an isomorphism. Now k_1^\vee is a cyclic $k_L[\Gamma/\Gamma_1]$ -module by Proposition 5.2.7 and the ideal $J\Omega(\Gamma)$ generated by the augmentation ideal J of $\Omega(\Gamma_1)$ is topologically nilpotent in the sense that $J^n \rightarrow 0$ as $n \rightarrow \infty$, because Γ_1 is assumed to be pro- p . In this situation we can apply the Nakayama Lemma for compact Λ -modules — see [3, Corollary to Theorem 3] — to deduce that k_∞^\vee is a cyclic $\Omega(\Gamma)$ -module: any lift of a $k_L[\Gamma/\Gamma_1]$ -module generator for k_1^\vee to k_∞^\vee will generate it as an $\Omega(\Gamma)$ -module.

Now $o_\infty^*/\pi o_\infty^* \cong k_\infty^\vee$ by Lemma 5.2.5. The $\Lambda(\Gamma)$ -module o_∞^* is profinite and $\pi^n \rightarrow 0$ as $n \rightarrow \infty$ in $\Lambda(\Gamma)$, so applying the Nakayama Lemma again, we conclude that o_∞^* is a cyclic $\Lambda(\Gamma)$ -module.

Since o_∞^* is a faithful $\Lambda(\Gamma)$ -module by Proposition 5.2.6 and since Γ is abelian, we deduce that o_∞^* must be a free $\Lambda(\Gamma)$ -module of rank 1.

(\Rightarrow) We reverse the argument above. Assume o_∞^* is a free $\Lambda(\Gamma)$ -module of rank 1. Then Lemma 5.2.5 implies that k_∞^\vee is a free $\Omega(\Gamma)$ -module of rank 1. Hence $(k_\infty^\vee)_{\Gamma_1}$ is a free $k[\Gamma/\Gamma_1]$ -module of rank 1. By Lemma 5.2.8 we have $(k_\infty^\vee)_{\Gamma_1} \cong \varprojlim (k_n^\vee)_{\Gamma_1}$ and the connecting maps in the tower $(k_n^\vee)_{\Gamma_1}$ are surjective, with the bottom term being $(k_1^\vee)_{\Gamma_1} = k_1^\vee$. Since this is a free $k_L[\Gamma/\Gamma_1]$ -module of rank 1 by Proposition 5.2.7, the natural map $(k_\infty^\vee)_{\Gamma_1} \rightarrow k_1^\vee$ from the inverse limit to the bottom term is a surjection between two free $k_L[\Gamma/\Gamma_1]$ -modules of rank 1. So it is also an isomorphism. Dualising shows that $k_1 \rightarrow k_\infty^{\Gamma_1}$ is an isomorphism as well. \square

Lemma 5.2.10. *In the situation of Proposition 5.2.9, suppose that o_∞^* is a free $\Lambda(\Gamma)$ -module of rank 1. Then L_n/L is tamely ramified for all $n \geq 1$.*

Proof. Consider the Γ_n -coinvariants of o_∞^* . These coinvariant must be a free rank 1 $o_L[\Gamma/\Gamma_n]$ -module by assumption. On the other hand, by construction, there's a surjective $o_L[\Gamma/\Gamma_n]$ -linear map

$$(o_\infty^*)_{\Gamma_n} \rightarrow o_n^*$$

(see the remark just before Proposition 5.2.6). Both sides are free o_L -modules of rank $[L_n : L]$, so this surjective map must actually be an isomorphism by the rank-nullity theorem. So, o_n^* is a free rank 1 $o_L[\Gamma/\Gamma_n]$ -module. But then using, for example [2, Lemma 5.4], we see that

$$o_n = \mathrm{Hom}_{o_L}(o_n^*, o_L) = \mathrm{Hom}_{o_L[\Gamma/\Gamma_n]}(o_n^*, o_L[\Gamma/\Gamma_n])$$

must also be a free rank 1 $o_L[\Gamma/\Gamma_n]$ -module. In other words, o_n has an integral normal basis, so by [33, Proposition 2.1] L_n/L must be tamely ramified. \square

The following result, which may be of independent interest, shows that the hypothesis that the action map $\rho : \Omega(\Gamma) \rightarrow \mathrm{End}_{\Omega(\Gamma)}(k_\infty^\vee)$ is an isomorphism has strong implications about ramification behaviour in the tower L_∞/L .

Lemma 5.2.11. *Suppose Γ is a torsionfree abelian pro- p group of finite rank, and that the action map $\rho : \Omega(\Gamma) \rightarrow \text{End}_{\Omega(\Gamma)}(k_\infty^\vee)$ is an isomorphism. Assume that $\Gamma_1 = \Gamma$. Then L_n/L is tamely ramified for all $n \geq 1$.*

Proof. Let $a \in k_\infty^{\Gamma_1}$ and consider the multiplication-by- a map $\ell_a : k_\infty \rightarrow k_\infty$. Since a is fixed by $\Gamma = \Gamma_1$, this map is $\Omega(\Gamma)$ -linear. By our assumption on ρ , we can find some $b \in \Omega(\Gamma)$ such that $\rho(b) = a$. Now a is algebraic over k_L and ρ is injective by assumption, so $b \in \Omega(\Gamma)$ must be algebraic over k_L as well. Since $\Gamma = \Gamma_1$, the mod- p Iwasawa algebra $\Omega(\Gamma)$ is a power series ring over k_L in finitely many variables. The only elements of such a power series ring that are algebraic over k_L are constants. Hence $b \in k_L$ and so $a \in k_L = k_1$ since $\Gamma = \Gamma_1$. Hence $k_\infty^{\Gamma_1} = k_1$. Now the result follows from Theorem 5.2.9 and Lemma 5.2.10. \square

Returning to the setting of §1.7, we have the following conclusion.

Corollary 5.2.12. *Suppose that $L = \mathbb{Q}_{p^2}$ and $\pi = p$, and let \mathcal{G} be the Lubin-Tate formal group attached to π . Let $\Gamma_L^{LT} = \text{Gal}(L(\mathcal{G}[p^\infty])/L)$. Then $o_L[[Z]]^{\psi_q=0}$ is not a free $o_L[[\Gamma_L^{LT}]]$ -module of rank 1.*

Proof. It is well known that L_n/L is not tamely ramified for any $n \geq 2$. Hence o_∞^* is not a free $\Lambda(\Gamma_L^{LT})$ -module of rank 1 by Lemma 5.2.10. Since \mathcal{G} is self-dual, the tower L_∞/L coincides with the one defined at Definition 2.7.1(1). The result now follows from Theorem 1.7.1. \square

5.3. The operator ψ and the span of the P_n . We now turn to some consequences of the Katz isomorphism for the span of the P_n , where P_n is the element of $\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p})$ given by $a \mapsto P_n(a \cdot \Omega)$. The Katz map $\mathcal{K}^* : \text{Hom}_{o_L}(\mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p}), S) \rightarrow S[[Z]]^{\psi_q\text{-int}}$ is then given by $\mu \mapsto \sum_{n \geq 0} \mu(P_n)Z^n$.

Proposition 5.3.1. *The L -span of the P_n is dense in the L -Banach space $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$.*

Proof. Let W denote the closure of the L -span of the P_n in $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$. If $W \neq \mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$, then it has a closed complement in $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$ and we can find a measure $\mu \neq 0$ that is zero on W (and hence on all of the P_n). This is a contradiction. \square

Remark 5.3.2. There is another proof of this result. Indeed, locally analytic functions are dense in $\mathcal{C}^0(o_L, \mathbb{C}_p)$ and for locally analytic functions, we have the generalized Mahler expansion of [28, Theorem 4.7]. So it is enough to prove that locally analytic and Gal continuous functions are dense in $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$.

A Gal-continuous function is determined by $(f(p^n))_{n=0}^\infty$ where each $f(p^n) \in L_\infty$ and $f(0) \in L$ and $f(p^n) \rightarrow f(0)$ (see also §§3.3-3.4). We can approximate each $f(p^n)$ by an element of L_∞ and this way, we can show that Gal-continuous locally constant functions are dense in the Gal-continuous functions. More precisely, given a sequence $\{f_n\}$ as above and some $k \geq 0$, we have $f_n - f_\infty \in p^k o_{\mathbb{C}_p}$ for all $n \geq n(k)$, so we replace these f_n by f_∞ , and approximate the others to within p^{-k} .

We now choose a coordinate X on LT such that $[p]_{\text{LT}}(X) = pX + X^q$. The polynomials P_i depend on the choice of coordinate. However, the \mathcal{O}_L -module $U_n = \bigoplus_{i=0}^n \mathcal{O}_L \cdot P_i$ is independent of the coordinate. Given this choice of coordinate, we have formulas and estimates for ψ_q in [15, §2A].

Lemma 5.3.3. *If $k \geq 1$, then $\psi_q(X^k) \in L[X]_{k-1}$.*

Proof. See [15, Proposition 2.2]. \square

Let $c^0(A)$ denote the set of sequences $\{c_n\}_{n \geq 0}$ with $c_n \in A$ and $c_n \rightarrow 0$ (with $A = \mathcal{O}_L$ or L).

Corollary 5.3.4. *The map $c^0(\mathcal{O}_L) \rightarrow \mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathcal{O}_{\mathbb{C}_p})$ given by $\{c_i\}_{i \geq 0} \mapsto \sum c_i P_i$ is injective, as well as the same map $c^0(L) \rightarrow \mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathbb{C}_p)$.*

Proof. Lemma 5.3.3 implies that for all $k \geq 0$, there exists $n = n(k)$ such that $p^n X^k \in \mathcal{O}_L[X]^{\psi_q\text{-int}}$. Let μ be the corresponding measure. We have $\mu(\sum_{i \geq 0} c_i P_i) = p^n c_k$ hence if $\sum_{i \geq 0} c_i P_i = 0$, then $c_k = 0$. The second assertion follows from the first. \square

Lemma 5.3.5. *If $k \geq 1$, then $\psi_q(p^k \cdot \mathcal{O}_L[X]_{q^k}) \subset p^{k-1} \cdot \mathcal{O}_L[X]_{q^{k-1}}$.*

Proof. This follows from [15, Proposition 2.2]. \square

Let $H_n \subset L[\Omega]$ denote the set of $P(\Omega)$ such that $\deg P \leq n$ and $P(a\Omega) \in \mathcal{O}_{\mathbb{C}_p}$ for all $a \in \mathcal{O}_L$. Obviously, $U_n = \bigoplus_{i=0}^n \mathcal{O}_L \cdot P_i(\Omega) \subset H_n$. Let $\mu_i : \mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathcal{O}_{\mathbb{C}_p}) \rightarrow L$ be the measure corresponding to X^i , so that $\mu_i(P_j) = \delta_{ij}$.

Proposition 5.3.6. *If $Q(\Omega) = \sum_{i=0}^n c_i P_i(\Omega) \in H_n$, then $c_i \in p^{-m} \mathcal{O}_L$ if $i \leq q^m$.*

Proof. We have $Q(\Omega) \in \mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathcal{O}_{\mathbb{C}_p})$. By Lemma 5.3.5, $p^m X^i \in \mathcal{O}_L[X]^{\psi_q\text{-int}}$ if $i \leq q^m$, and hence $p^m \mu_i \in \text{Hom}_{\mathcal{O}_L}(\mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathcal{O}_{\mathbb{C}_p}), \mathcal{O}_L)$ for all $0 \leq i \leq q^m$. Hence $p^m c_i \in \mathcal{O}_L$. \square

Corollary 5.3.7. *We have $H_{q^k} \subset p^{-k} U_{q^k}$.*

Let $\psi_p = p \cdot \psi_q$ so that $\psi_p(\mathcal{O}_L[X]) \subset \mathcal{O}_L[X]$.

Lemma 5.3.8. *$\psi_p(X^{q^k+(q-1)}) = X^k \bmod p$ and $\psi_p(X^m) = 0 \bmod p$ if $m \neq -1 \bmod q$.*

Proof. This follows from [15, Proposition 2.2]. \square

Corollary 5.3.9. *The map $c^0(L) \rightarrow \mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathbb{C}_p)$ is not surjective.*

Proof. By Corollary 5.3.4, it is injective. If it is a bijection, then the continuous dual of $\mathcal{C}_{\text{Gal}}^0(\mathcal{O}_L, \mathbb{C}_p)$ is naturally isomorphic to $\mathcal{O}_L[X][1/p]$ via the map $\mu \mapsto \sum_{n \geq 0} \mu(P_n) X^n$. However by the Katz isomorphism, the image of this map is $\mathcal{O}_L[X]^{\psi_q\text{-int}}[1/p]$.

Take $f(X) = 1 + X^{q-1} + X^{q^2-1} + \dots$. Lemma 5.3.8 implies that $\psi_p(f) = f \bmod p$ and hence $\psi_p^n(f) = f \bmod p$. We therefore have $\psi_q^n(f) \in p^{-n} f + p^{-(n-1)} \mathcal{O}_L[X]$ for all $n \geq 1$, so that $f(X)$ is not in $\mathcal{O}_L[X]^{\psi_q\text{-int}}[1/p]$. Hence $\mathcal{O}_L[X][1/p] \neq \mathcal{O}_L[X]^{\psi_q\text{-int}}[1/p]$. \square

In order to say more using Katz' result, we need to produce more elements of $o_L[[X]]^{\psi_q\text{-int}}$. There is $o_L[[X]]^{\psi_q=0}$, which contains X^i for $1 \leq i \leq q-2$ and $pX^{q-1} + (q-1)$ and hence $(\oplus_{i=1}^{q-2} X^i \cdot \varphi_q(o_L[[X]])) \oplus (pX^{q-1} + (q-1)) \cdot \varphi_q(o_L[[X]])$. If $f_n(X) \in (X \cdot o_L[[X]])^{\psi_q\text{-int}}$ and the b_n are in o_L , then $\sum_{n \geq 0} b_n \varphi_q^n(f_n) \in o_L[[X]]^{\psi_q\text{-int}}$ as well (the sum converges for the weak topology, and ψ_q is continuous for that topology). For example, if $f(X) \in (X \cdot o_L[[X]])^{\psi_q=0}$, then $\sum_{n \geq 0} \varphi_q^n(f) \in o_L[[X]]^{\psi_q=1}$.

Remark 5.3.10. We have

- (1) $\psi_q(X^i) = 0$ if $1 \leq i \leq q-2$ and $q+1 \leq i \leq 2q-3$ and $2q+1 \leq i \leq 3q-4$
- (2) $\psi_q(1) = 1$ and $\psi_q(X^{q-1}) = (1-q)/p$ and $\psi_q(X^q) = X$
- (3) $\psi_q(X^{2q-2}) = q-1$ and $\psi_q(X^{2q-1}) = X(1/p - 2p)$ and $\psi_q(X^{2q}) = X^2$
- (4) More generally, $\psi_q(X^k) = X\psi_q(X^{k-q}) - p\psi_q(X^{k+1-q})$

Lemma 5.3.11. *We have $p^k X^{q^k-1} \in o_L[[X]]^{\psi_q\text{-int}}$, but not $p^{k-1} X^{q^k-1}$.*

Proof. Recall that $\psi_q(X^{q-1}) = (1-q)/p$. This implies that $\psi_q(1/X) = \psi_q((X^{q-1} + p)/\varphi_q(X)) = 1/pX$. If $k \geq 1$, then

$$\binom{q^{k-1}}{i} \cdot p^i = \binom{q^{k-1}-1}{i-1} \cdot q^{k-1} p^i / i \in p^k o_L.$$

This implies that $\varphi_q(X^{q^{k-1}}) \in X^{q^k} + p^k X o_L[X]_{q^k-1}$. By Lemma 5.3.5, we have

$$\psi_q(X^{q^k-1}) = \psi_q\left(\frac{\varphi_q(X^{q^{k-1}}) + X^{q^k} - \varphi_q(X^{q^{k-1}})}{X}\right) \in \frac{X^{q^{k-1}-1}}{p} + o_L[[X]]^{\psi_q\text{-int}}.$$

This implies the Lemma by induction on k . □

Corollary 5.3.12. *There is an $h \in H$ in which the coefficient of P_{q^k-1} is in $p^{-k} o_L^\times$.*

Proof. Let $c_{q^k-1} \in \mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)^*$ be the linear form corresponding to X^{q^k-1} . There is an $f \in \mathcal{C}_{\text{Gal}}^0(o_L, o_{\mathbb{C}_p})$ such that $c_{q^k-1}(f) \in p^{-k} o_L^\times$ (if it was in $p^{1-k} o_L$ for all f , then $p^{k-1} c_{q^k-1}$ would be an integral linear form, and we'd have $p^{k-1} X^{q^k-1} \in o_L[[X]]^{\psi_q\text{-int}}$. This is not the case by Lemma 5.3.11). By Corollary 5.3.1, the L -span of the P_n is dense in $\mathcal{C}_{\text{Gal}}^0(o_L, \mathbb{C}_p)$. Therefore there is an $h \in H$ such that $\|f - h\| \leq p^{-1}$. We then have $c_{q^k-1}(h) \in p^{-k} o_L^\times$. □

6. OTHER CRITERIA

We indicate how to prove Theorems 1.8.1 and 1.8.2.

6.1. The Lubin-Tate derivative. As we said in the Introduction, Theorem 1.8.1 follows from Theorem 1.4.1 and Proposition 6.1.2 below.

Lemma 6.1.1. *The sum $\sum_{[p](\omega)=0} \omega^n$ is q if $n = 0$, it is 0 if $(q-1) \nmid n$, and it is $(q-1)(-p)^k$ if $n = (q-1)k$ with $k \geq 1$.*

Proof. Since $[p](T) = pT + T^q$, the sum is over 0 and the roots of $T^{q-1} = -p$. If λ is one of the roots, the set of all the roots is $\{\eta\lambda\}_{\eta^{q-1}=1}$. The result follows (for $n = 0$ it is a convention). \square

Proposition 6.1.2. *Assume that $L = \mathbb{Q}_{p^2}$ and that $\pi = p$. Let $\lambda = \Omega^{q-1}/p(q-1)! \in o_{\mathbb{C}_p}^\times$. If $f(Z) \in o_{\mathbb{C}_p}[[Z]]$, then $\varphi\psi_q(f) - \lambda \cdot D^{q-1}(f) \in o_{\mathbb{C}_p}[[Z]]$.*

Proof. Recall from [20, p. 667] that $f(Z \oplus Y) = \sum_{n \geq 0} Y^n P_n(\partial) f(Z)$. We have $\varphi\psi_q(f)(Z) = 1/q \cdot \sum_{[p](\omega)=0} f(Z \oplus \omega)$, so that

$$\varphi\psi_q(f)(Z) = \frac{1}{q} \sum_{[p](\omega)=0} \sum_{n \geq 0} \omega^n P_n(\partial) f(Z) = \frac{1}{q} \sum_{n \geq 0} \left(\sum_{[p](\omega)=0} \omega^n \right) P_n(\partial) f(Z).$$

By Lemma 6.1.1, the $\sum \omega^n$ for n not divisible by $q-1$ are zero, and the $\sum \omega^n$ for $n = (q-1)k$ are divisible by q except when $k = 1$. Hence

$$\varphi\psi_q(f) - \frac{1}{q}(q-1)(-p)P_{q-1}(\partial)(f) \in o_{\mathbb{C}_p}[[Z]].$$

The proposition now follows from the fact that

$$P_{q-1}(\partial) = \frac{\partial^{q-1}}{(q-1)!} = pD^{q-1} \cdot \frac{\Omega^{q-1}}{p(q-1)!} = pD^{q-1} \cdot \lambda. \quad \square$$

6.2. Changing the base field. We now turn to Theorem 1.8.2. If K is a subfield of L , we also have a character variety \mathfrak{X} for K ; write \mathfrak{X}_K and \mathfrak{X}_L . An L -analytic character $\eta : o_L \rightarrow \mathbb{C}_p^\times$ can be restricted to o_K , and it is then K -analytic. This gives a rigid analytic map $\mathfrak{X}_L \rightarrow \mathfrak{X}_K$. This map in turn gives rise to a map $\text{res}_{L/K} : \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}_K) \rightarrow \mathcal{O}_{\mathbb{C}_p}(\mathfrak{X}_L)$, which sends bounded functions to bounded functions, and $\mathcal{O}_M(\mathfrak{X}_K)$ to $\mathcal{O}_M(\mathfrak{X}_L)$ for all closed subfields $L \subset M \subset \mathbb{C}_p$.

Lemma 6.2.1. *On bounded functions, $\text{res}_{L/K} : \mathcal{O}_{\mathbb{C}_p}^b(\mathfrak{X}_K) \rightarrow \mathcal{O}_{\mathbb{C}_p}^b(\mathfrak{X}_L)$ is injective.*

Proof. Suppose that $f \in \mathcal{O}_{\mathbb{C}_p}^b(\mathfrak{X}_K)$ is zero on the restriction to o_K of every L -analytic character of o_L . Since o_K is a direct summand of o_L , every torsion character of o_K extends to a torsion character of o_L . Hence f is zero on all torsion characters of o_K . This implies that $f = 0$ as f is bounded. \square

If μ is a distribution on o_K , we define a distribution $\text{res}_{L/K}(\mu)$ on o_L as follows: if $f \in \mathcal{C}^{\text{an}}(o_L)$, we let $\text{res}_{L/K}(\mu)(f) = \mu(f|_{o_K})$. This is compatible with the above map if we view elements of $\mathcal{O}_{\mathbb{C}_p}(\mathfrak{X})$ as distributions.

Lemma 6.2.2. *If μ is a distribution on o_K , whose image under $\text{res}_{L/K}(\mu)$ is a measure on o_L , then there exists a measure $\tilde{\mu}$ on o_K such that $\mu = \tilde{\mu}$ on $\text{LC}(o_K)$.*

Proof. Let f be a locally constant function on o_K . Since o_K is a direct summand in o_L , we can extend f to a locally constant function \tilde{f} on o_L , in a way that the sup norm of \tilde{f} on o_L is the sup norm of f on o_K . Since $\text{res}_{L/K}(\mu)$ is a

measure, there exists C such that $\|\operatorname{res}_{L/K}(\mu)(g)\|_{o_L} \leq C \cdot \|g\|_{o_L}$ for all locally constant functions g on o_L . We then have

$$\|\mu(f)\|_{o_K} = \|\operatorname{res}_{L/K}(\mu)(\tilde{f})\|_{o_L} \leq C \cdot \|\tilde{f}\|_{o_L} = C \cdot \|f\|_{o_K}.$$

We can now let $\tilde{\mu}(f) = \mu(f)$ for any $f \in \operatorname{LC}(o_K)$. The above estimate shows that $\tilde{\mu}$ extends continuously to $\mathcal{C}^0(o_K)$. \square

Proposition 6.2.3. *If $\mathcal{O}_L^b(\mathfrak{X}_L) = L \otimes_{o_L} \Lambda(o_L)$, then $\mathcal{O}_L^b(\mathfrak{X}_K) = L \otimes_{o_K} \Lambda(o_K)$.*

Proof. If $\mu \in \mathcal{O}_L^b(\mathfrak{X}_K)$, then μ can be seen as a distribution on o_K , and it gives rise via $\operatorname{res}_{L/K}$ to an element of $L \otimes_{o_L} \Lambda(o_L)$. By Lemma 6.2.2, there is a measure $\tilde{\mu}$ on o_K such that $\mu = \tilde{\mu}$ on $\operatorname{LC}(o_K)$. The image of the distribution $\mu - \tilde{\mu}$ under $\operatorname{res}_{L/K}$ belongs to $L \otimes_{o_L} \Lambda(o_L)$ and is zero on locally constant functions, hence $\operatorname{res}_{L/K}(\mu - \tilde{\mu}) = 0$. By Lemma 6.2.1, $\mu = \tilde{\mu}$ and hence μ is a measure on o_K . \square

Theorem 6.2.4. *If L/K is finite and if $\Lambda_L(\mathfrak{X}_L) = o_L[\![o_L]\!]$, then $\Lambda_K(\mathfrak{X}_K) = o_K[\![o_K]\!]$.*

APPENDIX A. AN ALGORITHM FOR WHETHER THE $\sigma_{i,j}$ 'S SPAN $\operatorname{Int}(o_L, o_L)$

DRAGOȘ CRIȘAN AND JINGJIE YANG

Acknowledgements. The authors took part in an Oxford summer project together with Shashidhara Balla, Jonathan Medcalf and William Whitehead. Financial support from the LMS, Brasenose College, University College, Merton College and the Mathematical Institute, Oxford is gratefully acknowledged.

A.1. Introduction. Let $\mathbb{Q}_p \subseteq L \subsetneq \mathbb{C}_p$ be a field of finite degree d over \mathbb{Q}_p , o_L the ring of integers of L , $\pi \in o_L$ a fixed prime element, and $q := |o_L/\pi_L o_L|$ the dimension of the residue field. For an o_L -submodule S of $L[Y]$ and an integer n , let $S_n = \{f \in S : \deg(f) < n\}$.

Recall that the polynomials $P_n(Y)$ are defined by

$$\exp(Y \cdot \log_{\operatorname{LT}}(Z)) = \sum_{n=0}^{\infty} P_n(Y) Z^n.$$

We choose the coordinate Z such that $\log_{\operatorname{LT}}(Z) = \sum_{k=0}^{\infty} \pi^{-k} Z^{q^k}$.

Define the upper-triangular matrix $(\sigma_{i,j})_{i,j \geq 0}$ with entries in $L[Y]$ by

$$P_j(Ys) = \sum_{i=0}^j \sigma_{i,j}(Y) P_i(s).$$

By Lemmas 4.3.8 and 4.2.8, we know that $\sigma_{i,j}(Y) \in \operatorname{Int}(o_L, o_L)$ and that $\deg(\sigma_{i,j}(Y)) \leq j$. The question that we consider is whether the o_L -linear span of $\{\sigma_{i,j}(Y) : 0 \leq i \leq j\}$ equals $\operatorname{Int}(o_L, o_L)$. In this write-up we develop an algorithm to check whether $(\operatorname{Int}(o_L, o_L))_n$ is contained in the o_L -linear span of $\{\sigma_{i,j}(Y) : 0 \leq i \leq j < N\}$ for some fixed N , where for convenience we require $q-1 \mid N$.

A.2. Theory.

A.2.1. *Reduction to $\tau_{i,j}^{(a)}$.* To ease notation, for a fixed $a \in \{0, 1, \dots, q-2\}$, we denote $\underline{i} = a + (q-1)i$. By Proposition 4.3.9(2), there exist upper-triangular matrices $\tau_{i,j}^{(a)}(Y)$ such that

$$(19) \quad \sigma_{\underline{i}, \underline{j}}(Y) = Y^a \cdot \tau_{i,j}^{(a)}(Y^{q-1}).$$

Definition A.2.1. For a polynomial $P(x)$, we denote by $\gamma_n(P)$ the coefficient of x^n in P .

Definition A.2.2. Let M be the o_L -linear span of $\{\sigma_{i,j}(Y) : 0 \leq i \leq j\}$. For a fixed a , let $M^{(a)}$ be the o_L -linear span of $\{\sigma_{\underline{i}, \underline{j}}(Y) : 0 \leq i \leq j\}$. Let $S^{(a)}$ be the o_L -linear span of $\{\tau_{i,j}^{(a)}(Y) : 0 \leq i \leq j\}$.

Lemma A.2.3. Let $(f_b^{(a)})_{b \geq 0}$ be a regular basis for $S^{(a)}$ — that is, each $f_b^{(a)}$ has degree b . Then, $M = \text{Int}(o_L, o_L)$ if and only if for all $a \in \{0, 1, \dots, q-2\}$ and $b \geq 0$, we have

$$\nu_\pi(\gamma_b(f_b^{(a)})) = -w_q(a + b(q-1)).$$

Proof. For a fixed $a \in \{0, 1, \dots, q-2\}$, by (19), we have $\gamma_s(\sigma_{i,j}(Y)) = 0$ if $s \not\equiv j \pmod{q-1}$. So, by definition, $M = \bigoplus_{a=0}^{q-2} M^{(a)}$.

We write $S^{(a)}(Y^{q-1}) = \{f(Y^{q-1}) : f \in S^{(a)}\}$. Equation (19) shows that

$$M^{(a)} = Y^a \cdot N^{(a)}(Y^{q-1}).$$

Having chosen a regular basis $(f_b^{(a)})_{b \geq 0}$, these give rise to regular bases $(f_b^{(a)}(Y^{q-1}))_{b \geq 0}$ for $S^{(a)}(Y^{q-1})$.

So, we get regular bases $(Y^a f_b^{(a)}(Y^{q-1}))_{b \geq 0}$ for $M^{(a)}$ and thus a regular basis $\{Y^a f_b^{(a)}(Y^{q-1}) : a \in \{0, 1, \dots, q-2\}, b \geq 0\}$ for M .

Then, $M = \text{Int}(o_L, o_L)$ is equivalent to $\nu_\pi(\gamma_{a+b(q-1)}(Y^a f_b^{(a)}(Y^{q-1}))) = -w_q(a + b(q-1))$, which is equivalent to $\nu_\pi(\gamma_b(f_b^{(a)})) = -w_q(a + b(q-1))$. \square

Let $n = a + b(q-1)$, where a, b are integers, with $a \in \{0, 1, \dots, q-2\}$. The proof above shows that a polynomial of degree n with π -valuation of leading term equal to $-w_q(n)$ exists in M_N if and only a polynomial of degree b with the same valuation of leading term exists in $S_{N/(q-1)}^{(a)}$. So, the strategy will be to compute regular bases for $S_{N/(q-1)}^{(a)}$.

A.2.2. *A formula for $\tau_{i,j}^{(a)}$.* One advantage of this approach is that the matrices $\tau_{i,j}^{(a)}(Y)$ can be computed quickly. Recall Definition 4.3.3 (where we merely change notation, calling m by a instead):

Definition A.2.4. For each $j \geq i \geq 0$, let

$$Q_a(i, j) := \left\{ \mathbf{k} \in \mathbb{N}^\infty : \sum_{\ell=0}^{\infty} k_\ell = i, \sum_{\ell=1}^{\infty} k_\ell \left(\frac{q^\ell - 1}{q - 1} \right) = j - i \right\};$$

$$r_{i,j}^{(a)} := \sum_{\mathbf{k} \in Q_a(i,j)} \binom{i}{k_0; k_1; \dots} \cdot \pi^{-\sum_{\ell=1}^{\infty} \ell \cdot k_\ell}.$$

Define the upper triangular matrix $(D_{i,j})_{i,j}$ of coefficients as follows:

Definition A.2.5. Let $D_{i,j} = i! \gamma_i P_j(Y)$.

This does not depend on a . From Proposition 4.3.2, we obtain the following recursion formula, valid for $i \geq 1$:

$$D_{i,j} = \sum_{r \geq 0} \pi^{-r} D_{i-1, j-q^r},$$

with the initial conditions being $D_{0,j} = \delta_{0,j}$.

Now, by Proposition 4.3.5(2) it follows that $r_{i,j}^{(a)} = D_{i,j}$. To tie this back to $\tau_{i,j}^{(a)}$, we recall from Definition 4.3.11(3) the notation $\mathcal{D}_Y := \text{diag}(1, Y, Y^2, \dots)$. Then, Lemma 4.3.12 gives $\tau^{(a)} = (r^{(a)})^{-1} \cdot \mathcal{D}_Y \cdot r^{(a)}$. This gives a fast algorithm to compute the matrices $\tau^{(a)}$, as the recurrence relation for D allows us to compute $r^{(a)}$ easily.

A.2.3. Gaussian elimination over a (discrete) valuation ring. Let R be a (discrete) valuation ring and let A be an $m \times n$ matrix with entries in R . We define notions of elementary row operations and row echelon form over R , similarly to the definitions over a field.

Definition A.2.6. Given a matrix A as above, the elementary row operations are as follows.

- (1) Swap two rows.
- (2) Multiply an entire row by a unit in R .
- (3) Add an R -multiple of a row to another row.

Lemma A.2.7. *Performing elementary row operations on a matrix preserves its R -row span.*

Proof. For each elementary row operation on A , we define an $m \times m$ matrix B with entries in R such that the result of applying the elementary row operation on A is BA . Observe that in each case, B is invertible, so BA has the same R -row span as A . \square

Lemma A.2.8 (Gaussian Elimination). *Let A be a matrix as above. Assume that $m \geq n$ and that A has rank n . Then, one can perform a sequence of elementary row operations on A to produce an upper-triangular matrix of rank n .*

Proof. We will exhibit an algorithm that puts A in the required form.

We start with the leftmost column. As A has rank n , there is a non-zero entry on column 1. Pick the one with minimal valuation and swap rows, so that the entry on column 0 with minimal valuation is on position $(0, 0)$. Let the new matrix be B .

Then, for each row $i \geq 1$, subtract $\frac{b_{i0}}{b_{00}} \times (\text{row } 0)$ from row i . After all of these operations, the matrix has block form:

$$\left[\begin{array}{c|c} b_{00} & * \\ \hline 0 & A' \end{array} \right]$$

where $*$ denotes some $1 \times (n-1)$ matrix, and A' is an $(m-1) \times (n-1)$ matrix. Observe that, as A had rank n and the elementary row operations don't change the rank, A' will have rank $n-1$.

Now, we can inductively apply the same procedure to A' . Observe that all row operations on A' extend to row operations on the whole matrix that don't change the block structure (as the corresponding entries in the first column are all 0's). By construction, the end result is an upper-triangular matrix, which has the same rank as the initial matrix A . \square

A.3. Implementation. We focus on two fields L : the totally ramified extension $\mathbb{Q}_p(p^{1/d})$, and the unramified extension of degree d , where we take the prime p , the degree d , and the cutoff N as input parameters.

Fix $a \in \{0, 1, \dots, q-2\}$. First, we compute the matrices $(\tau^{(a)})_{0 \leq i \leq j < N/(q-1)}$ following the method discussed in Section A.2.2. Then, for $s = 0, \dots, N/(q-1) - 1$, we will appeal to the following result to inductively compute a basis $(g_b^{(a),s})_{0 \leq b \leq s}$ for the o_L -span of $\{\tau_{i,j}^{(a)} : 0 \leq i \leq j \leq s\}$, with each $g_b^{(a),s}$ having degree b .

Proposition A.3.1. Fix $s \geq 0$, and let $(g_b^{(a),s-1})_{0 \leq b \leq s-1}$ be a basis for the o_L -span of $\{\tau_{i,j}^{(a)} : 0 \leq i \leq j \leq s-1\}$ such that each $g_b^{(a),s-1}$ has degree b .

Record the coefficients of these polynomials $g_*^{(a),s-1}$ in s row vectors, and append $s+1$ new row vectors obtained from the coefficients of $\tau_{*,s}^{(a)}$ to obtain the $(2s+1) \times (s+1)$ matrix

$$B := \begin{pmatrix} Y^s & Y^{s-1} & & 1 \\ \bullet & * & \cdots & * \\ & \bullet & \cdots & * \\ & & \ddots & \vdots \\ & & & \bullet \\ * & * & \cdots & * \\ \vdots & \vdots & & \vdots \\ * & * & \cdots & * \end{pmatrix} \begin{matrix} \tau_{s,s}^{(a)} \\ g_{s-1}^{(a),s-1} \\ \\ g_0^{(a),s-1} \\ \tau_{0,s}^{(a)} \\ \\ \tau_{s-1,s}^{(a)} \end{matrix}$$

with coefficients in L . The \bullet 's are non-zero (where $B_{s,0} \neq 0$ because $\sigma_{\underline{s},\underline{s}} = Y^{\underline{s}}$ by Lemma 4.3.8 which by Equation 19 implies that $\tau_{s,s}^{(a)} = Y^s$), so B has rank $s+1$.

Bring the full-rank matrix B to upper-triangular form B' using Gaussian elimination over the discrete valuation ring \mathcal{O}_L as per Lemma A.2.8. Then

- (i) we can define the new polynomials $g_s^{(a),s}, g_{s-1}^{(a),s}, \dots, g_0^{(a),s}$ by reading off the first $s+1$ rows of B' , so that each $g_b^{(a),s}$ has degree b and $(g_b^{(a),s})_{0 \leq b \leq s}$ form a basis for the \mathcal{O}_L -span of $\{\tau_{i,j}^{(a)} : 0 \leq i \leq j \leq s\}$;
- (ii) for each $b = 0, \dots, s-1$, the π -adic valuation of the leading coefficient in the new polynomial $g_b^{(a),s}$ is at most that of the old polynomial $g_b^{(a),s-1}$.

Proof. By Lemma A.2.8 the upper-triangular matrix B' still has rank $s+1$, so it has only non-zero elements on its main diagonal. Hence for each $b = 0, 1, \dots, s$, the polynomial $g_b^{(a),s}$ obtained by reading off the b -th row has degree b . Then of course these polynomials are linearly independent. Also they are the only non-zero rows in B' , so by Lemma A.2.7 their \mathcal{O}_L -span is the same as that of the rows of B , which by construction is precisely the \mathcal{O}_L -span of $\{\tau_{i,j}^{(a)} : 0 \leq 1 \leq j \leq s\}$, giving (i).

Now fix $0 \leq b \leq s-1$, and consider what happens to the b -th column when we reduce B to B' . Observe that in the proof of Lemma A.2.8, when we operate on the j -th column for $j = 0, \dots, s-b-1$, as the row for $g_b^{(a),s-1}$ has a 0 entry in the j -th column, it is neither chosen to be the pivot row nor altered as we subtract off multiples of the pivot row. Thus when we operate on the $(s-b)$ -th column to determine the $(s-b)$ -th row and column of B' , the leading coefficient of $g_b^{(a),s-1}$ must be a candidate for the pivot. But the pivot $B'_{s-b,s-b}$ is chosen to have minimal valuation, so $\nu_\pi(\gamma_b(g_b^{(a),s-1})) \geq \nu_\pi(B'_{s-b,s-b})$. Now $B'_{s-b,s-b} = \gamma_b(g_b^{(a),s})$ by definition, giving (ii). \square

For b fixed, it follows that $\nu_\pi(\gamma_b(g_b^{(a),s}))$, $s = b, b+1, \dots$ is a non-increasing sequence. Moreover, as $g_b^{(a),s} \in S^{(a)}$ can be written as an \mathcal{O}_L -linear combination of the $f_i^{(a)}$'s and each $f_i^{(a)}$ is of degree i , we must have $g_b^{(a),s} = \sum_{0 \leq i \leq b} \lambda_i f_i^{(a)}$ for some $\lambda_i \in \mathcal{O}_L$; by looking at the leading coefficient, it follows that

$$\nu_\pi(\gamma_b(g_b^{(a),s})) \geq \nu_\pi(\gamma_b(f_b^{(a)})) \geq -w_q(a+b(q-1)).$$

These observations motivate us to look at the following

Definition A.3.2. For $n = a + b(q-1)$, let $s_0(n)$ be the minimal $s \geq b$ such that $(g_b^{(a),s})_{0 \leq b \leq s}$ satisfies $\nu_\pi(\gamma_b(g_b^{(a),s})) = -w_q(n)$, if such s exists; otherwise set $s_0(n) = \infty$.

Then whenever $s \geq s_0(n)$ in the computations, we can immediately conclude that the equality $\nu_\pi(\gamma_b(f_b^{(a)})) = -w_q(a+b(q-1))$ in Lemma A.2.3 holds for this $n = a + b(q-1)$.

We may thus make a small optimisation: at any stage s , if $s \geq s_0(a+b(q-1))$ for all $0 \leq b < d$ then we can just drop the last d columns when carrying out Gaussian elimination. Indeed for all $s' > s$ it is unnecessary to compute $(g_b^{(a),s'})_{0 \leq b < d}$ as the π -adic valuation of each leading term has already hit the desired minimum, and to compute the leading terms of $(g_b^{(a),s'})_{d \leq b \leq s'}$ we do not need the lower-order terms in the last d columns.




FIGURE 1. `extension = "3,2,800,ram"` — $s_0(n)$ in the quadratic ramified extension $\mathbb{Q}_3(\sqrt{3})$ for $n < 800$. Red points are the n 's for which $s_0(n) \geq 800$.

A.4. **Data.** For reference, the computations in Figure 1 took

- 227.04 seconds for D ;
- 616.45 seconds for $\tau^{(0)}$ and 616.43 seconds for $\tau^{(1)}$;
- 0.20 seconds for $s = 50$, 1.89 seconds for $s = 100$, 6.15 seconds for $s = 150$, 12.09 seconds for $s = 200$, etc. for $a = 0$, and slightly less for $a = 1$.

We see that $s_0(n) - b$ seems to depend on the p -adic digits of n ; we only managed to prove a special case of this pattern, which we will discuss below. Nonetheless, the data do suggest that $s_0(n)$ is finite for every n and hence that $\text{Int}(o_L, o_L)$ is spanned by the $\sigma_{i,j}$'s as an o_L -module.

A similar pattern emerges for larger p and unramified extensions: see Figures 2 and 3 below.

More data and plots can be found at our GitHub repository <https://github.com/Team-Konstantin/Bounded-Functions-on-Character-Varieties/tree/writeup>.




FIGURE 2. `extension = "17,2,3216,ram"` — $s_0(n)$ in the quadratic ramified extension $\mathbb{Q}_{17}(\sqrt{17})$ for $n < 3216$. Note that red points are the n 's for which $s_0(n) \geq 3216$ — not enough computation was done to unveil the pattern for the larger n 's!




FIGURE 3. `extension = "5,3,12524,unram"` — $s_0(n)$ in the cubic unramified extension of \mathbb{Q}_5 for $n < 12524$. Again, note how the red points — the n 's for which $s_0(n) \geq 12524$ — give the illusion of $s_0(n) - b$ decreasing.

A.5. Some results.

Definition A.5.1. Given a natural number n , let $s_q(n)$ be the sum of digits of n in base q .

Recall Definition A.3.2:

Definition. For $n = a + b(q - 1)$, let $s_0(n)$ be the minimal $s \geq b$ such that $(g_b^{(a),s})_{0 \leq b \leq s}$ satisfies $\nu_\pi(\gamma_b(g_b^{(a),s})) = -w_q(n)$, if such s exists; otherwise set $s_0(n) = \infty$.

We define the following more intuitive quantity:

Definition A.5.2. For $n = a + b(q - 1)$, let $\text{Cap}(n) = a + bs_0(n)$. Alternatively, $\text{Cap}(n)$ is the minimal $N \geq n$ such that the o_L -span of $\{\sigma_{i,j} : 0 \leq i \leq j \leq N\}$ contains a polynomial of degree n and π -valuation of the leading term $-w_q(n)$.

Here, the equivalence of the two definitions follows from the definition of $s_0(n)$.

Let $n = a + b(q - 1)$. Analysing the computational results, we are led to believe that, if $s_q(n) < p$, then $s_0(n) = b$. This is made clear by the following:

Theorem A.5.3. *Let n be a positive integer such that $s_q(n) < p$. Let $j = n$ and $i = s_q(n)$. Then $\sigma_{i,j}$ is a polynomial of degree n , with π -valuation of leading term equal to $-w_q(n)$.*

Recall the definition of the polynomials $c_n(Y)$ from [31]:

$$[Y](t) = \sum_{n=1}^{\infty} c_n(Y) t^n$$

Translating the definition of the polynomials $\sigma_{i,j}(Y)$ and using Lemma 4.3.8, we get:

$$([Y](t))^i = \left(\sum_{n=1}^{\infty} c_n(Y) t^n \right)^i = \sum_{j=i}^{\infty} \sigma_{i,j}(Y) t^j.$$

Using the binomial theorem, this gives:

$$\sigma_{i,j} = \sum_{n_1+n_2+\dots+n_i=j} c_{n_1} c_{n_2} \dots c_{n_i}$$

Of course, for $i = 1$ we obtain $\sigma_{1,j} = c_j$. So, the proof of the Theorem 3.1 in [31] shows that $\text{Cap}(n) = n$ for n equal to some power of q . We will extend this result to all n that have $s_q(n) < p$, where $s_q(n)$ is the sum of digits of n , written in base q . For this, we need the following lemma:

Lemma A.5.4. *Let n_1, n_2, \dots, n_i be positive integers. Then, $w_q(n_1) + w_q(n_2) + \dots + w_q(n_i) \leq w_q(n_1 + n_2 + \dots + n_i)$. Equality holds if and only if $s_q(n_1) + s_q(n_2) + \dots + s_q(n_i) = s_q(n_1 + n_2 + \dots + n_i)$, that is, if there is "no carrying" in the sum $n_1 + n_2 + \dots + n_i$.*

Proof. Direct calculations show that

$$w_q(n) = \frac{n - s_q(n)}{q - 1}$$

Substituting into our inequality, we need to prove

$$s_q(n_1) + s_q(n_2) + \dots + s_q(n_i) \geq s_q(n_1 + n_2 + \dots + n_i)$$

which can be checked by direct calculations or by induction. Equality holds in the initial inequality if and only if it holds here, which is to say there is "no carrying" in the sum $n_1 + n_2 + \dots + n_i$. \square

Now, we are ready for:

Proof of Theorem A.5.3. Recall that

$$\sigma_{i,j} = \sum_{n_1+n_2+\dots+n_i=j} c_{n_1} c_{n_2} \dots c_{n_i}$$

where each c_k is a polynomial of degree at most k , with π -valuation of the leading term at least $-w_q(n)$ (as it is in $\text{Int}(o_L, o_L)$).

Let's look at each of the terms $c_{n_1} c_{n_2} \dots c_{n_i}$. As each c_k has degree at most k , this contributes to the coefficient of Y^k in $\sigma_{i,j}$ if and only if $\deg(c_{n_1}) = n_1, \deg(c_{n_2}) = n_2, \dots, \deg(c_{n_i}) = n_i$. For the moment, assume this is the case. Then, the coefficient of Y^n in this product is the product of leading coefficients of the c_{n_i} 's, which has π -valuation at least $-(w_q(n_1) + w_q(n_2) + \dots + w_q(n_i))$. Now, using Lemma A.5.4, this is at least $-w_q(n_1 + n_2 + \dots + n_i) = -w_q(n)$, with equality if and only if $s_q(n_1) + s_q(n_2) + \dots + s_q(n_i) = s_q(n) = i$, so the n_i 's are powers of q . That is, the only contribution to the coefficient of Y^n in $\sigma_{i,j}$ that has small enough valuation comes from permutations of the unique way of writing n as a sum of i powers of q . In other words, if $n = b_r b_{r-1} \dots b_1 b_{0(q)}$ is the writing of n in base q , then the only terms that have a possible contribution are obtained when (n_1, n_2, \dots, n_i) is a permutation of $(q^0, q^0, \dots, q^1, \dots, q^r)$, where each q^k appears b_k times.

But, by [31], when k is a power of q , c_k is a polynomial of degree exactly k , with π -valuation of leading term exactly $-w_q(k)$. So, when (n_1, n_2, \dots, n_i) is a permutation as above, the product $c_{n_1} c_{n_2} \dots c_{n_i}$ is a polynomial of degree n , with π -valuation of leading term equal to $-w_q(n)$. Moreover, as proved before, if (n_1, n_2, \dots, n_i) is not such a permutation, the product $c_{n_1} c_{n_2} \dots c_{n_i}$ has the coefficient of Y^n either 0 or of π -valuation larger than $-w_q(n)$.

As there are $\binom{i}{b_0, b_1, \dots, b_r}$ such permutations, with $p \nmid \binom{i}{b_0, b_1, \dots, b_r}$ (because $i < p$ by the initial assumption on n), the final sum $\sigma_{i,j}$ has degree n , with π -valuation of leading term $-w_q(n)$. \square

Definition A.5.2 then gives:

Corollary A.5.5. *Let n be a positive integer such that $s_q(n) < p$. Then $\text{Cap}(n) = n$.*

A.6. SageMath Code. (tested on Sage 9.4)

```
1 extension = "3,2,100,ram" # Choose the extension to compute with
2 precision = 1000          # Choose the precision that Sage will use
3
4 parse = extension.split(',')

```

```

5 p = int(parse[0]) # Prime to calculate with
6 d = int(parse[1]) # Degree to calculate with
7 N = int(parse[2]) # Cutoff; must be divisible by q-1
8 ram = parse[3]
9
10
11 # Python imports
12 from time import process_time
13 import matplotlib.pyplot as plt
14 import numpy as np
15
16 # Definitions
17 from sage.rings.padic.padic_generic import ResidueLiftingMap
18 from sage.rings.padic.padic_generic import ResidueReductionMap
19 import sage.rings.padic.padic_extension_generic
20
21 power = p^d - 1
22 t_poly = ""
23
24 if ram == "ram":
25     t_poly = f"x^{d}-{p}"
26 else:
27     # generate poly for unramified case
28     Fp = GF(p)
29     Fp_t.<t> = PolynomialRing(Fp)
30     unity_poly = t^(power) - 1
31     factored = unity_poly.factor()
32     factored_str = str(factored)
33     start = factored_str.find("^"+str(d))
34     last_brac_pos = factored_str.find(")",start)
35     first_brac_pos = len(factored_str) \
36         - factored_str[::-1].find("(",len(factored_str)-start)
37     t_poly = factored_str[first_brac_pos:last_brac_pos].replace('t','x')
38
39
40 # Define the polynomial to adjoin a root from
41 Q_p = Qp(p,precision)
42 R_Qp.<x> = PolynomialRing(Q_p)
43 f_poly = R_Qp(t_poly)
44
45 # Define the p-adic field, its ring of integers and its residue field
46 # These dummy objects are a workaround to force the precision wanted
47 dummy1.<y> = Zp(p).ext(f_poly)
48 dummy2.<y> = Qp(p).ext(f_poly)
49
50 o_L.<y> = dummy1.change(prec=precision)
51 L.<y> = dummy2.change(prec=precision)
52 k_L = L.residue_field()
53 print(L)

```

```

54
55 # Find the generator of the unique maximal ideal in o_L.
56 Pi = o_L.uniformizer()
57
58 # Find f, e and q
59 f = k_L.degree()          # The degree of the residual field extension
60 e = L.degree()/k_L.degree() # The ramification index
61 q = p^f
62
63 # Do linear algebra over the ring of polynomials L[X]
64 # in one variable X with coefficients in the field L:
65 L_X.<X> = L[]
66 L_Y.<Y> = L[]
67
68 v = L.valuation()
69
70 # The subroutine Dmatrix calculates the following sparse matrix of coefficients.
71 # Let D[k,n] be equal to k! times the coefficient of Y^k in the polynomial P_n(Y).
72 # I compute this using the useful and easy recursion formula
73 #      
$$D[k,n] = \sum_{r \geq 0} \pi^{-r} D[k-1,n-q^r]$$

74 # that can be derived from Proposition 4.3.2.
75 # The algorithm is as follows: first make a zero matrix with S rows and columns
76 # (roughly, S is (q-1)*Size), then quickly populate it one row at a time,
77 # using the recursion formula.
78 def Dmatrix(S):
79     D = matrix(L, S,S)
80     D[0,0] = 1
81     for k in range(1,S):
82         for n in range(k,S):
83             r = 0
84             while n >= q^r:
85                 D[k,n] = D[k,n] + D[k-1,n-q^r]/Pi^r # the actual recursion
86             r = r+1
87     return D
88
89
90 # \Tau^{(m)} in Definition 4.3.11:
91 def TauMatrix(Size, m, D=None):
92     if D is None:
93         D = Dmatrix((q - 1) * (Size + 1))
94     R = matrix(L, Size,Size, lambda x,y: D[m + (q-1)*x, m + (q-1)*y])
95
96     # Define a diagonal matrix:
97     Diag = matrix(L_X, Size,Size, lambda x,y: kronecker_delta(x,y) * X^x)
98
99     # Compute the inverse of R:
100     S = R.inverse()
101
102     # Compute the matrix Tau using Lemma 4.3.12:

```

```

103     Tau = S * Diag * R
104
105     return Tau
106
107 def underscore(m, i):
108     return m + i*(q-1)
109
110 def w_q(n):
111     return (n - sum(n.digits(base=q))) / (q-1)
112
113 def compute_s(N, filename=None):
114     assert N%(q-1) == 0
115
116     t_start = process_time()
117     D = Dmatrix(N)
118     t_end = process_time()
119     print(f"D matrix: {t_end-t_start : .2f} sec")
120
121     s0_s = [-1 for _ in range(N)]
122
123     for a in range(q-1):
124         t_start = process_time()
125         Tau_a = TauMatrix(N//(q-1), a, D)
126         t_end = process_time()
127         print(f"a={a}, Tau matrix: {t_end-t_start : .2f} sec")
128
129         B_old = Matrix(0,0)
130         d = 0
131         for s in range(N // (q-1)):
132             t_start = process_time()
133
134             # 1. Use the non-zero rows from previous calculations
135             # 2. Add a 0 column to its left
136             # 3. Add rows corresponding to entries from the j_th column of Tau_
137             B = Matrix(L, 2*s-d+1, s-d+1)
138             B[0,0] = 1 # Tau_a[s, s]
139             B[1:s-d+1, 1:] = B_old
140             for i in [0 .. s-1]:
141                 coeffs = Tau_a[i, s].list()
142                 B[s-d+1+i, B.ncols()-len(coeffs)+d:] = vector(L, reversed(coeffs))
143
144             # Perform Gaussian elimination
145             i0 = 0
146             ks = []
147             for k in range(B.ncols()):
148                 valuation_row_pairs = [
149                     (v(B[i,k]), i) for i in range(i0, B.nrows()) if B[i,k] != 0
150
151                 if not valuation_row_pairs:

```

```

152         raise ValueError("B is not full-rank")
153     minv, i_minv = min(valuation_row_pairs)
154     ks.append(k)
155
156     # Swap the row of minimum valuation with the first bad row
157     B[i0, :], B[i_minv, :] = B[i_minv, :], B[i0, :]
158
159     # Divide the top row by a unit in o_L
160     u = B[i0, k] / Piint(e * v(B[i0, k]))
161     B[i0, :] /= u
162
163     # Cleave through the other rows
164     for i in range(i0 + 1, B.nrows()):
165         if v(B[i, k]) >= v(B[i0, k]):
166             B[i, :] -= B[i, k]/B[i0, k] * B[i0, :]
167
168     i0 += 1
169
170     d_is_updated = False
171     for b in [d .. s]:
172         n = a + b*(q-1)
173         if v(B[s-b, s-b]) * e == -w_q(n):
174             if s0_s[n] == -1:
175                 s0_s[n] = s
176             else:
177                 if not d_is_updated:
178                     d = b
179                     d_is_updated = True
180     B_old = B[:s-d+1, :s-d+1]
181
182     t_end = process_time()
183     print(f"a={a}, s={s}: {t_end-t_start : .2f} sec", end='\r')
184     if filename is not None:
185         with open(filename, 'w') as f:
186             f.write("n,s0\n")
187             for n, s0 in enumerate(s0_s):
188                 f.write(f"{n},{s0}\n")
189     print()
190
191     plt.style.use('bmh')
192     fig = plt.figure(figsize=(15,6), dpi=300)
193     for n, s0 in enumerate(s0_s):
194         if s0 != -1:
195             b = n // (q-1)
196             plt.plot(n, s0-b, 'x', c='C0')
197         else:
198             plt.plot(n, 0, 'x', c='C1')
199     plt.xlabel(r"$n = a + b(q-1)$")
200     plt.ylabel(r"$s_0(n) - b$")

```



```

201     plt.title(str(L))
202     plt.minorticks_on()
203     plt.grid(which='both')
204     plt.grid(which='major', linestyle='-', c='grey')
205
206     return s0_s, fig
207
208
209 s0_s = compute_s(N);

```

REFERENCES

- [1] K. Ardakov, *Prime ideals in nilpotent Iwasawa algebras*, Invent. Math. **190** (2012), no. 2, 439–503.
- [2] K. Ardakov and K. A. Brown, *Primeness, semiprimeness and localisation in Iwasawa algebras*, Trans. Amer. Math. Soc. **359** (2007), no. 4, 1499–1515.
- [3] P. N. Balister and S. Howson, *Note on Nakayama’s Lemma for compact Λ -modules*, Asian J. Math. **1** (1997), no. 2, 224–229.
- [4] L. Berger and P. Colmez, *Théorie de Sen et vecteurs localement analytiques*, Ann. Sci. Éc. Norm. Supér. (4) **49** (2016), no. 4, 947–970.
- [5] L. Berger, P. Schneider, and B. Xie, *Rigid character groups, Lubin-Tate theory, and (φ, Γ) -modules*, Mem. Amer. Math. Soc. **263** (2020), no. 1275, v+79.
- [6] M. Bhargava, *P -orderings and polynomial functions on arbitrary subsets of Dedekind rings*, J. Reine Angew. Math. **490** (1997), 101–127.
- [7] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis*, Grundlehren der mathematischen Wissenschaften, vol. 261, Springer-Verlag, Berlin, 1984.
- [8] N. Bourbaki, *Commutative algebra. Chapters 1–7*, Elements of Mathematics (Berlin), Springer-Verlag, Berlin, 1998.
- [9] O. Brinon and B. Conrad, *CM Summer school notes on p -adic Hodge theory*, 2009.
- [10] R. F. Coleman, *Division values in local fields*, Invent. Math. **53** (1979), no. 2, 91–116.
- [11] P. Colmez, *Représentations localement analytiques de $\mathbf{GL}_2(\mathbf{Q}_p)$ et (φ, Γ) -modules*, Represent. Theory **20** (2016), 187–248.
- [12] M. Demazure and P. Gabriel, *Introduction to algebraic geometry and algebraic groups*, North-Holland Mathematics Studies, vol. 39, North-Holland Publishing Co., Amsterdam-New York, 1980, Translated from the French by J. Bell.
- [13] M. Emerton, *Locally analytic vectors in representations of locally p -adic analytic groups*, Mem. Amer. Math. Soc. **248** (2017), no. 1175, iv+158.
- [14] L. Fourquaux, *Logarithme de Perrin-Riou pour des extensions associées à un groupe de Lubin-Tate*, preprint, 2008.
- [15] L. Fourquaux and B. Xie, *Triangulable \mathcal{O}_F -analytic (φ_q, Γ) -modules of rank 2*, Algebra Number Theory **7** (2013), no. 10, 2545–2592.
- [16] L. Gruson, *Fibrés vectoriels sur un polydisque ultramétrique*, Ann. Sci. École Norm. Sup. (4) **1** (1968), 45–89.
- [17] M. Hazewinkel, *Formal groups and applications*, AMS Chelsea Publishing, Providence, RI, 2012, Corrected reprint of the 1978 original.
- [18] M. Hopkins, *Lectures on Lubin-Tate spaces*, Arizona Winter School, 2019.
- [19] N. M. Katz, *Formal groups and p -adic interpolation*, Journées Arithmétiques de Caen (Univ. Caen, Caen, 1976), Astérisque No. 41–42, Soc. Math. France, Paris, 1977, pp. 55–65.
- [20] N. M. Katz, *Divisibilities, congruences, and Cartier duality*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **28** (1981), no. 3, 667–678 (1982).

- [21] S. Lang, *Cyclotomic fields I and II*, second ed., Graduate Texts in Mathematics, vol. 121, Springer-Verlag, New York, 1990, With an appendix by Karl Rubin.
- [22] H. Li and F. van Oystaeyen, *Zariskian filtrations*, *K-Monographs in Mathematics*, vol. 2, Kluwer Academic Publishers, Dordrecht, 1996.
- [23] J. Lubin and J. Tate, *Formal complex multiplication in local fields*, *Ann. of Math. (2)* **81** (1965), 380–387.
- [24] C. Perez-Garcia and W. H. Schikhof, *Locally convex spaces over non-Archimedean valued fields*, *Cambridge Studies in Advanced Mathematics*, vol. 119, Cambridge University Press, Cambridge, 2010.
- [25] B. Perrin-Riou, *Théorie d’Iwasawa p -adique locale et globale*, *Invent. Math.* **99** (1990), no. 2, 247–292.
- [26] T. Schmidt, *Picard groups in p -adic Fourier theory*, *Manuscripta Math.* **144** (2014), no. 1-2, 1–23.
- [27] P. Schneider, *Nonarchimedean functional analysis*, *Springer Monographs in Mathematics*, Springer-Verlag, Berlin, 2002.
- [28] P. Schneider and J. Teitelbaum, *p -adic Fourier theory*, *Doc. Math.* **6** (2001), 447–481.
- [29] P. Schneider and J. Teitelbaum, *Locally analytic distributions and p -adic representation theory, with applications to GL_2* , *J. Amer. Math. Soc.* **15** (2002), no. 2, 443–468.
- [30] E. de Shalit, *Mahler bases and elementary p -adic analysis*, *J. Théor. Nombres Bordeaux* **28** (2016), no. 3, 597–620.
- [31] E. de Shalit and E. Iceland, *Integer valued polynomials and Lubin-Tate formal groups*, *J. Number Theory* **129** (2009), no. 3, 632–639.
- [32] J. T. Tate, *p -divisible groups*, *Proc. Conf. Local Fields (Driebergen, 1966)*, Springer, Berlin, 1967, pp. 158–183.
- [33] L. Thomas, *On the Galois module structure of extensions of local fields*, *Actes de la Conférence “Fonctions L et Arithmétique”*, *Publ. Math. Besançon Algèbre Théorie Nr.*, vol. 2010, Lab. Math. Besançon, Besançon, 2010, pp. 157–194.

Konstantin Ardakov
 Mathematical Institute
 University of Oxford
 E-mail: ardakov@maths.ox.ac.uk
 URL: <http://people.maths.ox.ac.uk/ardakov/>

Laurent Berger
 UMPA, ENS de Lyon
 UMR 5669 du CNRS
 E-mail: laurent.berger@ens-lyon.fr
 URL: <https://perso.ens-lyon.fr/laurent.berger/>