

### More on Mayer-Vietoris

#### Exercise 1. Computations of homology groups

Consider the following topological spaces:

- (a) The 2-dimensional torus.
- (b) The Klein bottle.
- (c) The orientable surface of genus 2.

Compute their homology groups using the Mayer-Vietoris exact sequence. Generalize the result of (c) to the orientable surface of genus  $g$  using cellular homology.

#### Exercise 2. Homology of the surface of genus $g$ using Mayer-Vietoris

1. Let  $X_g$  be the complement in the 2-sphere of  $2g$  disjoint open discs  $D_1, \dots, D_{2g}$ . Compute the homology groups of  $X_g$ .
2. Compute the map in homology induced by the inclusion:

$$\bigsqcup_{i=1}^{2g} \partial D_i \rightarrow X_g.$$

3. Deduce the homology groups of the surface of genus  $g$ .

#### Exercise 3. Fundamental class of the sphere and higher Hurewicz maps

Let  $n \geq 1$  be an integer. A *fundamental class* of the sphere  $S^n$  is a generator of the group  $H_n(S^n)$ .

1. Let  $(U, V)$  be a topological cover of a space  $X$ . Give a method to compute the connecting map:

$$\partial : H_n(X) \rightarrow H_{n-1}(U \cap V)$$

of the Mayer-Vietoris exact sequence.

2. Using the first question, find a fundamental class of the sphere  $S^n$ .
3. Let  $(X, x)$  be a pointed topological space, construct a group homomorphism

$$\pi_n(X, x) \rightarrow H_n(X)$$

*Hint:* map  $f : S^n \rightarrow X$  to the pushforward by  $f$  of a fundamental class of  $S^n$ .

4. Show that this map is surjective when  $X = S^n$ .

#### Exercise 4. The Perron-Frobenius theorem

Use Brouwer's fixed point theorem to show that a square matrix with nonnegative entries has a real nonnegative eigenvalue.

#### Exercise 5. Homology of a product of spheres

Let  $m, n \geq 1$  be integers. Compute the homology groups of  $S^n \times S^m$  using

1. the Mayer-Vietoris exact sequence.
2. the cellular homology theorem.