

The slide features a decorative border composed of various elements: a dense arrangement of orange coins with skull and crossbones symbols, green coins, and whole oranges with green leaves. These elements are scattered around the edges of the slide, creating a vibrant and thematic frame.

Optimal disc and sphere packings

Daria Pchelina
LIP, équipe MC2

SIESTE seminar
16/10/2024



1 Introduction

2 Disc packings

3 Sphere packings

4 Conclusion



1 Introduction

2 Disc packings

3 Sphere packings

4 Conclusion

Optimal coin packings

Given infinite number of identical coins $\left(\textcircled{\text{☠}}\right)$,
how to place them on an infinite plane without overlap to maximize the covered surface?


coin packing:



Optimal coin packings

Given infinite number of identical coins $\left(\textcircled{\text{skull}} \right)$,
 how to place them on an infinite plane without overlap to maximize the covered surface?

coin packing:




Optimal coin packings

Given infinite number of identical coins $\left(\textcircled{\text{skull}}$



), how to place them on an infinite plane without overlap to maximize the covered surface?




hexagonal coin packing:



1910–1940

The hexagonal coin packing is optimal.

Introduction



Introduction



(proved optimal in 2000-2022)



Kepler conjecture, 1611

The "cannonball" packing is optimal:



(proved in 1998-2014)

Introduction




(proved optimal in 2000-2022)

Kepler conjecture, 1611

The "cannonball" packing is optimal:





(proved in 1998-2014)



$\mathbb{R}^8, \mathbb{R}^{24}$
(Viazovska, Fields Medal 2022)

Introduction




(proved optimal in 2000-2022)

Kepler conjecture, 1611

The "cannonball" packing is optimal:




(proved in 1998-2014)



$\mathbb{R}^8, \mathbb{R}^{24}$
(Viazovska, Fields Medal 2022)


combine different types of nanoparticles
self-assembly

new material



Paik et al 2015


phase separation



Cheon et al 2006


combine different types of nanoparticles
self-assembly

new material




Paik et al 2015

phase separation



Cheon et al 2006

Also in 3D:



Wu, Fan, Yin 2022

Chemists' question : **which sizes and concentrations allow for new materials?**



1 Introduction



2 Disc packings

3 Sphere packings

4 Conclusion



Definitions

Discs:

Packing P :
(in \mathbb{R}^2)

Definitions

Discs:



Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Definitions

Discs:



Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Definitions

Discs:



Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Definitions

Discs:



Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Definitions

Discs:



Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Definitions

Discs:



Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Definitions


Discs:

Packing P :
(in \mathbb{R}^2)

Density:

$$\delta(P) := \limsup_{n \rightarrow \infty} \frac{\text{area}([-n, n]^2 \cap P)}{\text{area}([-n, n]^2)}$$

Main Question


Given a finite set of discs (e.g., ,
what is the maximal density δ^* of a packing?

$$\delta^* := \sup_P \delta(P)$$


Optimal 2-disc packings

Theorem (Heppes 2000, 2003, Kennedy 2005, Bedaride and Fernique 2022)


Each of the following packings is optimal (densest) for discs of radii 1 and r :




$r \approx 0.63$ $\delta^* \approx 91.1\%$




$r \approx 0.54$ $\delta^* \approx 91.1\%$




$r \approx 0.53$ $\delta^* \approx 91.4\%$




$r \approx 0.41$ $\delta^* \approx 92\%$




$r \approx 0.38$ $\delta^* \approx 92\%$




$r \approx 0.34$ $\delta^* \approx 92.5\%$



$r \approx 0.28$ $\delta^* \approx 93.2\%$



$r \approx 0.15$ $\delta^* \approx 95\%$




$r \approx 0.1$ $\delta^* \approx 96\%$


Optimal 2-disc packings

Theorem (Heppes 2000, 2003, Kennedy 2005, Bedaride and Fernique 2022)


Each of the following packings is optimal (densest) for discs of radii 1 and r :




$r \approx 0.63$ $\delta^* \approx 91.1\%$




$r \approx 0.54$ $\delta^* \approx 91.1\%$




$r \approx 0.53$ $\delta^* \approx 91.4\%$




$r \approx 0.41$ $\delta^* \approx 92\%$




$r \approx 0.38$ $\delta^* \approx 92\%$




$r \approx 0.34$ $\delta^* \approx 92.5\%$



$r \approx 0.28$ $\delta^* \approx 93.2\%$




$r \approx 0.15$ $\delta^* \approx 95\%$



$r \approx 0.1$ $\delta^* \approx 96\%$


Connelly conjecture

Triangulated packings:




Conjecture (Connelly 2018)


If a finite set of discs allows **saturated** triangulated packings then one of them is optimal.




triangulated
saturated



non triangulated
saturated




triangulated
non saturated



non triangulated
non saturated


Connelly conjecture

Triangulated packings:




Conjecture (Connelly 2018)


If a finite set of discs allows **saturated** triangulated packings then one of them is optimal.




triangulated
saturated



non triangulated
saturated




triangulated
non saturated




non triangulated
non saturated

Theorem (●●● Fernique, Hashemi, Sizova 2019)

Discs of radii 1, r and s : there are 164 pairs (r, s) allowing triangulated packings.




Disc packings



164 (r, s) allowing triangulated packings:


- 15 cases: non saturated
- 16+16 cases: a **ternary** or **binary** triangulated packing is densest
- 45 cases: a **binary non triangulated** packing is denser




Disc packings

Theorem (Fernique, P 2023)


Each of the following packings is optimal for discs of radii 1, r and s :



1-disc packing




multi-size disc packing




Voronoi cell of a disc in a packing: set of points closer to this disc than to any other

FM-triangulation

1-disc packing



multi-size disc packing




Voronoi cell of a disc in a packing: set of points closer to this disc than to any other


Voronoi diagram of a packing: partition of the plane into Voronoi cells

FM-triangulation

1-disc packing



multi-size disc packing




Voronoi cell of a disc in a packing: set of points closer to this disc than to any other

Voronoi diagram of a packing: partition of the plane into Voronoi cells


FM-triangulation of a packing: dual graph of the Voronoi diagram

FM-triangulation

1-disc packing



multi-size disc packing




Voronoi cell of a disc in a packing: set of points closer to this disc than to any other

Voronoi diagram of a packing: partition of the plane into Voronoi cells


FM-triangulation of a packing: dual graph of the Voronoi diagram

FM-triangulation

1-disc packing



multi-size disc packing



Voronoi cell of a disc in a packing: set of points closer to this disc than to any other

Voronoi diagram of a packing: partition of the plane into Voronoi cells


FM-triangulation of a packing: dual graph of the Voronoi diagram

Density of a triangle Δ in a packing = its proportion covered by discs


$$\delta_{\Delta} = \frac{\text{area}(\Delta \cap P)}{\text{area}(\Delta)}$$

FM-triangulation

1-disc packing



multi-size disc packing



Voronoi cell of a disc in a packing: set of points closer to this disc than to any other


Voronoi diagram of a packing: partition of the plane into Voronoi cells

FM-triangulation of a packing: dual graph of the Voronoi diagram


Density of a triangle Δ in a packing = its proportion covered by discs

$$\delta_{\Delta} = \frac{\text{area}(\Delta \cap P)}{\text{area}(\Delta)}$$

Local density redistribution




P of density $\delta(P)$




P^* of density δ^*

Local density redistribution



P of density $\delta(P)$


$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$



P^* of density δ^*

$$\delta(\triangle) = \delta^*$$


Local density redistribution



P of density $\delta(P)$

$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$


$$\delta(P) \leq \delta^*$$



P^* of density δ^*

$$\delta(\triangle) = \delta^*$$


Local density redistribution



P of density $\delta(P)$

$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$

$$\delta(P) \leq \delta^*$$



P^* of density δ^*


$$\delta(\triangle) = \delta^*$$

Proof:

- the smallest angle of any Δ is at least $\frac{\pi}{6}$
- thus the largest angle is between $\frac{\pi}{3}$ and $\frac{2\pi}{3}$
- density of a triangle Δ : $\delta(\Delta) = \frac{\pi/2}{\text{area}(\Delta)}$
- the area of a triangle ABC with the largest angle \hat{A} : $\frac{|AB| \cdot |AC| \cdot \sin \hat{A}}{2} \geq \frac{2 \cdot 2 \cdot \frac{\sqrt{3}}{2}}{2} = \sqrt{3}$
- thus the density of ABC is less or equal to $\frac{\pi/2}{\sqrt{3}} = \delta^*$

$$2 > R = \frac{|AB|}{2 \sin \hat{C}} \geq \frac{1}{\sin \hat{C}} \implies \hat{C} > \frac{\pi}{6}$$


Local density redistribution



P of density $\delta(P)$


$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$

$$\delta(P) \leq \delta^*$$




P^* of density δ^*

$$\delta(\triangle) = \delta^*$$




P of density $\delta(P)$



P^* of density δ^*


Local density redistribution



P of density $\delta(P)$


$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$

$$\delta(P) \leq \delta^*$$



P^* of density δ^*

$$\delta(\triangle) = \delta^*$$




P of density $\delta(P)$

Triangles in P^* have different densities:


$$\delta(\triangle) < \delta^* < \delta(\triangle)$$

Hopeless to bound the density by δ^* in each triangle...




P^* of density δ^*

Local density redistribution




P of density $\delta(P)$

$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$



P^* of density δ^*


$$\delta(\triangle) = \delta^*$$



P of density $\delta(P) \leq \delta'(P)$


redistributed density δ' :

dense triangles
share their density
with neighbors



P^* of density δ^*


Local density redistribution



P of density $\delta(P)$


$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$

$$\delta(P) \leq \delta^*$$



P^* of density δ^*


$$\delta(\triangle) = \delta^*$$



P of density $\delta(P) \leq \delta'(P)$


redistributed density δ' :

dense triangles
share their density
with neighbors



P^* of density δ^*


Local density redistribution



P of density $\delta(P)$


$$\forall \Delta, \delta(\Delta) \leq \delta(\triangle) = \delta^*$$

$$\delta(P) \leq \delta^*$$



P^* of density δ^*

$$\delta(\triangle) = \delta^*$$




P of density $\delta(P) \leq \delta'(P)$

$$\forall \Delta, \delta'(\Delta) \leq \delta^*$$

$$\delta(P) \leq \delta'(P) \leq \delta^*$$

redistributed density δ' :

dense triangles
share their density
with neighbors



P^* of density δ^*


Verifying inequalities on compact sets

How to check $\delta'(\Delta) \leq \delta^*$ on each possible triangle Δ ? (there is a continuum of them)

Verifying inequalities on compact sets

How to check $\delta'(\Delta) \leq \delta^*$ on each possible triangle Δ ? (there is a continuum of them)

FM-triangulation properties + saturation \Rightarrow uniform bound on edge length




$$r_a + r_b \leq c \leq r_a + r_b + 2s$$

Verifying inequalities on compact sets

How to check $\delta'(\Delta) \leq \delta^*$ on each possible triangle Δ ? (there is a continuum of them)


FM-triangulation properties + saturation \Rightarrow uniform bound on edge length




$$r_a + r_b \leq c \leq r_a + r_b + 2s$$

- Interval arithmetic: to verify $\delta'(\Delta_{a,b,c}) \leq \delta^*$ for all $(a, b, c) \in [\underline{a}, \bar{a}] \times [\underline{b}, \bar{b}] \times [\underline{c}, \bar{c}]$, we verify $[\underline{\delta}, \bar{\delta}] \leq \delta^*$ where $[\underline{\delta}, \bar{\delta}] = \delta'(\Delta_{[\underline{a}, \bar{a}], [\underline{b}, \bar{b}], [\underline{c}, \bar{c}]})$


- If $\delta^* \in [\underline{\delta}, \bar{\delta}]$, recursive subdivision:




Our proof worked for these cases:




53




54




55




56




66




76




77




79




93




108




115




116




118



129




131




146


And these:




1




2




3




4




5




b_8




6




b_4




7




8




9




b_7




10




11




12




13




14




15



b_9




19






b_6

45 counter examples: *flip-and-flow* method






When the ratio of two discs is close enough to the ratio in a dense binary packing, we can pack these discs in a similar manner (non triangulated) and still get high density


ternary triangulated
packing



counter example
using only 2 discs






45 counter examples: *flip-and-flow* method






When the ratio of two discs is close enough to the ratio in a dense binary packing, we can pack these discs in a similar manner (non triangulated) and still get high density

ternary triangulated
packing



dense binary
packing





1 Introduction


2 Disc packings

3 Sphere packings


Conclusion

Kepler conjecture: -packings

3D close -packings:




$$\delta^* = \frac{\pi}{3\sqrt{2}}$$



Kepler conjecture: ●-packings

3D close ●-packings:



$$\delta^* = \frac{\pi}{3\sqrt{2}}$$

Hales, Ferguson, 1998–2014

(Conjectured by Kepler, 1611)


Close packings maximize the density.

- close packings maximize the density among lattice packings Gauss, 1831
- 18th problem of the Hilbert's list 1900
- 6 preprints by Hales and Ferguson ArXiv 1998
> 50000 + 137000 lines of code
- reviewing: 13 reviewers, 4 years... "99% certain" 1999–2003
- "short" version of the proof Annals of Mathematics 2005
- full version: 6 edited papers DCG 2006
- Flyspeck project: formal proof (HOL Light and Isabelle) 2003–2014
Forum of Mathematics, Pi 2017

sphere




cannonball packing



rock salt spheres



rock salt packing




Rock salt  -packings

rock salt spheres



rock salt packing



Rock salt  -packings

rock salt spheres




triangulated \rightarrow simplicial
(contact graph is a "tetrahedration")

Fernique, 2019

The only simplicial 2-sphere packings in 3D are rock salt packings.

rock salt packing



Rock salt -packings

rock salt spheres



triangulated \rightarrow simplicial
(contact graph is a "tetrahedration")

Fernique, 2019


The only simplicial 2-sphere packings in 3D are rock salt packings.

Salt conjecture

open problem

Rock salt packing is optimal $\delta^* \approx 79\%$

rock salt packing



Upper density bound for $\odot\bullet$ -packings in 2D

Florian, 1960

The density of a packing never exceeds the density in the following triangle:



Florian, 1960


The density of a packing never exceeds the density in the following triangle:

**Proof:**

- Dimension reduction ($3 \rightarrow 1$)


Fejes Tóth, Mólnar, 1958

For any triangle, there is a denser triangle with at least two contacts between discs.



Upper density bound for $\odot\bullet$ -packings in 2D


Florian, 1960

The density of a packing never exceeds the density in the following triangle: **Proof:**


- Dimension reduction ($3 \rightarrow 1$)

Fejes Tóth, Mólnar, 1958

For any triangle, there is a denser triangle with at least two contacts between discs.




- Function analysis




Upper density bound for $\text{orange} \cdot \text{blue}$ -packings


FM-triangulation (triangles)




FM-simplicial partition (tetrahedra)



densest triangle




densest tetrahedra



Upper density bound for $\odot\bullet$ -packingsTheorem, $r = \sqrt{2} - 1$

in progress

Each of the following tetrahedra is densest among the tetrahedra with the same spheres:




$$\delta_{1111} \approx 0.7209$$




$$\delta_{11rr} \approx 0.8105$$



$$\delta_{1rrr} \approx 0.8065$$



$$\delta_{rrrr} \approx 0.7847$$




$$\delta_{111r} \approx 0.8125$$

Upper density bound for $\odot\bullet$ -packingsTheorem, $r = \sqrt{2} - 1$

in progress


Each of the following tetrahedra is densest among the tetrahedra with the same spheres:




$$\delta_{1111} \approx 0.7209$$




$$\delta_{11rr} \approx 0.8105$$



$$\delta_{1rrr} \approx 0.8065$$




$$\delta_{rrrr} \approx 0.7847$$



$$\delta_{111r} \approx 0.8125$$

Proof:

- Dimension reduction ($6 \rightarrow 4$):
tetrahedron deflation + sphere sliding



Upper density bound for $\textcircled{\small r} \textcircled{\small r} \textcircled{\small r} \textcircled{\small r}$ -packingsTheorem, $r = \sqrt{2} - 1$

in progress


Each of the following tetrahedra is densest among the tetrahedra with the same spheres:




$$\delta_{1111} \approx 0.7209$$




$$\delta_{11rr} \approx 0.8105$$



$$\delta_{1rrr} \approx 0.8065$$




$$\delta_{rrrr} \approx 0.7847$$



$$\delta_{111r} \approx 0.8125$$

Proof:

- Dimension reduction ($6 \rightarrow 4$):
tetrahedron deflation + sphere sliding



Upper density bound for $\textcircled{\small r} \textcircled{\small r} \textcircled{\small r} \textcircled{\small r}$ -packingsTheorem, $r = \sqrt{2} - 1$

in progress


Each of the following tetrahedra is densest among the tetrahedra with the same spheres:




$$\delta_{1111} \approx 0.7209$$




$$\delta_{11rr} \approx 0.8105$$



$$\delta_{1rrr} \approx 0.8065$$




$$\delta_{rrrr} \approx 0.7847$$



$$\delta_{111r} \approx 0.8125$$

Proof:


- Dimension reduction ($6 \rightarrow 4$):
tetrahedron deflation + sphere sliding



Upper density bound for $\textcircled{\text{orange}}\textcircled{\text{blue}}$ -packingsTheorem, $r = \sqrt{2} - 1$

in progress

Each of the following tetrahedra is densest among the tetrahedra with the same spheres:




$$\delta_{1111} \approx 0.7209$$




$$\delta_{11rr} \approx 0.8105$$



$$\delta_{1rrr} \approx 0.8065$$




$$\delta_{rrrr} \approx 0.7847$$



$$\delta_{111r} \approx 0.8125$$

Proof:

- Dimension reduction ($6 \rightarrow 4$):
tetrahedron deflation + sphere sliding
- Computer-assisted proof for tetrahedra with 2 contacts:
recursive subdivision + interval arithmetic
 ≈ 1000 lines of code



11h on 96 CPUs

Why the computations are so slow

interval arithmetic + huge formulas \rightarrow loss of precision

Why the computations are so slow

interval arithmetic + huge formulas → loss of precision

Example: to compute the support sphere radius, we need to solve $Ar^2 + Br + C = 0$

$$\begin{aligned}
 A = & -4a^2b^2d^2 + 4a^2c^2d^2 + 4b^2c^2d^2 - 4c^4d^2 - 4c^2d^4 + 4a^2b^2e^2 - 4b^4e^2 - 4a^2c^2e^2 + 4b^2c^2e^2 + 4b^2d^2e^2 + 4c^2d^2e^2 - 4b^2e^4 - 4a^4f^2 + 4a^2b^2f^2 + 4a^2c^2f^2 - 4b^2c^2f^2 + 4a^2d^2f^2 + 4c^2d^2f^2 + 4a^2e^2f^2 + 4b^2e^2f^2 - 4d^2e^2f^2 \\
 & - 4a^2f^4 + 4d^4r_a^2 - 8d^2e^2r_a^2 + 4e^4r_a^2 - 8d^2f^2r_a^2 - 8e^2f^2r_a^2 + 4f^4r_a^2 - 8b^2d^2r_xr_y + 8c^2d^2r_xr_y + 8b^2e^2r_xr_y - 8c^2e^2r_xr_y - 16a^2f^2r_xr_y + 8b^2f^2r_xr_y + 8c^2f^2r_xr_y + 8d^2f^2r_xr_y + 8e^2f^2r_xr_y - 8f^4r_xr_y + 4b^4r_y^2 - 8b^2c^2r_y^2 + 4c^4r_y^2 \\
 & - 8b^2f^2r_y^2 - 8c^2f^2r_y^2 + 4f^4r_y^2 - 8a^2d^2r_xr_z + 8c^2d^2r_xr_z + 8a^2e^2r_xr_z - 16b^2e^2r_xr_z + 8c^2e^2r_xr_z + 8d^2e^2r_xr_z - 8e^4r_xr_z + 8a^2f^2r_xr_z - 8c^2f^2r_xr_z + 8e^2f^2r_xr_z - 8a^2b^2r_yr_z + 8a^2c^2r_yr_z + 8b^2c^2r_yr_z - 8c^4r_yr_z - 16c^2d^2r_yr_z \\
 & + 8b^2e^2r_yr_z + 8c^2e^2r_yr_z + 8a^2f^2r_yr_z + 8c^2f^2r_yr_z - 8e^2f^2r_yr_z + 4a^4r_z^2 - 8a^2c^2r_z^2 + 4c^4r_z^2 - 8a^2e^2r_z^2 + 4e^4r_z^2 - 8a^2d^2r_xr_w + 8b^2d^2r_xr_w - 8d^4r_xr_w - 8d^2e^2r_xr_w + 8a^2f^2r_xr_w + 8b^2f^2r_xr_w - 8d^2f^2r_xr_w - 8a^4r_zr_w + 8a^2b^2r_zr_w + 8d^2e^2r_zr_w \\
 & + 8a^2f^2r_zr_w - 8b^2f^2r_zr_w + 8d^2f^2r_zr_w + 8a^2b^2r_yr_w - 8b^4r_yr_w - 8a^2c^2r_yr_w + 8b^2c^2r_yr_w + 8b^2d^2r_yr_w + 8c^2d^2r_yr_w - 16b^2e^2r_yr_w + 8a^2f^2r_yr_w + 8b^2f^2r_yr_w - 8d^2f^2r_yr_w - 8a^4r_zr_w + 8a^2b^2r_zr_w + 8a^2c^2r_zr_w - 8b^2c^2r_zr_w \\
 & + 8a^2d^2r_zr_w + 8c^2d^2r_zr_w + 8a^2e^2r_zr_w + 8b^2e^2r_zr_w - 8d^2e^2r_zr_w - 16a^2f^2r_zr_w + 4a^4r_w^2 - 8a^2b^2r_w^2 + 4b^4r_w^2 - 8a^2c^2r_w^2 - 8b^2d^2r_w^2 + 4d^4r_w^2 \\
 \\
 B = & -4c^2d^4r_x + 4b^2d^2e^2r_x + 4c^2d^2e^2r_x - 4b^2e^4r_x + 4a^2d^2f^2r_x + 4c^2d^2f^2r_x + 4a^2e^2f^2r_x + 4b^2e^2f^2r_x - 8d^2e^2f^2r_x - 4a^2f^4r_x + 4d^4r_x^2 - 8d^2e^2r_x^2 + 4e^4r_x^2 - 8d^2f^2r_x^2 - 8e^2f^2r_x^2 + 4f^4r_x^2 + 4b^2c^2d^2r_y - 4c^4d^2r_y - 4b^4e^2r_y \\
 & + 4b^2c^2e^2r_y + 4a^2b^2f^2r_y + 4a^2c^2f^2r_y - 8b^2c^2f^2r_y + 4c^2d^2f^2r_y + 4b^2e^2f^2r_y - 4a^2f^4r_y - 4b^2d^2r_y^2 + 4c^2d^2r_y^2 + 4b^2e^2r_y^2 - 4c^2e^2r_y^2 - 8a^2f^2r_y^2 + 4b^2f^2r_y^2 + 4c^2f^2r_y^2 + 4d^2f^2r_y^2 + 4e^2f^2r_y^2 - 4f^4r_y^2 + 4b^4r_xr_y - 4b^2c^2r_xr_y + 4c^2d^2r_xr_y + 4b^2e^2r_xr_y - 4c^2e^2r_xr_y - 8a^2f^2r_xr_y^2 + 4b^2f^2r_xr_y^2 + 4c^2f^2r_xr_y^2 + 4d^2f^2r_xr_y^2 + 4e^2f^2r_xr_y^2 - 4f^4r_xr_y^2 + 4b^4r_xr_y^2 - 8b^2c^2r_xr_y^2 + 4c^4r_xr_y^2 - 8b^2f^2r_xr_y^2 - 8c^2f^2r_xr_y^2 + 4f^4r_xr_y^2 + 4a^2c^2d^2r_z - 4c^4d^2r_z + 4a^2b^2e^2r_z \\
 & - 8a^2c^2e^2r_z + 4b^2c^2e^2r_z + 4c^2d^2e^2r_z - 4b^2e^4r_z - 4a^4f^2r_z + 4a^2c^2f^2r_z + 4a^2e^2f^2r_z - 4a^2d^2f^2r_z + 4c^2d^2f^2r_z + 4a^2e^2f^2r_z - 8b^2e^2f^2r_z + 4c^2e^2f^2r_z + 4d^2e^2f^2r_z - 4e^4f^2r_z + 4a^2f^2r_z^2 + 4a^2c^2f^2r_z^2 - 4c^2e^2f^2r_z^2 + 4e^4f^2r_z^2 \\
 & - 4a^2b^2r_x^2r_z + 4a^2c^2r_x^2r_z + 4b^2c^2r_x^2r_z - 4c^4r_x^2r_z - 8c^2e^2r_x^2r_z + 4b^2e^2r_x^2r_z + 4c^2e^2r_x^2r_z + 4a^2f^2r_x^2r_z + 4c^2f^2r_x^2r_z - 4e^2f^2r_x^2r_z - 4a^2d^2r_x^2r_z + 4c^2d^2r_x^2r_z + 4a^2e^2r_x^2r_z - 8b^2e^2r_x^2r_z + 4c^2e^2r_x^2r_z + 4d^2e^2r_x^2r_z - 4e^4r_x^2r_z + 4a^2d^2e^2r_x^2r_z - 4e^4e^2r_x^2r_z \\
 & + 4a^2f^2r_xr_z^2 - 4c^2f^2r_xr_z^2 + 4e^2f^2r_xr_z^2 - 4a^2b^2r_yr_z^2 + 4a^2c^2r_yr_z^2 + 4b^2c^2r_yr_z^2 - 4c^4r_yr_z^2 - 8c^2e^2r_yr_z^2 + 4b^2e^2r_yr_z^2 + 4c^2e^2r_yr_z^2 + 4a^2f^2r_yr_z^2 + 4c^2f^2r_yr_z^2 - 4e^2f^2r_yr_z^2 + 4a^2r_z^2r_w + 4a^2b^2r_z^2r_w + 4a^2c^2d^2r_w + 4b^2c^2d^2r_w - 4c^2d^4r_w + 4a^2b^2e^2r_w - 4b^4e^2r_w + 4b^2d^2e^2r_w - 4a^4f^2r_w + 4a^2b^2f^2r_w + 4a^2d^2f^2r_w + 4a^2e^2f^2r_w + 4b^2d^2f^2r_w - 8b^2e^2f^2r_w + 4a^2f^2r_w^2 + 4a^2c^2f^2r_w^2 - 4a^2e^2f^2r_w^2 + 4b^2d^2f^2r_w^2 - 4d^4f^2r_w^2 - 4a^4r_zr_w^2 + 4a^2b^2r_zr_w^2 + 4a^2c^2r_zr_w^2 \\
 & - 4b^2c^2r_zr_w^2 + 4a^2d^2r_zr_w^2 + 4a^2e^2r_zr_w^2 + 4b^2e^2r_zr_w^2 - 4d^2e^2r_zr_w^2 - 8a^2f^2r_zr_w^2 + 4a^2d^2r_xr_w^2 + 4b^2d^2r_xr_w^2 - 8c^2d^2r_xr_w^2 - 4d^4r_xr_w^2 - 4a^2e^2r_xr_w^2 + 4b^2e^2r_xr_w^2 + 4d^2e^2r_xr_w^2 + 4a^2f^2r_xr_w^2 - 4b^2f^2r_xr_w^2 \\
 & + 4d^2f^2r_xr_w^2 + 4a^2b^2r_yr_w^2 - 4b^4b^2r_yr_w^2 - 4a^2c^2r_yr_w^2 + 4b^2c^2r_yr_w^2 + 4b^2d^2r_yr_w^2 + 4c^2d^2r_yr_w^2 - 8b^2e^2r_yr_w^2 + 4a^2f^2r_yr_w^2 + 4b^2f^2r_yr_w^2 - 4d^2f^2r_yr_w^2 - 4a^4r_zr_w^2 + 4a^2b^2r_zr_w^2 + 4a^2c^2r_zr_w^2 \\
 & - 4b^2c^2r_zr_w^2 + 4a^2d^2r_zr_w^2 + 4c^2d^2r_zr_w^2 + 4a^2e^2r_zr_w^2 + 4b^2e^2r_zr_w^2 - 4d^2e^2r_zr_w^2 - 8a^2f^2r_zr_w^2 + 4a^4r_w^3 - 8a^2b^2r_w^3 + 4b^4r_w^3 - 8a^2c^2r_w^3 - 8b^2d^2r_w^3 + 4d^4r_w^3 \\
 \\
 C = & c^4d^4 - 2b^2c^2d^2e^2 + b^4e^4 - 2a^2c^2d^2f^2 - 2a^2b^2e^2f^2 + a^4f^4 - 2c^2d^4r_x^2 + 2b^2d^2e^2r_x^2 + 2c^2d^2e^2r_y^2 - 2b^2e^4r_x^2 + 2a^2d^2f^2r_x^2 + 2c^2d^2f^2r_x^2 + 2a^2e^2f^2r_x^2 + 2b^2e^2f^2r_x^2 - 4d^2e^2f^2r_x^2 - 2a^2f^4r_x^2 + d^4r_x^4 - 2d^2e^2r_x^4 + e^4r_x^4 \\
 & - 2d^2f^2r_x^4 - 2e^2f^2r_x^4 + f^4r_x^4 + 2b^2c^2d^2r_y^2 - 2c^2d^2e^2r_y^2 - 2b^2e^2r_y^2 + 2b^2c^2e^2r_y^2 + 2a^2b^2f^2r_y^2 + 2a^2c^2f^2r_y^2 - 4b^2c^2f^2r_y^2 + 2c^2d^2f^2r_y^2 + 2b^2e^2f^2r_y^2 - 2a^2f^4r_y^2 - 2b^2d^2r_y^2 + 2c^2d^2r_y^2 + 2b^2e^2r_y^2 - 2c^2e^2r_y^2 - 4a^2f^2r_y^2 \\
 & + 2b^2f^2r_y^2 + 2c^2f^2r_y^2 + 2d^2f^2r_y^2 + 2e^2f^2r_y^2 - 2f^4r_y^2 + b^4r_y^4 - 2b^2c^2r_y^4 + c^4r_y^4 - 2b^2f^2r_y^4 - 2c^2f^2r_y^4 + f^4r_y^4 + 2a^2c^2d^2r_z^2 - 2c^2d^2e^2r_z^2 + 2b^2c^2e^2r_z^2 + 2c^2d^2e^2r_z^2 - 2b^2e^4r_z^2 + 2b^2c^2e^2r_z^2 + 2b^2d^2e^2r_z^2 - 2b^2e^2r_z^4 \\
 & - 2a^2f^2r_z^2 + 2a^2c^2f^2r_z^2 + 2a^2e^2f^2r_z^2 - 2a^2d^2f^2r_z^2 + 2c^2d^2f^2r_z^2 + 2a^2e^2f^2r_z^2 - 2e^2f^2r_z^4 + e^4r_z^4 - 2c^2e^2r_z^4 + e^2f^2r_z^4 + e^4r_z^4 - 2c^2e^2r_z^4 + e^2f^2r_z^4 + e^4r_z^4 - 2b^2e^2r_w^2 + 2b^2d^2e^2r_w^2 - 2a^2f^2r_w^2 + 2a^2b^2f^2r_w^2 + 2a^2d^2f^2r_w^2 + 2a^2e^2f^2r_w^2 + 2b^2d^2f^2r_w^2 - 4c^2d^2r_w^2 - 2d^4r_w^2 - 2a^2e^2r_w^2 + 2b^2e^2r_w^2 + 2d^2e^2r_w^2 + 2a^2f^2r_w^2 + 2b^2f^2r_w^2 - 2d^2f^2r_w^2 - 2a^4r_zr_w^2 + 2a^2b^2r_zr_w^2 + 2a^2c^2r_zr_w^2 - 2b^2c^2r_zr_w^2 + 2d^2e^2r_zr_w^2 \\
 & + 2a^2e^2r_zr_w^2 + 2b^2e^2r_zr_w^2 - 2d^2e^2r_zr_w^2 - 4a^2f^2r_zr_w^2 + a^4r_w^3 - 2a^2b^2r_w^3 + b^4r_w^3 - 2a^2c^2r_w^3 - 2b^2d^2r_w^3 + d^4r_w^3
 \end{aligned}$$

Why the computations are so slow

interval arithmetic + huge formulas → loss of precision

Example: to compute the support sphere radius, we need to solve $Ar^2 + Br + C = 0$

Thanks to dimension reduction:

compute with fixed radii and edge lengths, then "simplify"

$r_x = r_y = r_z = r_w = 1, a = b = c = 2$:

$$A_{1111} = 4(d^2 - e^2)^2 + 4f^4 + ((d^2 - 8)e^2 - 8d^2)f^2$$

$$B_{1111} = 8(d^2 - e^2)^2 + 8f^4 + 2((d^2 - 8)e^2 - 8d^2)f^2$$

$$C_{1111} = d^2e^2f^2$$




1 Introduction

2 Disc packings

3 Sphere packings


4 Conclusion

Techniques




properties of triangulations

Geometry:




... and "tetrahedrizations"



differential geometry

Computer assistance:



case analysis
Python, C++

$$A_{1111} = 4(d^2 - e^2)^2 + 4f^4 + ((d^2 - 8)e^2 - 8d^2)f^2$$


$$B_{1111} = 8(d^2 - e^2)^2 + 8f^4 + 2((d^2 - 8)e^2 - 8d^2)f^2$$

$$C_{1111} = d^2e^2f^2$$

$$8 \left(\begin{array}{l} \arctan \left(\frac{\sqrt{-4(d^2 - e^2)^2 - 4f^4 - ((d^2 - 8)e^2 - 8d^2)f^2}}{(d+2)e^2 + 2d^2 + (d^2 + 4d - 2)f^2} \right) \\ + \arctan \left(\frac{\sqrt{-4(d^2 - e^2)^2 - 4f^4 - ((d^2 - 8)e^2 - 8d^2)f^2}}{(d+2)f^2 - 2d^2 - 2e^2 + (d^2 + 4d)f} \right) \\ + \arctan \left(\frac{\sqrt{-4(d^2 - e^2)^2 - 4f^4 - ((d^2 - 8)e^2 - 8d^2)f^2}}{(d+2)f^2 - 2d^2 - 2e^2 + (d^2 + 4d)f} \right) \\ - \arctan \left(\frac{\sqrt{-4(d^2 - e^2)^2 - 4f^4 - ((d^2 - 8)e^2 - 8d^2)f^2}}{2(d^2 + e^2 + f^2 - 3d)} \right) \end{array} \right)$$

$$\delta_{1111} = \frac{\quad}{\sqrt{-4(d^2 - e^2)^2 - 4f^4 - ((d^2 - 8)e^2 - 8d^2)f^2}}$$

symbolic calculus
SageMath




interval arithmetic
MPFI (RIF SageMath)
Boost (C++)




Open questions: packings and tilings


triangulated packings



~




tilings by triangles
with local rules




density = weighted proportion of tiles

Open questions: packings and tilings


triangulated packings



~



tilings by triangles
with local rules



density = weighted proportion of tiles

Triangulated Packing Problem

algebraic numbers represented by polynomials and intervals

excludes hexagonal packing

Given k disc radii $\overbrace{r_1, \dots, r_k}$, is there a triangulated packing of density $> \frac{\pi}{2\sqrt{3}}$

$\forall r_1, \dots, r_k$ with triangulated packings, one is periodic
(Wang algorithm: search for a period)

\Rightarrow

decidable


$\exists r_1, \dots, r_k$ whose triangulated packings are all aperiodic

\Rightarrow

undecidable?


Open questions: packings and tilings

triangulated packings



~

tilings by triangles
with local rules



density = weighted proportion of tiles

Dense Packing Problem

algebraic numbers represented by polynomials and intervals

Given k disc radii $\overbrace{r_1, \dots, r_k}$, is there a

excludes hexagonal packing

packing of density $> \frac{\pi}{2\sqrt{3}}$

$\forall r_1, \dots, r_k$ with dense packings, one is periodic
(interval arithmetic and subdivision until needed precision)

\Rightarrow

decidable

$\exists r_1, \dots, r_k$ whose dense packings are all aperiodic


\Rightarrow

not possible!

Other “spherical” questions: from pollen grains to kissing problem

Tammes 1930: configuration of pores on a pollen grain

spherical codes




maximize the number of spherical caps of a given radius on a sphere

Other “spherical” questions: from pollen grains to kissing problem

Tammes 1930: configuration of pores on a pollen grain

spherical codes




maximize the number of spherical caps of a given radius on a sphere
place n points on a sphere to maximize the distance between two nearest points

Other “spherical” questions: from pollen grains to kissing problem

Tammes 1930: configuration of pores on a pollen grain

spherical codes



maximize the number of spherical caps of a given radius on a sphere

place n points on a sphere to maximize the distance between two nearest points


find the smallest possible radius of a central sphere tangent to n unit spheres

solved for $n = 3, \dots, 14$, and 24 (1943 – 2015)

Other “spherical” questions: from pollen grains to kissing problem

Tammes 1930: configuration of pores on a pollen grain

spherical codes




maximize the number of spherical caps of a given radius on a sphere

place n points on a sphere to maximize the distance between two nearest points

find the smallest possible radius of a central sphere tangent to n unit spheres

solved for $n = 3, \dots, 14$, and 24 (1943 – 2015)


kissing number: how many unit spheres can touch the same unit central sphere in \mathbb{R}^d
 solved for $d = 2 : 6, 3 :$



Other “spherical” questions: from pollen grains to kissing problem

Tammes 1930: configuration of pores on a pollen grain

spherical codes




maximize the number of spherical caps of a given radius on a sphere

place n points on a sphere to maximize the distance between two nearest points

find the smallest possible radius of a central sphere tangent to n unit spheres

solved for $n = 3, \dots, 14$, and 24 (1943 – 2015)


kissing number: how many unit spheres can touch the same unit central sphere in \mathbb{R}^d
 solved for $d = 2 : 6$, $3 : 12$ (1953), $4 : 24$ (2003), $8 : 240$, $24 : 196560$ (1979)



Other “spherical” questions: from pollen grains to kissing problem

Tammes 1930: configuration of pores on a pollen grain

spherical codes





maximize the number of spherical caps of a given radius on a sphere

place n points on a sphere to maximize the distance between two nearest points

find the smallest possible radius of a central sphere tangent to n unit spheres

solved for $n = 3, \dots, 14$, and 24 (1943 – 2015)

kissing number: how many unit spheres can touch the same unit central sphere in \mathbb{R}^d
 solved for $d = 2 : 6, 3 : 12$ (1953), $4 : 24$ (2003), $8 : 240, 24 : 196560$ (1979)




dodecahedral conjecture

smallest Voronoi cell in sphere packing
 (proved in 2010)


Thank you for your attention!

How to find triangulated packings

packing is triangulated



each disc has a "corona"





How to find triangulated packings

packing is triangulated



each disc has a "corona"



To find disc sizes with triangulated packings, we run through all possible combinations of symbolic coronas of two discs (finite number):

symbolic corona

$$\begin{array}{ccc} r & 1 & \\ r & 1 & r \\ 1 & r & 1 \end{array}$$



(Fernique, Hashemi, Sizova 2019)

How to find triangulated packings

packing is triangulated




each disc has a "corona"



To find disc sizes with triangulated packings, we run through all possible combinations of symbolic coronas of two discs (finite number):

symbolic corona



value of r

$$6 \times \widehat{11r} + 1 \times \widehat{r1r} = 2\pi$$

$$r \approx 0.63$$

(Fernique, Hashemi, Sizova 2019)